Supplementary Table 1 Number of IL-2 doses

mM						mRCC					
Cycle #	N	Mean	Median	Minimum	Maximum	Cycle #	N	Mean	Median	Minimum	Maximum
1	170	9.37	9	3	14	1	191	8.78	9	1	15
2	157	7.83	8	1	13	2	174	6.64	6	1	14
3	60	8.7	9	2	14	3	52	8.35	8	2	14
4	56	7	7	1	14	4	48	5.96	6	1	11
5	24	8.92	9	4	14	5	15	7.73	7	5	14
6	24	5.75	5.5	1	14	6	13	5.23	5	3	8
7	5	7.6	8	4	9	7	5	5.6	5	4	7
8	3	8	8	7	9	8	5	5.2	5	4	7
9	2	9	9	8	10	9	1	5	5	5	5
10	2	8.5	8.5	7	10						
11	1	10	10	10	10						
12	1	9	9	9	9						

Supplementary Table 2 Therapies prior to IL-2

Prior Therapy	mM	mRCC
	(N)	(N)
0.1.011	10	2
Only Chemotherapy	19	3
Only Immunotherapy	44	1
Omy miniunomerapy	44	1
Only Targeted Therapy	0	21
Chemotherapy &	11	1
Immunotherapy		
Chemotherapy & Targeted	0	1
Therapy		
Immunotherapy & Targeted	1	3
minunomerapy & rangeted	1	3
Therapy		
Therapy		
Chemotherapy &	2	0
13		
Imuunotherapy & Targeted		
Therapy		
		20
Total	77	30

Supplementary Table 3 Response to IL-2 by prior therapy for mM

# of Prior Therapy	CR	PR	SD	PD	Missing	Total
	(N)	(N)	(N)	(N)	(N)	(N)
1	3	3	10	38	1	55
2	0	2	6	9	1	18
3 or more	0	0	1	3	0	4


Prior systemic treatments include chemotherapy, immunotherapy (including IFN-a), targeted therapy. It does not include radiation, surgery, and other treatments.

Supplementary Table 4 Response to IL-2 by prior therapy for mRCC

# of Prior	CR	PR	SD	PD	Missing	Total
Therapy	(N)	(N)	(N)	(N)	(N)	(N)
1	1	1	3	13	1	19
2	0	0	2	5	2	9
3 or more	0	0	0	2	0	2

Prior systemic treatments include chemotherapy, immunotherapy (including IFN-a), targeted therapy. It does not include radiation, surgery, and other treatments.

Supplementary Figure 1



Supplementary Figure 1 a Overall survival in mM patients with and without prior systemic therapy. **b** Overall survival in mRCC patients with and without prior systemic therapy. Prior systemic therapy included chemotherapy, targeted therapy, and/or immunotherapy.