Supplemental Online Content

Gyamfi-Bannerman C, Clifton RG, Tita ATN, et al. Eunice Kennedy Shriver Maternal-Fetal Medicine Units Network. Neurodevelopmental outcomes after late preterm antenatal corticosteroids: the ALPS follow-up study. *JAMA*. doi:10.1001/jama.2024.4303

Trial Sites and Study Personnel

- **eTable 1.** Maternal/Neonatal Characteristics for Enrolled vs. Non-Enrolled Follow-Up Participants
- **eeTable 2.** Participants Excluded for Death or Developmental Delay
- **eTable 3.** Comparison of Original Cohorts and Cohorts Remaining at ≥ 6 years
- eTable 4. Subgroup Analysis among Preterm and Term Deliveries

This supplemental material has been provided by the authors to give readers additional information about their work.

Trial Sites and Study Personnel

The members of the *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network that participated in this study are as follows:

Clinical Centers:

Main centers are listed below. Personnel at subsites have the subsite listed in brackets.

University of Alabama at Birmingham, Birmingham, AL – Alan T. N. Tita, MD, PhD, Donna Dunn, PhD, CNM, FNP-BC, Nicole Burrell, MPH, Kristy Domnanovich, PhD, Brian M. Casey, MD

The Ohio State University, Columbus, OH – Kara M. Rood, MD, Anna Bartholomew, MPH, BSN, Maged M. Costantine, MD, RN, CCRP, Renee Devlin, PhD, Aubrey Fowler, BS, Hannah Cackovic Henson, APRN-CNM, Paige Cackovic, BS, Jessica Cohen, BS, Alexis Neri, MPH, Sounali Perez, BS, Kayla McDaniel, BS, Samantha Wiegand, MD [Miami Valley Hospital] David S. McKenna, MD [Miami Valley Hospital], E. Kaye Snow, BSN, RN [Miami Valley Hospital], Kathleen Fennig, MS, BSN, RN [Miami Valley Hospital], Amanda Beeman, PsyD

University of Utah Health Sciences Center, Salt Lake City, UT –Torri D. Metz, MD, MS, Amber Sowles, RN, BSN, Michael W. Varner, MD, M. Sean Esplin, MD [Intermountain Health], Kristina Feldman, PhD, Lauren Perez, PhD, BCBA, Natalie Buerger, PhD, Kelly Stout, PhD, Emily Parker, PhD, Magenta Silberman, MEd, Amanda Nelsen, RN, BSN, Kim Hill, RN, BSN

Brown University, Providence, RI – Dwight J. Rouse, MD, Donna Allard, ADN, RNC, Nina K. Ayala, MD, Victoria Watson, MS, Angelica DeMartino, BSN, MS, Betty Vohr, MD

Columbia University, New York, NY — Cynthia Gyamfi-Bannerman, MD, MS, Sabine Z. Bousleiman, Rn, MSN, MSPH, Ronald J. Wapner, MD, Rupa Ravi, MPH, Annette Zygmunt, PhD, Adisa Velovic, MA, Brandy Firman BSPH, BSDMS, Mitzie Grant, PhD, Matthew Hoffman, MD, MPH [Christiana Care], Carrie A. Kitto [Christiana Care], Elizabeth Higley, PhD [Christiana Care], Kristy T.S. Palomares, MD, PhD [Saint Peter's University Hospital], Imene Beche, MBS [Saint Peter's University Hospital], Maria Ridley, PsyD [Saint Peter's University Hospital], Daniel Skupski, MD [New York-Presbyterian/Queens], Rosalyn Chan-Akeley MD, MPH, [New York-Presbyterian/Queens]

University of Texas Medical Branch, Galveston, TX – George R. Saade, MD, Ashley Salazar, DNP-WHNP, Amelia A. Nounes, RN, MSN, Lisa Thibodeaux, RNC-OB, BSN, Karen E. Smith, PhD

MetroHealth Medical Center-Case Western Reserve University, Cleveland, OH – Kelly Gibson, MD, Eugenia Sweet, RN, Wendy Dalton BSN, RN, Bailey Diaz, BA, Melissa Kinas BSN, RN, David Hackney, MD [University Hospitals], LuAnn Polito BS, JD, AND, Abigail Pierse, BS, Lisa Ramirez, PhD

University of Texas Health Science Center at Houston-Children's Memorial Hermann Hospital, Houston, TX – Suneet P. Chauhan, MD, Hon Dsc, Felecia Ortiz, RN, BSN, Kimberly Rennie, PhD, Dinorah Zanger, PhD

University of North Carolina at Chapel Hill, Chapel Hill, NC – John M. Thorp, Jr., MD, T. Michael O'Shea, MD, MPH, Kelly Clark, BSN, RN, Sally Timlin, MSN, RN, Kathy Lloyd, BS, Carmen Beamon, MD, MPH [WakeMed Health & Hospitals], Hannah Nunn, BA [WakeMed Health & Hospitals], Emily B. Prince, PhD., Adrianne A. Harris, PhD, Jaclyn Starritt, PhD, Laura Kirby, PhD, Maria Barber, MS

Northwestern University, Chicago, IL – William A. Grobman, MD, MBA, Gail Mallett, RN, MS, Elizabeth Rangel, RN, MSN, Beth Plunkett MD [NorthShore University HealthSystem], Kathy Kearns, RN BSN [NorthShore University HealthSystem], Mercedes Ramos-Brinson, BS, Annamarie Russo, MEd, CCRP, Mary Frese, PhD

University of Colorado, Denver, CO – Kent D. Heyborne, MD, Jocelyn Phipers RN BSN, Sarah Schwartz MSN, RN, CNEn, Betsy Burke, RN BSN, Michelle Kleman, MA, LPC CSP [Children's Hospital Colorado], Kellie Roesser, MS, CCRP [Children's Hospital Colorado], Ashly Dalton, MEd, Leah Grande, PhD

Duke University, Durham, NC – Geeta K. Swamy, MD, Jennifer Ferrara, RN, Lena Fried, Brenna L. Hughes, MD, MSc, Kathryn E. Gustafson, PhD, Caitlin Stone

Stanford University, Palo Alto, CA – Yasser Y. El-Sayed, MD, Christine Joyce Lee, RN, Casey Krueger, PhD, Ryan Lucash, PhD, Beth Earhart, PhD, Jessica Patel, PhD, Anna Girsen, MD, PhD

Data Coordinating Center:

The George Washington University Biostatistics Center, Washington, DC – Rebecca G. Clifton, PhD, Elizabeth A. Thom, PhD, Jessica A. de Voest, PhD, MPH, Danielle E. Kincaid, MPH, Crystal Nwachuku, MSHS, Katia J. Barrett, MA, PMP, Pramiksha Marcharchand

<u>NIH</u>:

Eunice Kennedy Shriver National Institute of Child Health and Human Development, Bethesda, MD – Monica Longo, MD, PhD, Uma M. Reddy, MD, MPH, Menachem Miodovnik, MD

Steering Committee Chair:

University of Texas at Austin, Austin, TX - George A. Macones, MD, MSCE

eTable1. Maternal/Neonatal Characteristics for Enrolled vs. Non-Enrolled Follow-Up Participants

Characteristic	Enrolled in Follow-up	Not Enrolled in		
	Study*	Follow-up Study*		
	(N=1,026)	(N=1,805)		
Race/ethnicity [†]				
American Indian / Alaskan Native	2 (0.2%)	5 (0.3%)		
Asian	32 (3.1%)	61 (3.4%)		
Hispanic	309 (30.1%)	544 (30.1%)		
Native Hawaiian / Pacific Islander	3 (0.3%)	6 (0.3%)		
Non-Hispanic Black	258 (25.1%)	470 (26.0%)		
Non-Hispanic White	417 (40.6%)	691 (38.3%)		
More than one self-reported race	2 (0.2%)	5 (0.3%)		
Unknown / Not Reported	3 (0.3%)	23 (1.3%)		
Age, y	29 (6.2)	28 (6.2)		
College degree or higher (maternal)	322/1,005 (32.0%)	406/1,766 (23.0%)		
Private insurance [‡]	444 (43.3%)	616 (34.1%)		
Employed	445/1,024 (43.5%)	703/1,798 (39.1%)		
Number of prior pregnancies >=20 weeks				
No prior pregnancy	417 (40.6%)	720 (39.9%)		

1 prior pregnancy	277 (27.0%)	458 (25.4%)		
>=2 prior pregnancies	332 (32.4%)	627 (34.7%)		
Smoked during pregnancy	108 (10.5%)	282 (15.6%)		
Asthma requiring medication during pregnancy	87 (8.5%)	168 (9.3%)		
Preeclampsia or gestational hypertension	325 (31.7%)	550 (30.5%)		
Gestational diabetes	113 (11.0%)	193 (10.7%)		
Indication for trial entry				
Preterm labor with intact membranes	289 (28.2%)	503 (27.9%)		
Ruptured membranes	227 (22.1%)	393 (21.8%)		
Expected delivery for gestational hypertension	276 (26.9%)	451 (25.0%)		
or preeclampsia				
Expected delivery for fetal growth restriction	30 (2.9%)	60 (3.3%)		
Expected delivery for oligohydramnios	25 (2.4%)	65 (3.6%)		
Expected delivery for other indication	179 (17.4%)	333 (18.4%)		
Treatment Assignment				
Active	522 (50.9%)	907 (50.2%)		
Placebo	504 (49.1%)	898 (49.8%)		
Cesarean delivery	325/1,026 (31.7%)	560/1,801 (31.1%)		
Gestational age at delivery, weeks§	36 (35.3,36.7)	36 (35.4,36.7)		

Small for gestational age (<10th percentile)	169/1,026 (16.5%)	306/1,801 (17.0%)
Major congenital anomaly [¶]	8/1,026 (0.8%)	24/1,801 (1.3%)
Sex		
Female	470/1,026 (45.8%)	861/1,801 (47.7%)
Male	556/1,026 (54.2%)	940/1,801 (52.1%)

^{*}Data are mean (standard deviation), median (Q1,Q3), or n (%) unless otherwise specified.

eTable2. Participants Excluded for Death or Developmental Delay

Exclusion	Betamethasone	Placebo	P-value
	(N=1,429)	(N=1,402)	
Infant/Neonatal Death	6 (0.42%)	4 (0.29%)	0.75
Childhood Death	6 (0.42%)	4 (0.29%)	0.75
Maternal Death	6 (0.42%)	4 (0.29%)	0.75
Severe Autism/Developmental Delay	12 (0.84%)	13 (0.93%)	0.80

[†]Maternal race/ethnicity was based on self-report.

[‡]Those without private insurance have government-assisted insurance or are self-pay/uninsured.

[§]Gestational age at delivery missing in 4.

[¶]Although the presence of a major congenital anomaly was an exclusion criterion in the ALPS trial, these disorders were not discovered until birth.

eTable3. Comparison of Original Cohorts and Cohorts Remaining at ≥ 6 years

Characteristic	Original Cohort (N=2,831)			Unweighted Analysis Cohort (N=1,026)			Weighted Analysis Cohort (N=2,832)*		
	Betamethasone (N=1,429)	Placebo (N=1,402)	Standardized Mean Difference	Betamethasone (N=522)	Placebo (N=504)	Standardized Mean Difference	Betamethasone (N=1,430)	Placebo (N=1,402)	Standardized Mean Difference
Race/ethnicity			0.116			0.250			0.258
American Indian / Alaskan Native	3 (0.2%)	4 (0.3%)		2 (0.4%)	0 (0%)		5 (0.3%)	0 (0%)	
Asian	56 (3.9%)	37 (2.6%)		21 (4.0%)	11 (2.2%)		54 (3.7%)	27 (1.9%)	
Hispanic	405 (28.3%)	448 (32.0%)		147 (28.2%)	162 (32.1%)		411 (28.7%)	470 (33.5%)	
Native Hawaiian / Pacific Islander	3 (0.2%)	4 (0.3%)		0 (0%)	2 (0.4%)		0 (0%)	7 (0.5%)	
Non-Hispanic Black	6 (0.4%)	3 (0.2%)		3 (0.6%)	0 (%)		8 (0.5%)	0 (0%)	
Non-Hispanic White	363 (25.4%)	365 (26.0%)		124 (23.8%)	134 (26.6%)		370 (25.9%)	404 (28.8%)	
More than one self-reported race	580 (40.6%)	528 (37.7%)		222 (42.5%)	195 (38.7%)		575 (40.2%)	496 (35.4%)	
Unknown / Not Reported	13 (0.9%)	13 (0.9%)		3 (0.6%)	0 (0%)		8 (0.5%)	(0%)	
Age, y	29 (6.3)	28 (6.1)	0.127	29 (6.3)	29 (6.1)	0.065	29 (6.3)	28 (6.1)	0.126
College degree or higher (maternal)	381 (27.3%)	347 (25.2%)	0.047	164 (32.0%)	158 (32.1%)	0.003	403 (28.7%)	372 (27.2%)	0.033
Private insurance	558 (39.0%)	502 (35.8%)	0.067	228 (43.7%)	216 (42.9%)	0.017	558 (39.0%)	504 (35.9%)	0.064
Employed	577 (40.5%)	571 (40.8%)	0.005	217 (41.7%)	228 (45.3%)	0.074	564 (39.5%)	581 (41.5%)	0.041
Number of prior pregnancies >=20 weeks			0.061			0.118			0.130
No prior pregnancy	565 (39.5%)	572 (40.8%)		209 (40.0%)	208 (41.3%)		567 (39.7%)	574 (40.9%)	

1 prior pregnancy	360 (25.2%)	375 (26.7%)		131 (25.1%)	146 (29.0%)		357 (25.0%)	411 (29.3%)	
>=2 prior pregnancies	504 (35.3%)	455 (32.5%)		182 (34.9%)	150 (29.8%)		506 (35.4%)	418 (29.8%)	
Smoked during pregnancy	204 (14.3%)	186 (13.3%)	0.029	55 (10.5%)	53 (10.5%)	0.001	205 (14.3%)	189 (13.4%)	0.026
Asthma requiring medication during pregnancy	129 (9.0%)	126 (9.0%)	0.001	50 (9.6%)	37 (7.3%)	0.080	141 (9.8%)	106 (7.5%)	0.081
Preeclampsia or gestational hypertension	435 (30.4%)	440 (31.4%)	0.020	168 (32.2%)	157 (31.2%)	0.022	463 (32.4%)	422 (30.1%)	0.049
Gestational diabetes	153 (10.7%)	153 (10.9%)	0.007	54 (10.3%)	59 (11.7%)	0.043	149 (10.4%)	160 (11.4%)	0.033
Indication for trial entry			0.041			0.142			0.163
Preterm labor with intact membranes	400 (28.0%)	392 (28.0%)		143 (27.4%)	146 (29.0%)		405 (28.3%)	435 (31.0%)	
Ruptured membranes	316 (22.1%)	304 (21.7%)		119 (22.8%)	108 (21.4%)		319 (22.3%)	291 (20.7%)	
Expected delivery for gestational hypertension or preeclampsia	356 (24.9%)	371 (26.5%)		150 (28.7%)	126 (25.0%)		416 (29.1%)	338 (24.1%)	
Expected delivery for fetal growth restriction	46 (3.2%)	44 (3.1%)		13 (2.5%)	17 (3.4%)		35 (2.4%)	51 (3.6%)	
Expected delivery for oligohydramnios	48 (3.4%)	42 (3.0%)		15 (2.9%)	10 (2.0%)		39 (2.7%)	28 (2.0%)	
Expected delivery for other indication	263 (18.4%)	249 (17.8%)		82 (15.7%)	97 (19.2%)		218 (15.2%)	261 (18.6%)	
Cesarean delivery	454 (31.8%)	431 (30.8%)	0.022	164 (31.4%)	161 (31.9%)	0.011	447 (31.3%)	432 (30.8%)	0.010
Gestational age at delivery, weeks	36.1 (1.1)	36.1 (1.2)	0.006	36.1 (1.1)	36.1 (1.2)	0.015	36.1 (1.1)	36.1 (1.2)	0.001
Small for gestational age (<10th percentile)	255 (17.9%)	220 (15.7%)	0.058	90 (17.2%)	79 (15.7%)	0.042	248 (17.3%)	233 (16.6%)	0.020
Major congenital anomaly†	11 (0.8%)	21 (1.5%)	0.069	1 (0.2%)	7 (1.4%)	0.136	3 (0.2%)	20 (1.4%)	0.134

Sex			0.051			0.048			0.057
Female	690 (48.4%)	641 (45.8%)		233 (44.6%)	237 (47.0%)		632 (44.2%)	660 (47.1%)	
Male	737 (51.6%)	759 (54.2%)		289 (55.4%)	267 (53.0%)		798 (55.8%)	743 (52.9%)	

^{*}Accounts for maternal age, insurance status, and smoking during pregnancy. N's may not total original cohort because of rounding.

[†]Although the presence of a major congenital anomaly was an exclusion criterion in the ALPS trial, these disorders were not discovered until birth.

eTable4. Subgroup Analysis among Preterm and Term Deliveries

Outcome	Betamethasone (N=479)	Placebo (N=470)	Unadjusted*	Adjusted* [†]	
Primary Outcome					
DAS-II General Conceptual Ability < 85 [‡]			Relative Risk	Relative Risk	
Preterm	67/398 (16.8%)	67/391 (17.1%)	0.98 (0.72,1.34)	0.98 (0.73,1.30)	
Term	15/81 (18.5%)	20/79 (25.3%)	0.73 (0.40,1.33)	0.83 (0.49,1.42)	
Secondary Outcomes	(N=522)	(N=504)			
Social Responsiveness Scale t-score > 65§			Relative Risk	Relative Risk	
Preterm	61/432 (14.1%)	62/421 (14.7%)	0.96 (0.69,1.33)	0.93 (0.67,1.28)	
Term	9 (10.5%)	10 (12.1%)	0.87 (0.37,2.03)	0.90 (0.39,2.08)	
Gross Motor Function Classification System > Level I [¶]					
Preterm	1 (0.2%)	1 (0.2%)			
Term	0	0			
Child Behavior Checklist [#]					
Total problems t-score**			Mean Difference	Mean Difference	
Preterm	48.7 (11.2)	48.7 (11.3)	-0.05 (-1.54,1.44)	-0.38 (-1.85,1.09)	
Term	47.5 (9.9)	50.8 (10.9)	-3.28 (-6.63,0.06)	-3.25 (-6.54,0.04)	

^{*}Data are n (%) or relative risk (95% CI) for categorical outcomes and mean (standard deviation) or mean difference (95% CI) for continuous outcomes.

*Child Behavior Checklist total problems t-score ranges: < 60 normal, 60-63 borderline (84th-90th percentile), and ≥ 64 clinical (> 90th percentile) with higher scores indicating greater problems.

**Child Behavior Checklist missing in 3.

[†]Adjusted for maternal age, child age, sex, maternal education, and center.

 $^{^{\}dagger}$ DAS-II denotes Differential Ability Scales 2nd Edition. The DAS-II General Conceptual Ability score has a mean of 100 and a standard deviation of 15; ≥ 85 average and < 85 below average with higher scores indicating greater abilities.

[§]Social Responsiveness Scale ranges: < 60 normal, 60-65 mild, 66-75 moderate, and ≥ 76 severe with higher scores indicating greater problems.

[¶]The Gross Motor Function Classification System has 5 levels and increasing level indicates greater functional impairment: walks without limitations, walks with limitations, walks with hand-held mobility device, self-mobility with limitations; may use powered mobility, and transported in a manual wheelchair.