

Supporting Information

Metabolic profiling in plasma and brain induced by 17 β -estradiol supplementation in ovariectomized mice

So Hwi Yang ^{a,1}, Ye Jin Kim ^{a,1}, Hye Rim Yang ^a, Sang Un Park ^b, Jae Geun kim ^{a,*}, and Jae Kwang Kim^{a,c,*}

^aDivision of Life Sciences, College of Life Sciences, and Bioengineering, Incheon National University, Incheon 22012, Republic of Korea


^b Department of Crop Science, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon 34134, Republic of Korea

^c Convergence Research Center for Insect Vectors, Incheon National University, Incheon 22012, Republic of Korea

*Corresponding author Division of Life Sciences, College of Life Sciences and Bioengineering, Incheon National University, Incheon 22012, Republic of Korea

E-mail address kjkpj@inu.ac.kr (J. K. Kim), thgnltkfd12@inu.ac.kr (S. H. Yang), 201721047@inu.ac.kr (Y. J. Kim), jgkim@inu.ac.kr (J. G. Kim), hr.yang0414@gmail.com (H. R. Yang), supark@cnu.ac.kr (S. U. Park)

(A) Plasma



(B) Brain




Figure S1. Pathway and enrichment analyses of (A) plasma and (B) brain metabolites affected by 17β -estradiol supplementation in the OVX mouse model. Pathway analysis was conducted using KEGG human metabolic pathways.