

Supplementary table 2.**Summary of healthy blood donors and their involvement in individual experiments**

Donor ID	Gender	Age	Fig.2a	Fig.2b	Fig.2c	Fig.2d	Fig.3a	Fig.3b	Fig.3c	Fig.3d	Fig.4	Fig.7
1	female	36	x	x			x	x	x			
2	female	36		x			x	x	x	x		x
3	female	26			x							
4	female	30					x	x				
5	female	42			x							
6	female	27	x									
7	female	44									x	x
8	female	48	x								x	
9	female	26									x	
10	female	28										x
11	female	33									x	
12	female	32										x
13	female	38									x	
14	female	32										x
15	female	39			x		x	x				
16	female	32					x	x				
17	female	46					x	x			x	
18	female	21			x							

19	female	45			x							
20	male	30		x			x	x	x	x		
21	male	42	x	x	x		x	x	x	x		
22	male	37		x	x		x	x	x	x		x
23	male	27	x	x								x
24	male	28	x									x

Melanoma-associated fibroblasts impair CD8+ T cell function and modify expression of immune checkpoint regulators via increased arginase activity

Barbara Érsek^{1,2*}, Pálma Silló^{3*}, Ugur Cakir³, Viktor Molnár⁴, András Bencsik¹, Balázs Mayer³, Eva Mezey⁵, Sarolta Kárpáti³, Zoltán Pósi^{1§} and Krisztián Németh^{3§}

Corresponding author:

Zoltán Pósi

Department of Genetics, Cell and Immunobiology, Semmelweis University

4 Nagyvarad ter, VII/709, Budapest, H-1089, Hungary

Phone: +36-1-210-2930 Ext. 56435; Fax: +36-1-303-6968

E-mail: pos.zoltan@med.semmelweis-univ.hu