

SUPPLEMENTARY MATERIAL

TABLE 5. A 3 x 3 paired sample contingency table illustrating the proportion of CKD cats with differing classification of cortico-medullary definition in left and right kidneys between baseline and repeated ultrasound scans.

				<i>Repeated</i>			n	<i>P</i> -value
				Maintained	Poorly maintained	Loss		
Cortico-medullary definition (left)	<i>Baseline</i>	All	Maintained	1 (4%)	1 (4%)	0 (0%)	27	.11
			Poorly maintained	5 (19%)	9 (33%)	1 (4%)		
			Loss	0 (0%)	2 (7%)	8 (30%)		
	<i>Baseline</i>	Normocalcemia	Maintained	0 (0%)	0 (0%)	0 (0%)	14	.23
			Poorly maintained	3 (21%)	5 (36%)	1 (7%)		
			Loss	0 (0%)	1 (7%)	4 (29%)		
	<i>Baseline</i>	Hypercalcemia	Maintained	1 (8%)	1 (8%)	0 (0%)	13	.42
			Poorly maintained	2 (15%)	4 (31%)	0 (0%)		
			Loss	0 (0%)	1 (8%)	4 (31%)		
Cortico-medullary definition (right)	<i>Baseline</i>	All	Maintained	0 (0%)	2 (8%)	0 (0%)	26	.43
			Poorly maintained	5 (19%)	8 (31%)	3 (12%)		
			Loss	0 (0%)	3 (12%)	5 (19%)		
	<i>Baseline</i>	Normocalcemia	Maintained	0 (0%)	0 (0%)	0 (0%)	13	.07
			Poorly maintained	3 (23%)	6 (46%)	0 (0%)		
			Loss	0 (0%)	1 (8%)	3 (23%)		
	<i>Baseline</i>	Hypercalcemia	Maintained	0 (0%)	2 (15%)	0 (0%)	13	.79
			Poorly maintained	2 (15%)	2 (15%)	3 (23%)		
			Loss	0 (0%)	2 (15%)	2 (15%)		

Abbreviation: n, number of cats.