

SUPPLEMENTARY MATERIAL

TABLE 6. A 3 x 3 paired sample contingency table illustrating the proportion of CKD cats with differing classification of nephrocalcinosis in left and right kidneys between baseline and repeated ultrasound scans.

				<i>Repeated</i>			n	<i>P</i> -value
				Absent	Present (suspected)	Present		
Nephrocalcinosis (left)	<i>Baseline</i>	All	Absent	11 (41%)	2 (7%)	0 (0%)	27	.59
			Present (suspected)	0 (0%)	3 (11%)	3 (11%)		
			Present	1 (4%)	1 (4%)	6		
		Normocalcemia	Absent	8 (57%)	1 (7%)	0 (0%)	14	1
			Present (suspected)	0 (0%)	1 (7%)	1 (7%)		
			Present	1 (7%)	0 (0%)	2 (14%)		
		Hypercalcemia	Absent	3 (23%)	1 (8%)	0 (0%)	13	.42
			Present (suspected)	0 (0%)	2 (15%)	2 (15%)		
			Present	0 (0%)	1 (8%)	4 (31%)		
Nephrocalcinosis (right)	<i>Baseline</i>	All	Absent	13 (50%)	4 (15%)	1 (4%)	26	.1
			Present (suspected)	1 (4%)	3 (12%)	2 (8%)		
			Present	0 (0%)	1 (4%)	1 (4%)		
		Normocalcemia	Absent	9 (69%)	1 (8%)	1 (8%)	13	.37
			Present (suspected)	0 (0%)	1 (8%)	0 (0%)		
			Present	0 (0%)	0 (0%)	1 (8%)		
		Hypercalcemia	Absent	4 (31%)	3 (23%)	0 (0%)	13	.3
			Present (suspected)	1 (8%)	2 (15%)	2 (15%)		
			Present	0 (0%)	1 (8%)	0 (0%)		

Abbreviation: n, number of cats.