

Supplementary Table 1 Characteristics of GC patients

Variables		All patients (n=117)	bsPD-L1-negative (n=100)	bsPD-L1-positive (n=17)	p value
Age	Median (range)	75 (37 - 93)	75 (37-93)	77 (58-88)	0.963
Gender					0.234
	Male	83 (70.9%)	73 (73.0%)	10 (58.8%)	
	Female	34 (29.1%)	27 (27.0%)	7 (41.2%)	
Smoking					0.831
	Never	44 (37.6%)	38 (38.0%)	6 (35.3%)	
	Current or Former	73 (62.4%)	62 (62.0%)	11 (64.7%)	
Histology					0.656
	Differentiated	63 (53.9%)	53 (53.0%)	10 (58.8%)	
	Undifferentiated	54 (46.1%)	47 (47.0%)	7 (41.2%)	
T					0.484
	-1	36 (30.8%)	32 (32.0%)	4 (23.5%)	
	2-	81 (69.2%)	68 (68.0%)	13 (76.5%)	
N					0.262
	0	61 (52.1%)	50 (50.0%)	11 (64.7%)	
	1-	56 (47.9%)	50 (50.0%)	6 (35.3%)	
M					0.520
	0	105 (89.7%)	89 (89.0%)	16 (94.1%)	
	1	12 (10.3%)	11 (11.0%)	1 (5.9%)	
Disease stage					0.713
	I - II	71 (60.7%)	60 (60.0%)	11 (64.7%)	
	III - IV	46 (39.3%)	40 (40.0%)	6 (35.3%)	