



Supporting Information

for

Cholesterol nanoarchaeosomes for alendronate targeted delivery as an anti-endothelial dysfunction agent

Horacio Emanuel Jerez, Yamila Roxana Simioni, Kajal Ghosal, Maria Jose Morilla and Eder Lilia Romero

Beilstein J. Nanotechnol. **2024**, *15*, 517–534. [doi:10.3762/bjnano.15.46](https://doi.org/10.3762/bjnano.15.46)

Additional figures




Figure S1: Cytokine release in the compartments of the mild inflammation model before (pre) and after (post) 24 h of crosstalk incubation: (A) apical (HUVECs) and (B) basolateral (THP-1 macrophages). Values of IL-6 and TNF- α are shown on the left Y axis, and values of IL-8 are shown on the right Y axis. Data are expressed as mean \pm SD ($n = 2$).




Figure S2: Cytokine release in the compartments of the pronounced inflammation model before (pre) and after (post) 24 h of cross-talk incubation: (A) apical (HUVECs) and (B) basolateral (FCs). Values of IL-6 and TNF- α are shown on the left Y axis, and values of IL-8 are shown on the right Y axis. Data are expressed as mean \pm SD ($n = 2$).




Figure S3: Cytokine release in the upper compartment of the mild inflammation model after incubation with void vesicles. Values of IL-6 and TNF- α are shown on the left Y axis, and values of IL-8 are shown on the right Y axis. Data are expressed as mean \pm SD ($n = 2$).