

SUPPLEMENTARY DIGITAL MATERIAL 1

Supplementary Table I.—Lymphocytes levels in patients with post-polio myelitis syndrome (PPS) and controls.

Variable	PPS patients (N.=47)	Controls (N.=27)	P value
T lymphocytes (%)	72 (64; 78)	68 (63; 78)	0.66
T lymphocytes (N/mm ³)	1294 (892; 1848)	1481 (1148; 1739)	0.46
Th1 cells (%)	35 (29; 43)	35 (31; 38)	0.43
Th1 cells (N/mm ³)	403 (257; 558)	365 (274; 622)	0.88
Th2 cells (%)	11 (9; 14)	13 (10; 15)	0.32
Th2 cells (N/mm ³)	123 (98; 168)	130 (99; 200)	0.44
Th17 cells (%)	16 (13; 20)	17 (12; 20)	0.98
Th17 cells (N/mm ³)	172 (112; 250)	182 (121; 249)	0.86
T regulatory (%)	7 (6; 9)	7 (6; 8)	0.45
T regulatory (N/mm ³)	66 (49; 83)	65 (47; 86)	0.89
B lymphocytes (%)	15 (13; 20)	15 (12; 18)	0.87
B lymphocytes (N/mm ³)	283 (200; 384)	308 (212; 473)	0.26
NK lymphocytes (%)	13 (8; 18)	14 (10; 19)	0.61
NK lymphocytes (N/mm ³)	225 (140; 327)	309 (147; 380)	0.20

Values are medians (interquartile range). Alpha risk: 0.002.