

SUPPLEMENTARY DIGITAL MATERIAL 2

Supplementary Table S2: II.—Cytokines levels in patients with post-polio myelitis syndrome (PPS) and controls.

Variable	PPS patients (N.=47)	Controls (N.=27)	P value
GM-CSF	0.20 (0.06; 0.28)	0.16 (0.05; 0.28)	0.52
IFN- $\gamma$	294 (11; 47)	27 (2; 48)	0.69
IL-1 $\beta$	0.65 (0.37; 1.16)	0.72 (0.27; 1.28)	0.89
IL-4	0.30 (0.23; 0.38)	0.29 (0.19; 0.43)	0.88
IL-6	0.48 (0.08; 1.41)	0.53 (0.31; 1.62)	0.42
IL-8	8·1 (6.1; 12.3)	7·5 (5.0; 14.3)	0.36
IL-10	0.32 (0.14; 0.56)	0.29 (0.06; 0.61)	0.71
IL-13	2·9 (1.7; 8.6)	1·7 (1.7; 9.2)	0.40
IL-17	1·7 (0.02; 4.1)	1·5 (0.02; 3.7)	0.72
IL-21	5·5 (1.3; 12.9)	3·7 (0.3; 22.8)	0.69
IL-22	0.9 (0.3; 1.9)	0.7 (0.3; 1.2)	0.51
IL-23	6·9 (4.7; 10.5)	8·1 (5.7; 12.1)	0.31
TGF- $\beta$ 1	4252 (2867; 5869)	3919 (3192; 5269)	0.87
MIP-3a	49 (26; 88)	57 (35; 190)	0.13
MCP-1	240 (159; 289)	203 (144; 226)	0.02
TNF- $\alpha$	1·4 (0.6; 1.9)	1·2 (0.4; 2.8)	0.62
RANTES	40,654 (28,043; 65,985)	46,372 (29,599; 65,748)	0.38

All values are expressed in pg/mL as medians (interquartile range). Alpha risk: 0.002.