

Type 1 Diabetes Human Enteroid Studies Reveal Major Changes In The Intestinal Epithelial Compartment

Vishwesh Bharadiya¹, Yan Rong¹, Zixin Zhang¹, Ruxian Lin¹, Anthony Lawrence Guerrero², C. Ming Tse¹, Mark Donowitz^{1,3}, and Varsha Singh^{1, #}

¹Divisions of Gastroenterology & Hepatology, Department of Medicine, ²Department of Pediatrics, ³Department of Physiology, the Johns Hopkins University School of Medicine, Baltimore, Maryland, 21205, USA

#correspondance: Varsha Singh, Divisions of Gastroenterology & Hepatology, Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, 21205, USA

email: vsingh11@jhmi.edu




Fig S1. T1D enteroids retain pathological changes after multiple passaging

a. qRT-PCR analysis in duodenal enteroids from lower passage (p8) and higher passage (p30) showing differences between HS and T1D. T1D enteroids retain pathological differences in signature genes tested between the two groups. Results are shown as mean \pm SEM. * $P < 0.05$ vs control. Analysis of differences was determined by Student's unpaired t-tests. N=3 independent experiment was performed on enteroids from 1 donor from each group.

Fig S2. The original images of blots.

Figure 3f

