


Figure S1: Comparison of kidney function at follow-up based on race/APOL1 status of the participants using the race inclusive CKD-EPI 2012 creatinine-cystatin C GFR calculator



**Table S1:** Comparison of baseline characteristics of participants with high-risk APOL1 genotype who did and did not have follow-up data on kidney function at 25 years

	With follow-up (N= 45)	Without follow-up (N=65)	p-value
Age, years	50±5	51±5	0.3
Women	27 (60%)	39 (60%)	>1.0
Body mass index, kg/m <sup>2</sup>	27±3	26±4	0.2
Systolic blood pressure, mmHg	118±11	116±12	0.35
Diastolic blood pressure, mmHg	75±8	74±8	0.5
eGFR-creatinine, mL/min/1.73m <sup>2</sup>	104±11	104±9	1.0
College graduate	20 (44%)	22 (34%)	0.3
Employed full- or part-time	39 (87%)	51 (78%)	0.3
Current smoker	9 (20%)	31 (48%)	0.01
Family history of diabetes	5 (11%)	9 (14%)	0.7
Family history of hypertension	11 (24%)	15 (23%)	0.9
Established care with a primary care physician	41 (91%)	49 (75%)	0.03

Data represented as mean±standard deviation

The above data was obtained via publicly available ARIC dataset (not USRDS)