

Local recurrence

Group by BD type	Study name	Statistics for each study			Event rate and 95% CI					
		Event rate	Lower limit	Upper limit						Relative weight
С	lacopini, 2015	0.050	0.003	0.475	- -	-	—1	- 1	- 1	7.79
	Kasuga, 2021	0.045	0.003	0.448	-	_	_			7.83
	Kinoshita, 2019	0.020	0.001	0.251	-	_		- 1		8.04
	Matsumoto, 2019	0.042	0.003	0.425	-	_	_	- 1		7.86
	Manta, 2021	0.009	0.001	0.134	-			- 1		8.12
;	Matsui, 2021	0.029	0.002	0.336	-					7.96
;	Nishio, 2020	0.013	0.001	0.175	-	-				8.09
:	Suzuki, 2017	0.031	0.004	0.191	-	– I				15.89
;	Yang DH, 2019	0.133	0.034	0.405	-	-	-			28.43
	Pooled	0.043	0.020	0.090	•					
	Prediction Interval									
+CD	Kochhar, 2018	0.071	0.004	0.577	-	_				24.03
+CD	Lightner, 2021	0.020	0.001	0.251	_	_				25.36
+CD	Ngamruengphong, 2022	0.022	0.003	0.142		.		- 1		50.61
C+CD	Pooled	0.029	0.007	0.108		- 1		- 1		
C+CD	Prediction Interval									
/erall	Pooled	0.039	0.020	0.075	•	- 1		- 1		
/erall	Prediction Interval									
					0.00	0.25	0.50	0.75	1.00	

overall I2 = 0%UC = 0%UC+CD = 0%

Supplementary Fig. 5. Forest plot for local recurrence. CI, confidence interval; UC, ulcerative colitis; CD, Crohn's disease.