

Supplementary material has been published as submitted. It has not been copyedited, or typeset by Acta Dermato-Venereologica

Table SI. Detailed version of characteristics of children 0-17 years with atopic dermatitis, food allergy, allergic rhinoconjunctivitis, and/or asthma

	All participants	Practicing specialist % (n/N of total)				Hospital % (n/N of total)		
		Total	Pediatrician	Dermatologist	Otorhino- laryngologist	Total	Pediatric department	Dermatology department
Gender, boys	63 % (175/277)	61 % (68/112)	58 % (58/100)	71 % (5/7)	100 % (5/5)	65 % (107/165)	65 % (79/122)	65 % (28/43)
Age, years, median (95% CI)	10 (9-10)	10 (9-12)	10 (9-12)	0 (0-2)	13 (-)	9 (8-10)	10 (9-11)	4 (3-9)
Parents living together, Yes	82 % (227/277)	83 % (93/112)	81 % (82/101)	100 % (6/6)	100 % (5/5)	81 % (134/165)	79 % (96/121)	86 % (38/44)
Household income								
< 400.000 DKK/year	5 % (15/277)	5 % (5/111)	5 % (5/100)	0 % (0/6)	0 % (0/5)	6 % (10/166)	4 % (5/122)	11 % (5/44)
400.000-800.000 DKK/year	34 % (93/277)	34 % (38/111)	34 % (34/100)	50 % (3/6)	20 % (1/5)	33 % (55/166)	30 % (37/122)	41 % (18/44)
> 800.000 DKK/year	48 % (134/277)	49 % (54/111)	48 % (48/100)	50 % (3/6)	60 % (3/5)	48 % (80/166)	51 % (62/122)	41 % (18/44)
Do not know/ do not want to disclose	13 % (35/277)	13 % (14/111)	13 % (13/100)	0 % (0/6)	20 % (1/5)	13 % (21/166)	15 % (18/122)	7 % (3/44)
Parents educational level								
Mothers								
- Primary school	1 % (3/277)	0 % (0/112)	0 % (0/101)	0 % (0/6)	0 % (0/5)	2 % (3/165)	2 % (2/122)	2 % (1/43)
- High school and vocational school	9 % (24/277)	11 % (12/112)	10 % (10/101)	0 % (0/6)	40 % (2/5)	7 % (12/165)	7 % (8/122)	9 % (4/43)
- Short and medium (bachelor) cycle higher education	47 % (129/277)	50 % (56/112)	51 % (52/101)	17 % (1/6)	60 % (3/5)	44 % (73/165)	45 % (55/122)	42 % (18/43)
- Long cycle higher (master) education	40 % (112/277)	37 % (41/112)	36 % (36/101)	83 % (5/6)	0 % (0/5)	43 % (71/165)	42 % (51/122)	47 % (20/43)
Fathers								
- Primary school	3 % (9/272)	3 % (3/110)	2 % (2/99)	0 % (0/6)	20 % (1/5)	4 % (6/162)	3 % (4/120)	5 % (2/42)

- High school and vocational school	22 % (60/272)	29 % (32/110)	30 % (30/99)	0 % (0/6)	40 % (2/5)	17 % (28/162)	14 % (17/120)	26 % (11/42)
- Short and medium (bachelor) cycle higher education	27 % (74/272)	27 % (30/110)	28 % (28/99)	33 % (2/6)	0 % (0/5)	27 % (44/162)	28 % (33/120)	26 % (11/42)
- Long cycle higher (master) education	44 % (120/272)	37 % (41/110)	35 % (35/99)	67 % (4/6)	40 % (2/5)	49 % (79/162)	54 % (65/120)	33 % (14/42)
Family history of atopic diseases								
- AD	36 % (98/274)	32 % (36/110)	32 % (31/98)	57 % (4/7)	20 % (1/5)	38 % (62/164)	31 % (38/122)	57 % (24/42)
- Food allergy	29 % (79/274)	28 % (31/110)	31 % (30/98)	0 % (0/7)	20 % (1/5)	29 % (48/164)	26 % (32/122)	38 % (16/42)
- ARC	62 % (169/274)	60 % (66/110)	60 % (59/98)	57 % (4/7)	60 % (3/5)	63 % (103/164)	62 % (76/122)	64 % (27/42)
- Asthma	34 % (94/274)	34 % (38/110)	37 % (36/98)	14 % (1/7)	20 % (1/5)	34 % (56/164)	34 % (42/122)	33 % (14/42)
- None	18 % (49/274)	17 % (19/110)	17 % (17/98)	29 % (2/7)	0 % (0/5)	18 % (30/164)	21 % (26/122)	9 % (4/42)

Data are summarized as percent (number of total) or median (95% confidence interval)

AD: atopic dermatitis; ARC: allergic rhinoconjunctivitis; CI: confidence interval; n: number.