

**Supplemental Table 3. Unstandardized Parameter Estimates and R-square Results for the Selected SEM Structural Model**

	Estimate	S.E.	Sig.	R-square	Sig
<b>Drug Acquisition Risk by:</b>					
Used with new people	1.000	0.000	-	0.837	***
Used in a new location	0.921	0.072	***	0.711	***
Had new source/supplier	0.895	0.067	***	0.670	***
<b>Drug Use Risk by:</b>					
Used different amounts	1.000	0.000	-	0.435	***
Used alone	0.750	0.182	***	0.245	*
Used multiple drugs	1.151	0.163	***	0.577	***
Used with benzodiazepines	0.862	0.162	***	0.324	**
Used with alcohol	0.832	0.142	***	0.301	***
Used in a different way	0.840	0.168	***	0.307	***
First use in a while	0.487	0.157	***	0.103	NS
Fentanyl preference	0.730	0.174	***	0.232	*
<b>Psychosocial Risk by:</b>					
Trauma exposure	1.000	0.000	-	0.266	*
Homelessness	0.865	0.268	***	0.199	**
Serious psychological distress	1.075	0.299	***	0.307	*
<b>Drug Acquisition Risk on Psychosocial Risk</b>	1.212	0.348	***	-	
<b>Drug Use Risk on Psychosocial Risk</b>	0.726	0.24	**	-	
<b>Past-Year Overdose on:</b>					
Drug acquisition risk	0.114	0.155	NS	-	
Drug use risk	0.434	0.218	*	-	
<b>Drug Acquisition Risk with Drug Use Risk</b>	0.153	0.079	NS	-	
<b>Total Indirect Effect of Psychosocial Risk on Past-Year Overdose</b>	0.454	0.176	**	-	
<b>Any Past-Year Overdose</b>	-	-	-	0.131	*

**Note.** Figures shown are unstandardized parameter estimates, standard errors, and the r-square statistic based on the final SEM structural model.

- = Not estimated; NS = Non-significant; \* = < .05; \*\* = < .01; \*\*\* = < .001.