


# Sexual health and testosterone concentration in male lymphoma survivors: A systematic review

Supplementary data provided by Behringer et al.

Title: Sexual quality of life in Hodgkin Lymphoma: A longitudinal analysis by the German Hodgkin Study Group

Behringer, Karolin  
Müller, H  
Flechtner, H.-H.  
Brillant, C  
Halbsguth, T V  
Thielen, I  
Eichenauer, D. A.  
Schober, T  
Nisters-Backes, H  
Fuchs, M  
Engert, A  
Borchmann, P

## Supplementary data for adult men with Hodgkin lymphoma:



HM, 28.11.2022

Mean values of sexual functioning in males of the HD10 trial (normal reference level= 83)

time	Mean	N	Std.-deviation
1	76,5636	485	29,71529
2	58,5594	344	33,39190
3	70,2827	338	32,61789
4	77,3558	158	28,89504
5	75,2739	142	30,39000
6	79,6296	120	27,22926
7	83,9506	108	23,64761
8	80,8279	102	27,84510
9	79,9458	82	30,16184

Mean values of sexual functioning in males of the HD11 trial (normal reference level= 83)

time	Mean	N	Std.-deviation
1	71,8891	467	30,54669
2	59,3281	296	31,42782
3	68,6343	288	29,96943
4	77,6930	131	25,94816
5	80,5333	125	23,76608
6	81,2500	96	23,87494
7	80,1235	90	24,16561
8	80,8081	88	26,88751
9	83,0303	55	20,38210

Mean values of sexual functioning in males of the HD12 trial (normal reference level= 83)

time	Mean	N	Std.-deviation
1	66,2160	641	33,67431
2	52,7051	382	34,34764
3	67,4155	230	32,62390
4	73,3200	167	31,14312
5	72,6667	150	32,86093
6	74,7346	157	30,54873
7	72,5146	133	30,63058
8	77,4854	114	29,24673
9	69,6732	85	33,43847