Sexual health and testosterone concentration in male lymphoma survivors: A systematic review

Supplementary data provided by Gan et al.

Title: Erectile dysfunction in male lymphoma survivors in a Southeast Asian country

Gan, GG Ng, D.L.C. Leong, Y.C.

Supplementary data for adult men with HL treated with chemotherapy.

Descriptive Statistics

	N	Minimum	Maximum	Mean	Std. Deviation
Age group	104	1	2	1.61	.491
Age	104	18	85	54.93	15.345
Valid N (listwise)	104				

Case Processing Summary

	Cases					
	Valid		Missing		Total	
	N	Percent	N	Percent	N	Percent
ED score all	104	100.0%	0	0.0%	104	100.0%

Descriptives

			Statistic	Std. Error
ED score all	Mean	11.91	.838	
	95% Confidence Interval	Lower Bound	10.25	
	forMean	Upper Bound	13.58	
	5% Trimmed Mean		11.86	
	Median		15.00	
	Variance	73.109		
	Std. Deviation	8.550		
	Minimum	0		
	Maximum	25		
	Range			
Interquartile Range			16	
	Skewness		205	.237
	Kurtosis		-1.475	.469

ED group

					Cumulative
		Frequency	Percent	Valid Percent	Percent
Valid	No ED	13	12.5	12.5	12.5
	Yes ED	56	53.8	53.8	66.3
	not sexually active (< 5)	35	33.7	33.7	100.0
	Total	104	100.0	100.0	