

Tables S1 + S2 Long-term (12 months) results of linear mixed models for pairwise comparisons of treatment groups for the primary outcome measures

Table S1 depression (PHQ-9)

Fixed Effects	Comparison					
	Human guided versus care as usual		Computer guided versus care as usual		Human guided versus computer guided	
Parameter	B (SE)	P value	B (SE)	P value	B (SE)	P value
Intercept	<i>8.34 (0.27)</i>	<i><.001</i>	<i>8.34 (0.27)</i>	<i><.001</i>	<i>7.90 (0.28)</i>	<i><.001</i>
Time	<i>4.03 (0.87)</i>	<i><.001</i>	<i>4.03 (0.87)</i>	<i><.001</i>	<i>6.30 (0.83)</i>	<i><.001</i>
Condition	-0.44 (0.38)	.25	-0.22 (0.40)	.57	-0.22 (0.40)	.58
Timexcondition	2.27 (1.20)	.06	2.71 (1.22)	.03	-0.44 (1.15)	.70

Note. Italicized values are statistically significant (P<.002)

Table S2 anxiety (GAD-7)

Fixed Effects	Comparison					
	Human guided versus care as usual		Computer guided versus care as usual		Human guided versus computer guided	
Parameter	B (SE)	P value	B (SE)	P value	B (SE)	P value
Intercept	<i>7.67 (0.24)</i>	<i><.001</i>	<i>7.67 (0.24)</i>	<i><.001</i>	<i>6.72 (0.25)</i>	<i><.001</i>
Time	<i>3.66 (0.74)</i>	<i><.001</i>	<i>3.66 (0.74)</i>	<i><.001</i>	<i>5.37 (0.79)</i>	<i><.001</i>
Condition	-0.95 (0.34)	<.01	-0.46 (0.36)	.20	-0.49 (0.36)	.17
Timexcondition	1.71 (1.06)	.11	1.88 (1.09)	.09	-0.17 (1.10)	.88

Note. Italicized values are statistically significant (P<.002)