

Supplementary Online Content

Correll CU, Johnston K, Turkoz I, et al. Three-year outcomes of 6-month paliperidone palmitate in adults with schizophrenia: an open-label extension study of a randomized clinical trial. *JAMA Netw Open*. 2024;7(7):e2421495. doi:10.1001/jamanetworkopen.2024.21495

eTable 1. CGI-S Change From Baseline to Month 36 by Age

eTable 2. CGI-S Change From Baseline at Each Visit by Body Mass Index

eTable 3. CGI-S Change From Baseline at Each Visit by Duration of Illness

eTable 4. CGI-S Change From Baseline at Each Visit by Sex

eTable 5. PANSS Total Score Change From Baseline to Month 36 by Age

eTable 6. PANSS Total Score Change From Baseline at Each Visit by Body Mass Index

eTable 7. PANSS Total Score Change From Baseline at Each Visit by Duration of Illness

eTable 8. PANSS Total Score Change From Baseline at Each Visit by Sex

eTable 9. PSP Total Score Change From Baseline at Each Visit by Age

eTable 10. PSP Total Score Change From Baseline at Each Visit by Body Mass Index

eTable 11. PSP Total Score Change From Baseline at Each Visit by Duration of Illness

eTable 12. PSP Total Score Change From Baseline at Each Visit by Sex

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable 1. CGI-S Change from Baseline (DB) to Month 36 by Age^a

	18-25 years	>25-50 years	>50 years
Month 36			
N	7	77	15
LS-Mean (SE)	-0.22 (0.140)	-0.29 (0.055)	-0.26 (0.108)
Diff of LS-Means (SE) 18-25 Years - others		0.07 (0.147)	0.05 (0.173)
95% CI of difference		(-0.220; 0.357)	(-0.293; 0.388)
<i>p</i> -value		0.642	0.784
Diff of LS-Means (SE) >25-50 Years - > 50 Years			-0.02 (0.111)
95% CI of difference			(-0.239; 0.198)
<i>p</i> -value			0.851

Abbreviations: CGI-S, Clinical Global Impression-Severity scale; DB, double-blind study; df, degrees of freedom; LS, least square; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by age group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline CGI-S Score as a fixed effect covariate; age group, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and age group-by-visit interaction. The correlation of the repeated measures is modeled with an AR(1) covariance structure.

eTable 2. CGI-S Change from Baseline (DB) at Each Visit by Body Mass Index^a

	LS Mean Difference				df	P-value
	<30 kg/m ² Estimate (SE)	≥30 kg/m ² Estimate (SE)	(<30 kg/m ² vs ≥30 kg/m ²) Estimate (SE)	(<30 kg/m ² vs ≥30 kg/m ²) 95% CI		
Model Estimates/Treatment Comparison (BMI <30 kg/m² vs ≥30 kg/m²) at Each Visit						
Week 4	-0.05 (0.054)	-0.04 (0.075)	-0.01 (0.083)	(-0.175; 0.151)	302	0.881
Month 2	-0.06 (0.054)	-0.01 (0.075)	-0.05 (0.083)	(-0.215; 0.111)	302	0.532
Month 3	-0.12 (0.054)	-0.06 (0.075)	-0.06 (0.083)	(-0.218; 0.108)	302	0.506
Month 4	-0.12 (0.054)	-0.06 (0.075)	-0.06 (0.083)	(-0.218; 0.108)	302	0.506
Month 5	-0.14 (0.054)	-0.09 (0.075)	-0.05 (0.083)	(-0.214; 0.112)	302	0.539
Month 6	-0.14 (0.054)	-0.09 (0.075)	-0.05 (0.083)	(-0.214; 0.112)	302	0.539
Month 7	-0.18 (0.054)	-0.12 (0.075)	-0.06 (0.083)	(-0.222; 0.105)	302	0.481
Month 8	-0.19 (0.054)	-0.12 (0.075)	-0.07 (0.083)	(-0.233; 0.093)	302	0.398
Month 9	-0.19 (0.054)	-0.15 (0.075)	-0.04 (0.083)	(-0.206; 0.121)	302	0.609
Month 10	-0.21 (0.054)	-0.09 (0.075)	-0.12 (0.083)	(-0.285; 0.042)	302	0.144
Month 11	-0.23 (0.054)	-0.15 (0.075)	-0.08 (0.083)	(-0.241; 0.085)	302	0.349
Month 12	-0.30 (0.054)	-0.12 (0.075)	-0.18 (0.083)	(-0.339; 0.013)	302	0.034
Month 15	-0.21 (0.054)	-0.12 (0.076)	-0.09 (0.083)	(-0.258; 0.069)	306	0.258
Month 18	-0.24 (0.054)	-0.12 (0.076)	-0.12 (0.083)	(-0.286; 0.042)	310	0.145
Month 21	-0.28 (0.054)	-0.19 (0.077)	-0.08 (0.084)	(-0.247; 0.085)	322	0.336

Month 24	-0.28 (0.054)	-0.16 (0.077)	-0.12 (0.085)	(-0.283; 0.051)	326	0.173
Month 27	-0.28 (0.055)	-0.18 (0.078)	-0.10 (0.086)	(-0.272; 0.066)	338	0.230
Month 30	-0.28 (0.055)	-0.21 (0.079)	-0.07 (0.087)	(-0.240; 0.100)	346	0.419
Month 33	-0.32 (0.055)	-0.14 (0.080)	-0.18 (0.088)	(-0.350; - 0.005)	357	0.043
Month 36	-0.31 (0.056)	-0.18 (0.081)	-0.13 (0.088)	(-0.303; 0.045)	362	0.145

Abbreviations: BMI, body mass index; CGI-S, Clinical Global Impression-Severity scale; DB, double-blind study; df, degrees of freedom; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by BMI group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline CGI-S Score as a fixed effect covariate; BMI group, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and BMI group-by-visit interaction. The correlation of the repeated measures is modeled with an AR(1) covariance structure.

eTable 3. CGI-S Change from Baseline (DB) at Each Visit by Duration of Illness^a

	LS Mean Difference				df	P-value
	≤3 Years Estimate (SE)	>3 Years Estimate (SE)	(≤3 Years vs >3 Years) Estimate (SE)	(≤3 Years vs >3 Years) 95% CI		
Model Estimates/Treatment Comparison (≤3 Years vs >3 Years) at Each Visit						
Week 4	-0.02 (0.071)	-0.07 (0.058)	0.06 (0.083)	(-0.109; 0.219)	298	0.509
Month 2	0.01 (0.071)	-0.08 (0.058)	0.09 (0.083)	(-0.069; 0.259)	298	0.257
Month 3	-0.07 (0.071)	-0.13 (0.058)	0.06 (0.083)	(-0.106; 0.223)	298	0.485
Month 4	-0.07 (0.071)	-0.13 (0.058)	0.06 (0.083)	(-0.106; 0.223)	298	0.485
Month 5	-0.15 (0.071)	-0.13 (0.058)	-0.02 (0.083)	(-0.189; 0.139)	298	0.765
Month 6	-0.15 (0.071)	-0.13 (0.058)	-0.02 (0.083)	(-0.189; 0.139)	298	0.765
Month 7	-0.15 (0.071)	-0.18 (0.058)	0.02 (0.083)	(-0.142; 0.186)	298	0.791
Month 8	-0.13 (0.071)	-0.20 (0.058)	0.07 (0.083)	(-0.091; 0.238)	298	0.379
Month 9	-0.18 (0.071)	-0.19 (0.058)	0.01 (0.083)	(-0.158; 0.170)	298	0.942
Month 10	-0.15 (0.071)	-0.20 (0.058)	0.05 (0.083)	(-0.119; 0.210)	298	0.585
Month 11	-0.15 (0.071)	-0.23 (0.058)	0.08 (0.083)	(-0.083; 0.245)	298	0.333
Month 12	-0.24 (0.071)	-0.26 (0.058)	0.02 (0.083)	(-0.143; 0.185)	298	0.800
Month 15	-0.18 (0.071)	-0.20 (0.058)	0.02 (0.084)	(-0.146; 0.183)	302	0.825
Month 18	-0.12 (0.071)	-0.25 (0.058)	0.12 (0.084)	(-0.042; 0.288)	304	0.144
Month 21	-0.24 (0.072)	-0.27 (0.058)	0.03 (0.085)	(-0.139; 0.194)	314	0.748
Month 24	-0.18 (0.073)	-0.28 (0.058)	0.10 (0.085)	(-0.066; 0.270)	322	0.235

Month 27	-0.23 (0.073)	-0.27 (0.059)	0.04 (0.086)	(-0.126; 0.212)	329	0.616
Month 30	-0.23 (0.074)	-0.29 (0.059)	0.06 (0.087)	(-0.115; 0.226)	336	0.522
Month 33	-0.22 (0.076)	-0.30 (0.059)	0.09 (0.088)	(-0.089; 0.259)	356	0.336
Month 36	-0.26 (0.078)	-0.29 (0.059)	0.03 (0.090)	(-0.147; 0.207)	373	0.743

Abbreviations: CGI-S, Clinical Global Impression-Severity scale; DB, double-blind study; df, degrees of freedom; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by duration of illness group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline CGI-S Score as a fixed effect covariate; duration of illness, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and duration of illness-by-visit interaction. The correlation of the repeated measures is modeled with an AR(1) covariance structure.

eTable 4. CGI-S Change from Baseline (DB) at Each Visit by Sex^a

	LS Mean Difference				df	P-value
	Female Estimate (SE)	Male Estimate (SE)	(Female vs Male) Estimate (SE)	(Female vs Male) 95% CI		
Model Estimates/Treatment Comparison (Female - Male) at Each Visit						
Week 4	-0.12 (0.073)	-0.03 (0.052)	-0.09 (0.079)	(-0.249; 0.063)	316	0.240
Month 2	-0.15 (0.073)	-0.01 (0.052)	-0.13 (0.079)	(-0.287; 0.024)	316	0.097
Month 3	-0.15 (0.073)	-0.10 (0.052)	-0.05 (0.079)	(-0.203; 0.108)	316	0.551
Month 4	-0.20 (0.073)	-0.07 (0.052)	-0.12 (0.079)	(-0.280; 0.032)	316	0.118
Month 5	-0.20 (0.073)	-0.11 (0.052)	-0.09 (0.079)	(-0.244; 0.068)	316	0.268
Month 6	-0.20 (0.073)	-0.11 (0.052)	-0.09 (0.079)	(-0.244; 0.068)	316	0.268
Month 7	-0.28 (0.073)	-0.12 (0.052)	-0.15 (0.079)	(-0.310; 0.001)	316	0.051
Month 8	-0.33 (0.073)	-0.11 (0.052)	-0.22 (0.079)	(-0.375; - 0.064)	316	0.006
Month 9	-0.30 (0.073)	-0.14 (0.052)	-0.17 (0.079)	(-0.325; - 0.013)	316	0.034
Month 10	-0.36 (0.073)	-0.11 (0.052)	-0.25 (0.079)	(-0.401; - 0.090)	316	0.002
Month 11	-0.41 (0.073)	-0.12 (0.052)	-0.29 (0.079)	(-0.442; - 0.131)	316	<0.001
Month 12	-0.41 (0.073)	-0.18 (0.052)	-0.23 (0.079)	(-0.382; - 0.070)	316	0.005
Month 15	-0.38 (0.073)	-0.11 (0.052)	-0.27 (0.079)	(-0.428; - 0.117)	317	<0.001
Month 18	-0.49 (0.073)	-0.09 (0.052)	-0.40 (0.080)	(-0.559; - 0.246)	322	<0.001
Month 21	-0.49 (0.074)	-0.16 (0.052)	-0.33 (0.080)	(-0.491; - 0.176)	330	<0.001
Month 24	-0.46 (0.074)	-0.16 (0.053)	-0.30 (0.081)	(-0.463; - 0.146)	337	<0.001

Month 27	-0.49 (0.074)	-0.16 (0.053)	-0.34 (0.081)	(-0.498; -0.179)	342	<0.001
Month 30	-0.44 (0.075)	-0.20 (0.053)	-0.24 (0.081)	(-0.398; -0.077)	344	0.004
Month 33	-0.47 (0.075)	-0.20 (0.054)	-0.27 (0.082)	(-0.435; -0.111)	356	0.001
Month 36	-0.44 (0.076)	-0.21 (0.054)	-0.23 (0.083)	(-0.392; -0.066)	361	0.006

Abbreviations: CGI-S, Clinical Global Impression-Severity scale; DB, double-blind study; df, degrees of freedom; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by sex. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline CGI-S Score as a fixed effect covariate; sex, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and sex-by-visit interaction. The correlation of the repeated measures is modeled with an AR(1) covariance structure.

eTable 5. PANSS Total Score Change from Baseline (DB) to Month 36 by Age^a

	18-25 years	>25-50 years	>50 years
Month 36			
N	8	78	15
LS-Mean (SE)	-2.1 (3.19)	-2.4 (1.11)	-1.8 (2.48)
Diff of LS-Means (SE) 18-25 Years - others		0.4 (3.36)	-0.3 (4.02)
95% CI of difference		(-6.33; 7.07)	(-8.29; 7.75)
<i>p</i> -value		0.913	0.947
Diff of LS-Means (SE) >25-50 Years - > 50 Years			-0.6 (2.68)
95% CI of difference			(-5.97; 4.70)
<i>p</i> -value			0.813

Abbreviations: DB, double-blind study; df, degrees of freedom; LS, least square; PANSS, Positive and Negative Syndrome Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by age group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PANSS Total Score as a fixed effect covariate; age group, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and age group-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.

eTable 6. PANSS Total Score Change from Baseline (DB) at Each Visit by Body Mass Index^a

	LS Mean Difference				df	P-value
	<30 kg/m ² Estimate (SE)	≥30 kg/m ² Estimate (SE)	(<30 kg/m ² vs ≥30 kg/m ²) Estimate (SE)	(<30 kg/m ² vs ≥30 kg/m ²) 95% CI		
Model Estimates/Treatment Comparison (BMI <30 kg/m² vs ≥30 kg/m²) at Each Visit						
Month 3	-1.1 (0.47)	-1.4 (0.62)	0.3 (0.65)	(-1.01; 1.56)	115	0.678
Month 6	-2.5 (0.61)	-3.0 (0.86)	0.5 (0.96)	(-1.42; 2.38)	122	0.620
Month 9	-3.3 (0.61)	-2.5 (0.86)	-0.8 (0.96)	(-2.70; 1.10)	124	0.405
Month 12	-4.2 (0.67)	-3.5 (0.97)	-0.7 (1.10)	(-2.91; 1.44)	121	0.507
Month 24	-3.4 (1.08)	-0.4 (1.61)	-3.0 (1.89)	(-6.73; 0.76)	118	0.118
Month 36	-3.2 (1.15)	-0.7 (1.74)	-2.6 (2.04)	(-6.63; 1.48)	77	0.209

Abbreviations: BMI, body mass index; DB, double-blind study; df, degrees of freedom; PANSS, Positive and Negative Syndrome Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by BMI group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PANSS Total Score as a fixed effect covariate; BMI group, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and BMI group-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.

eTable 7. PANSS Total Score Change from Baseline (DB) at Each Visit by Duration of Illness^a

	LS Mean Difference				df	P-value
	≤3 Years Estimate (SE)	>3 Years Estimate (SE)	(≤3 Years vs >3 Years) Estimate (SE)	(≤3 Years vs >3 Years) 95% CI		
Model Estimates/Treatment Comparison (≤3 Years vs >3 Years) at Each Visit						
Month 3	-1.4 (1.11)	-1.7 (0.90)	0.3 (1.30)	(-2.27; 2.86)	246	0.824
Month 6	-2.6 (1.11)	-3.2 (0.90)	0.6 (1.30)	(-1.97; 3.16)	246	0.647
Month 9	-2.5 (1.11)	-3.9 (0.90)	1.4 (1.30)	(-1.16; 3.97)	246	0.282
Month 12	-4.6 (1.11)	-4.4 (0.90)	-0.2 (1.30)	(-2.77; 2.35)	246	0.872
Month 24	-3.3 (1.12)	-2.9 (0.91)	-0.4 (1.32)	(-2.99; 2.20)	256	0.765
Month 36	-4.3 (1.19)	-2.9 (0.93)	-1.5 (1.39)	(-4.19; 1.27)	290	0.294

Abbreviations: DB, double-blind study; df, degrees of freedom; PANSS, Positive and Negative Syndrome Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by duration of illness group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PANSS Total Score as a fixed effect covariate; duration of illness, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and duration of illness-by-visit interaction. The correlation of the repeated measures is modeled with an AR(1) covariance structure.

eTable 8. PANSS Total Score Change from Baseline (DB) at Each Visit by Sex^a

	LS Mean Difference				df	P-value
	Female Estimate (SE)	Male Estimate (SE)	(Female vs Male) Estimate (SE)	(Female vs Male) 95% CI		
Model Estimates/Treatment Comparison (Female - Male) at Each Visit						
Month 3	-1.2 (0.64)	-1.2 (0.46)	-0.1 (0.64)	(-1.34; 1.20)	115	0.915
Month 6	-2.9 (0.86)	-2.5 (0.60)	-0.5 (0.95)	(-2.34; 1.41)	123	0.624
Month 9	-4.3 (0.85)	-2.5 (0.60)	-1.8 (0.94)	(-3.61; 0.09)	123	0.062
Month 12	-4.7 (0.96)	-3.7 (0.67)	-1.0 (1.08)	(-3.17; 1.11)	120	0.344
Month 24	-2.8 (1.60)	-2.4 (1.10)	-0.3 (1.88)	(-4.06; 3.40)	118	0.861
Month 36	-2.8 (1.70)	-2.3 (1.17)	-0.5 (2.01)	(-4.50; 3.52)	75	0.807

Abbreviations: DB, double-blind study; df, degrees of freedom; PANSS, Positive and Negative Syndrome Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by sex. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PANSS Total Score as a fixed effect covariate; sex, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and sex-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.

eTable 9. PSP Total Score Change from Baseline (DB) at Each Visit by Age^a

	18-25 years	>25-50 years	>50 years
Month 36			
N	8	79	15
LS-Mean (SE)	4.5 (2.59)	2.9 (0.97)	2.6 (2.07)
Diff of LS-Means (SE) 18-25 Years - others		1.6 (2.72)	1.9 (3.27)
95% CI of difference		(-3.84; 6.96)	(-4.60; 8.38)
p-value		0.567	0.564
Diff of LS-Means (SE) >25-50 Years - > 50 Years			0.3 (2.19)
95% CI of difference			(-4.02; 4.67)
p-value			0.881

Abbreviations: DB, double-blind study; df, degrees of freedom; LS, least square; PSP, Personal and Social Performance Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by age group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PSP Total Score as a fixed effect covariate; age group, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and age group-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.

eTable 10. PSP Total Score Change from Baseline (DB) at Each Visit by Body Mass Index^a

	LS Mean Difference				df	P-value
	<30 kg/m ² Estimate (SE)	≥30 kg/m ² Estimate (SE)	(<30 kg/m ² vs ≥30 kg/m ²) Estimate (SE)	(<30 kg/m ² vs ≥30 kg/m ²) 95% CI		
Model Estimates/Treatment Comparison (BMI <30 kg/m² vs ≥30 kg/m²) at Each Visit						
Month 3	0.7 (0.64)	0.1 (0.85)	0.6 (0.89)	(-1.13; 2.42)	114	0.472
Month 6	1.1 (0.73)	1.1 (1.02)	-0.1 (1.12)	(-2.30; 2.13)	116	0.939
Month 9	2.5 (0.73)	1.0 (1.02)	1.5 (1.12)	(-0.76; 3.67)	123	0.196
Month 12	3.1 (0.75)	1.4 (1.04)	1.8 (1.14)	(-0.49; 4.04)	115	0.123
Month 24	3.2 (0.98)	1.2 (1.43)	2.0 (1.63)	(-1.28; 5.19)	116	0.234
Month 36	3.7 (1.00)	1.7 (1.47)	2.0 (1.68)	(-1.33; 5.34)	97	0.236

Abbreviations: BMI, body mass index; DB, double-blind study; df, degrees of freedom; PSP, Personal and Social Performance Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by BMI group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PSP Total Score as a fixed effect covariate; BMI group, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and BMI group-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.

eTable 11. PSP Total Score Change from Baseline (DB) at Each Visit by Duration of Illness^a

	≤3 Years	>3 Years	Difference	Difference	df	P-value
	Estimate (SE)	Estimate (SE)	(≤3 Years vs >3 Years)	(≤3 Years vs >3 Years)		
			Estimate (SE)	95% CI		
Model Estimates/Treatment Comparison (≤3 Years vs >3 Years) at Each Visit						
Month 3	0.4 (0.78)	0.6 (0.70)	-0.2 (0.90)	(-1.97; 1.60)	114	0.838
Month 6	0.5 (0.95)	1.4 (0.79)	-0.9 (1.12)	(-3.09; 1.33)	117	0.433
Month 9	0.9 (0.95)	2.5 (0.79)	-1.6 (1.12)	(-3.84; 0.59)	123	0.149
Month 12	2.6 (0.98)	2.6 (0.80)	-0.0 (1.15)	(-2.29; 2.27)	115	0.993
Month 24	3.3 (1.40)	2.4 (1.02)	1.0 (1.65)	(-2.28; 4.25)	119	0.552
Month 36	3.8 (1.44)	2.8 (1.04)	1.0 (1.70)	(-2.39; 4.34)	97	0.566

Abbreviations: DB, double-blind study; df, degrees of freedom; PSP, Personal and Social Performance Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by duration of illness group. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PSP Total Score as a fixed effect covariate; duration of illness, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and duration of illness-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.

eTable 12. PSP Total Score Change from Baseline (DB) at Each Visit by Sex^a

	LS Mean Difference				df	P-value
	Female Estimate (SE)	Male Estimate (SE)	(Female vs Male) Estimate (SE)	(Female vs Male) 95% CI		
Model Estimates/Treatment Comparison (Female - Male) at Each Visit						
Month 3	0.6 (0.86)	0.5 (0.63)	0.2 (0.88)	(-1.55; 1.92)	114	0.834
Month 6	0.9 (1.02)	1.2 (0.73)	-0.3 (1.09)	(-2.47; 1.86)	115	0.782
Month 9	2.5 (1.03)	1.8 (0.73)	0.7 (1.10)	(-1.48; 2.88)	122	0.525
Month 12	2.7 (1.04)	2.6 (0.74)	0.2 (1.13)	(-2.08; 2.39)	113	0.891
Month 24	3.4 (1.41)	2.3 (0.98)	1.1 (1.61)	(-2.09; 4.29)	116	0.494
Month 36	3.6 (1.44)	2.9 (1.01)	0.7 (1.65)	(-2.60; 3.95)	94	0.684

Abbreviations: df, degrees of freedom; PSP, Personal and Social Performance Scale; SE, standard error.

^aPer Mixed Model Repeated Measures ANCOVA, change from baseline (DB) at each visit by sex. The estimate, standard error, 95% CI, and *p*-value are based on a repeated measures mixed effects ANCOVA model with baseline PSP Total Score as a fixed effect covariate; sex, country, and visit (scheduled visit assessment) as fixed effect (categorical) factors; and sex-by-visit interaction. The correlation of the repeated measures is modeled with an unstructured covariance structure.