

Supplementary Table S2. Evolution of urinary symptoms—paired results.

	Mean ± SD	[IQR]	(95%CI)
Dysuria			
Pre-op	0.6 ± 1.8	[0; 0]	(0.1; 1.1)
Six months	0.5 ± 1.8	[0; 0]	(0.0; 1.0)
Difference	−0.1 ± 1.4	[0; 0]	(−0.5; 0.3)
P-value	0.611		
Urinary urgency			
Pre-op	1.8 ± 3.0	[0; 5]	(1.0; 2.6)
Six months	1.1 ± 2.5	[0; 0]	(0.4; 1.7)
Difference	−0.8 ± 2.7	[−0.3; 0]	(−1.5; −0.0)
P-value	0.026		
Urge incontinence			
Pre-op	0.5 ± 1.4	[0; 0]	(0.1; 0.9)
Six months	0.4 ± 1.5	[0; 0]	(0.1; 0.8)
Difference	−0.1 ± 1.6	[0; 0]	(−0.5; 0.3)
P-value	0.77		
Stress incontinence			
Pre-op	0.9 ± 2.0	[0; 0]	(0.4; 1.4)
Six months	0.8 ± 1.8	[0; 0]	(0.3; 1.2)
Difference	−0.1 ± 2.1	[0; 0]	(−0.7; 0.4)
P-value	0.62		
Incomplete bladder emptying			
Pre-op	2.0 ± 2.8	[0; 5]	(1.3; 2.7)
Six months	1.2 ± 2.4	[0; 2]	(0.6; 1.9)
Difference	−0.8 ± 2.4	[−1.3; 0]	(−1.4; −0.1)
P-value	0.012		