

<b>Subclinical Cut-Off Scores</b>			
<b>Domain</b>	<b>Instrument</b>	<b>Cut-Off</b>	<b>References</b>
Depression	Patient Health Questionnaire-9 for Adolescents (PHQ-A)	5≤/27	[1]
Bipolar	Altman Self-Rating Mania Scale, Child version (ASRM)	6≤/20	[2–4]
Anxiety	Generalized Anxiety Disorder 7 (GAD-7)	5≤/21	[5]
Obsessive-compulsive disorder	Obsessive Compulsive Disorder Scale (SOCS)	6≤/14	[6,7]
Psychotic-like experiences	Community Assessment of Psychic Experiences - Positive (CAPE-P15)	1.47≤/45	[8,9]
Alcohol/Substance abuse	CRAFFTd	2≤/6	[10]
Eating behavior	Eating Disorder Examination-Questionnaire adapted for children (ChEDE-Q8)	2.38≤/6	[12]
Conduct disorder	Strength and Difficulties Questionnaire, self-report (SDQ-S), subscale conduct disorders	4≤/10	[13,14]
Non-suicidal self-injurious behavior	Screening questions (frequency, expectation, associated characteristics), Diagnostic Interview for Mental Disorders in children and adolescents, Kinder-DIPS, Additional module NSSV	Item 1 (yes) Item 2 (5 or more days)	[15]

## References

1. Johnson JG, Harris ES, Spitzer RL, Williams JBW. The patient health questionnaire for adolescents. *Journal of Adolescent Health*. 2002;30(3):196–204.
2. Altman E, Hedeker D, Peterson JL, Davis JM. A comparative evaluation of three self-rating scales for acute mania. *Biological Psychiatry*. 2001;50(6):468–71.
3. Haussleiter IS, Lorek S, Zeschel E, Mattausch S, Meyers M, Juckel G. At-risk symptoms of bipolar disorder in a university student cohort. *Psychiatry Research*. 2018;265:324–33.
4. Meyer TD, Crist N, La Rosa N, Ye B, Soares JC, Bauer IE. Are existing self-ratings of acute manic symptoms in adults reliable and valid?—A systematic review. *Bipolar Disorders*. 2020;22(6):558–68.
5. Löwe B, Decker O, Müller S, Brähler E, Schellberg D, Herzog W, et al. Validation and Standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the General Population. *Medical Care*. 2008;46(3):266–74.
6. Piqueras JA, Rodríguez-Jiménez T, Ortiz AG, Moreno E, Lázaro L, Godoy A. Validation of the Short Obsessive–Compulsive Disorder Screener (SOCS) in children and adolescents. *BJPsych open*. 2015;1(1):21–6.
7. Zijlmans J, Marhe R, van der Ende J, Verhulst FC, Popma A, Tiemeier H, et al. Children with Obsessive–Compulsive Symptomology in the General Population: Different Subtypes? *Journal of Developmental & Behavioral Pediatrics*. 2017;38(7):476.
8. Capra C, Kavanagh DJ, Hides L, Scott J. Brief screening for psychosis-like experiences. *Schizophrenia Research*. 2013;149(1–3):104–7.

9. Knight C, Stochl J, Sonesson E, Russo DA, Jones PB, Perez J. Revisiting CAPE-P15 cut-off values to increase sensitivity for detecting psychotic experiences in primary care. *Schizophrenia Research*. 2020;216:507–10.
10. Tossmann P, Kasten L, Lang P, Strüber E. Bestimmung der konkurrenten Validität des CRAFFT-d: Ein Screeninginstrument für problematischen Alkoholkonsum bei Jugendlichen. *Zeitschrift für Kinder- und Jugendpsychiatrie und Psychotherapie*. 2009;37(5):451–9.
11. Kliem S, Schmidt R, Vogel M, Hiemisch A, Kiess W, Hilbert A. An 8-item short form of the Eating Disorder Examination-Questionnaire adapted for children (ChEDE-Q8). *International Journal of Eating Disorders*. 2017;50(6):679–86.
12. Becker A, Wang B, Kunze B, Otto C, Schlack R, Hölling H, et al. Normative Data of the Self-Report Version of the German Strengths and Difficulties Questionnaire in an Epidemiological Setting. *Zeitschrift für Kinder- und Jugendpsychiatrie und Psychotherapie*. 2018 Nov 1;46(6):523–33.
13. Goodman R. Psychometric Properties of the Strengths and Difficulties Questionnaire. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2001 Nov;40(11):1337–45.
14. In-Albon T, Kraus L. Nichtsuizidale Selbstverletzungen [Non-suicidal self-injury]. Universität Koblenz-Landau, Klinische Psychologie und Psychotherapie des Kindes- und Jugendalters. 2017.