medRxiv preprint doi: https://doi.org/10.1101/2024.08.01.24311376; this version posted August 3, 2024. The copyright holder for this preprint (which was not certified by peer review) is the author/funder, who has granted medRxiv a license to display the preprint in perpetuity. It is made available under a CC-BY-NC 4.0 International license.

12	4.6	4.8	4.9	5.0	5.1	5.3	5.4	5.5	5.7	5.9	6.1	
0 - T	4.2	4.4	4.5	4.6	4.7	4.8	5.0	5.1	5.3	5.5	5.7	
Ì	3.9	4.0	4.1	4.2	4.3	4.4	4.6	4.7	4.9	5.1	5.3	
 -	3.5	3.6	3.7	3.8	3.9	4.0	4.2	4.3	4.5	4.7	4.9	
	3.1	3.2	3.3	3.4	3.5	3.7	3.8	3.9	4.1	4.3	4.5	
	2.7	2.8	2.9	3.0	3.1	3.3	3.4	3.5	3.7	3.9	4.1	
17 - 77	2.4	2.5	2.6	2.7	2.8	2.9	3.0	3.1	3.3	3.5	3.7	
N C7 -	2.1	2.2	2.2	2.3	2.4	2.5	2.6	2.8	2.9	3.1	3.3	
24 - 1	1.8	1.8	1.9	2.0	2.1	2.2	2.3	2.4	2.6	2.7	2.9	
й СУ -	1.5	1.6	1.6	1.7	1.8	1.8	1.9	2.1	2.2	2.3	2.5	
V	1.3	1.3	1.4	1.4	1.5	1.5	1.6	1.7	1.9	2.0	2.2	

269	SD of Differences Between	StimIII and StimV Pairs	Across Censoring Thresholds
			5

Supplemental Figure 1. Heatmap depicting the standard deviation of total deviation
differences between pairs of stimulus size III and stimulus size V points across varying
censoring thresholds where at least one stimulus was censored. For the optimal
censoring threshold of 21 dB for stimulus size III and 24 for stimulus size V, the
standard deviation is 1.9 dB.

276

270

277 References