

Supplementary Material

Attributional style in Borderline Personality Disorder is associated with self-esteem and loneliness

Schulze, A. ^{1*}, Rommelfanger, B. ¹, Schendel, E. ¹, Schott, H. ¹, Lerchl, A. ¹, Vonderlin, R. ²,
Lis, S ^{1,2}.

¹Department of Clinical Psychology, Central Institute of Mental Health, Medical Faculty
Mannheim, University of Heidelberg, Germany

²Department of Psychosomatic Medicine and Psychotherapy, Central Institute of Mental
Health, Medical Faculty Mannheim, University of Heidelberg, Germany

Table S1*Results of the 2x2x3 rm-ANOVA*

effect	df _{1,2}	<i>F</i>	<i>p</i>	η^2_p
group	1, 62	10.40	.002	.144
attribution dimension	2, 124	9.69	<.001	.135
attribution dimension * group	2, 124	9.86	<.001	.137
valence	1, 62	.05	.817	.001
valence * group	1, 62	47.81	<.001	.435
attribution dimension * valence	2, 124	27.62	<.001	.308
attribution dimension * valence * group	2, 124	3.28	.042	.050

*Note: p's are reported for Greenhouse-Geisser corrected degrees of freedom***Table S2***Comparison between groups*

attribution dimension	scenario	<i>t</i> (62)	<i>p</i>	95% CI		effect size Cohen's <i>d</i>
internality	positive	6.112	<.001	0.79	1.55	1.53
	negative	-5.681	<.001	-1.53	-0.74	-1.42
stability	positive	1.298	.199	-0.14	0.68	0.32
	negative	-6.293	<.001	-2.01	-1.04	-1.57
globality	positive	1.199	.235	-0.21	0.84	0.30
	negative	-6.689	<.001	-2.04	-1.10	-1.67

Note: Bonferroni corrected threshold is $p < .008$

Figure S1

Association between attributional style with duration of loneliness and interpersonal functioning

