

Supplementary Information

American Journal of Cardiovascular Drugs

Strategy for an early simultaneous introduction of four-pillars of heart failure therapy: results from a single center experience

Paolo Severino^{1†*}, Andrea D'Amato^{1†}, Silvia Prosperi¹, Marco Valerio Mariani¹, Vincenzo Myftari¹, Aurora Labbro Francia¹, Claudia Cestiè¹, Elisa Tomarelli¹, Giovanna Manzi¹, Lucia Ilaria Birtolo¹, Stefanie Marek-Iannucci¹, Viviana Maestrini¹, Massimo Mancone¹, Roberto Badagliacca¹, Francesco Fedele¹, Carmine Dario Vizza¹.

¹ Department of Clinical, Internal, Anesthesiology and Cardiovascular Sciences, Sapienza University of Rome, Viale del Policlinico 155, 00161 Rome, Italy.

† These Authors contributed equally

* Corresponding Author: Email paolo.severino@uniroma1.it

Table S1. Baseline features of the total population and each group before the propensity score matching (PMS) analysis at hospital admission.

Parameter	Overall population (N=320)	Group 1 (N= 139)	Group 2 (N=181)	p-value
Age, years (SD)	71.3 (±10.9)	69.7 (±13.2)	73 (±11.7)	0.01
Male gender, n (%)	87 (27)	29 (20.9)	58 (32.3)	0.03
Arterial hypertension, n (%)	263 (82)	111 (79.9)	152 (84)	0.38
Type II diabetes mellitus, n (%)	60 (18.7)	22 (15.9)	38 (21)	0.25
Dyslipidemia, n (%)	160 (50)	58 (41.7)	102 (56.5)	0.013
CVD family history, n (%)	106 (33)	34 (24.5)	72 (40)	0.004
Smoking habit, n (%)	95 (30)	47 (33.8)	48 (26.7)	0.17
eGFR, ml/min (SD)	60.4 (±24.1)	62.2 (±24)	58.4 (±23.5)	0.15
LVEDD, mm (SD)	55.6 (±7.5)	58 (±4.2)	51.8 (±7.2)	< 0.001
IVS, mm (SD)	11.1 (±1.7)	11 (±1.1)	11.3 (±1.7)	0.07
PW, mm (SD)	10.2 (±1.2)	10 (±1.3)	10.5 (±1.4)	0.001
LVEF, % (SD)	33.1 (±10.2)	31.1 (± 9.9)	35.3 (±12.7)	0.001

CVD cardiovascular disease, eGFR estimated glomerular filtration rate, Hb hemoglobin, LVEDD left ventricular end diastolic diameter, IVS interventricular septum, PW posterior wall, LVEF left ventricle ejection fraction, NYHA New York Heart Association.