Supplementary material

Journal: Drugs - Real World Outcomes (DRWO)

Manuscript Title: Demographic and Clinical Characteristics of Patients With Crohn's Disease and Ulcerative Colitis Experiencing Changes in Bowel Urgency Severity – A Real-World Study

Author list: James D. Lewis¹, Theresa H. Gibble², Mingyang Shan², Xian Zhoa³, April N. Naegeli², Ghadeer K. Dawwas⁴

Corresponding author:

James D. Lewis, M.D., M.S.C.E.

University of Pennsylvania

Center for Clinical Epidemiology and Biostatistics

7th Floor, Blockley Hall, 423 Guardian Drive

Philadelphia, Pennsylvania 19104-6021, USA

E-mail: lewisjd@pennmedicine.upenn.edu; fax: (215) 573-5325

¹ University of Pennsylvania, Philadelphia, USA

² Eli Lilly and Company, Indianapolis, USA

³ Syneos Health, Morrisville, North Carolina, USA

⁴Vanderbilt University Medical Center, Tennessee, USA

 Table S1. List of patient-reported outcomes.

Questionnaires	Choice of response		
During the last week, do you have any of the	□ Fatigue		
following symptoms? Check all that apply	☐ Felt depressed or blue		
What is your average (typical) number of bowel movements that you had in a single day last week?			
Over the last 3 days, how would you describe the frequency of your stools?	☐ Normal number of stools		
	☐ 1–2 stools more than normal		
	☐ 3–4 stools more than normal		
	□ >4 stools more than normal		
	☐ Not applicable, I have an ostomy		
Over the last 3 days, how would you	□ Formed		
describe the consistency of most of your bowel movement	☐ Soft or semi-formed		
	☐ Mostly or all liquid		
Over the last 3 days, how much rectal	☐ No blood seen		
bleeding have you experienced?	☐ Blood less than 50% of the time		
	☐ Blood 50% or more of the time		
Over the last 3 days, how much abdominal pain have you had?	□ None		
	□ Mild		
	☐ Moderate		
	□ Severe		
Over the last 3 days, how much urgency have you had before bowel movements?	☐ None. I can wait 15 minutes or longer to have a bowel movement.		

	☐ Mild. I need to get to the bathroom within 5–15 minutes.			
	☐ Moderate. I need to get to the bathroom within 2–5 minutes.			
	☐ Moderately severe. I need to get to the bathroom in less than 2 minutes.			
	☐ Severe. Sometimes I am unable to make it to the bathroom in time.			
	☐ Not applicable. I have an ostomy.			
During the last months, have you had nighttime bowel movement?	□ Yes			
	□ No			
During the last month, have you had leakage of your stool while sleeping?	□ Yes			
	□ No			
During the last month, have you had any	□ Yes			
leakage of stool while you were awake?	□ No			
Please rate your general well-being over the last 3 days	☐ Generally well			
lust 5 days	☐ Slightly under par			
	□ Poor			
	□ Very poor			
	☐ Terrible			
Physician's global assessment of the current disease status	□ Quiescent			
	□ Mild			
	☐ Moderate			
	□ Severe			

Table S2: Patients with and without 6 months urgency data by enrollment urgency data

	Crohn's Disease			Ulcerative Colitis		
Urgency at Baseline	Missing urgency at 6 months, n (%)	With urgency at 6 months, n (%)	P- value	Missing urgency at 6 months, n (%)	With urgency at 6 months, n (%)	P- value
No urgency	604 (35%)	282 (34%)		324 (36%)	188 (43%)	
Mild	547 (31%)	283 (34%)		264 (30%)	125 (29%)	
Moderate	285 (16%)	162 (20%)		129 (15%)	50 (12%)	
Moderate severe or severe	316 (18%)	105 (13%)	0.002	174 (20%)	72 (17%)	0.073

Fig. S1: Patient Attrition Flow Chart

