Supplemental Digital Content 6. This cross table demonstrates how often the most important goal domain was chosen at the primary test and also as the most important or as secondary goal domain at the retest. The values correspond to the number of patients and the percentage of the row total, except for the "Row total" column, where the percentages correspond to the percentage of the column total.

Most important goal domain chosen at primary test	Most important or secondary goal domain chosen at the retest								
	Pain	Activities	Flexibility/ Mobility	Work	Tingling	Strength	Appearance	Numbness/ Sensation	Row total
Pain	31 (89%)	2 (6%)	1 (3%)	0 (0%)	0 (0%)	0 (0%)	1 (3%)	0 (0%)	35 (34%)
Activities	5 (21%)	19 (79%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	24 (24%)
Flexibility/Mobility	1 (7%)	3 (20%)	9 (60%)	0 (0%)	0 (0%)	0 (0%)	2 (13%)	0 (0%)	15 (15%)
Work	1 (13%)	1 (13%)	0 (0%)	5 (63%)	0 (0%)	0 (0%)	0 (0%)	1 (13%)	8 (8%)
Tingling	0 (0%)	1 (17%)	0 (0%)	0 (0%)	5 (80%)	0 (0%)	0 (0%)	0 (0%)	6 (6%)
Strength	2 (33%)	1 (17%)	0 (0%)	0 (0%)	0 (0%)	2 (33%)	1 (17%)	0 (0%)	6 (6%)
Appearance	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	2 (100%)	0 (0%)	2 (2%)
Numbness/Sensation	0 (0%)	1 (20%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	4 (80%)	5 (5%)
No treatment goal	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (100%)	0 (0%)	0 (0%)	0 (0%)	1 (1%)
Column total	40 (39%)	28 (28%)	10 (10%)	5 (5%)	6 (6%)	2 (2%)	6 (6%)	5 (5%)	102 (100%)