Supplemental material

BMJ Publishing Group Limited (BMJ) disclaims all liability and responsibility arising from any reliance placed on this supplemental material which has been supplied by the author(s)

J NeuroIntervent Surg

Online supplemental figure S1

Flow-chart of patient selection

Supplemental material

BMJ Publishing Group Limited (BMJ) disclaims all liability and responsibility arising from any reliance placed on this supplemental material which has been supplied by the author(s)

J NeuroIntervent Surg

Online supplemental table S1

Breakdown of detected CVF (after senior rating) in terms of patients (n=20)

and combined double examinations (n=27)

Pseudonyms	CVF pos. at LD-DSM*	CVF pos. at LD-CTM*
1	0	1
2	1	2
3	0	6
4	0	1
5	1	1
6	0	2
7	0	1
8	0	1
9	0	0
10	0	1
11	0	1
12	0	1
13	3	3
14	1	2
15	1	2
16	0	1
17	1	1
18	1	1
19	0	1
20	0	1
Total number of patients with ≥ 1 CVF	**7**	**19**
21 (follow-up #1)	1	1
22 (follow-up #3)	0	3
23 (follow-up #13)	0	3
24 (follow-up #16)	0	2
25 (bilateral #3)	see #3	
26 (bilateral #11)	see #11	
27 (bilateral #22)	see #22	
Total number of CVF	**10**	**39**

*CVF: CSF-venous fistula, LD-DSM: lateral decubitus digital subtraction myelography, LD-CTM: lateral decubitus CT-myelography, * number of detected CVFs after senior rating (in cases of disagreement between rater 1 and 2)*
Pseudonyms 1-20: 20 patients at initial combined DSM/CTM investigation; pseudonyms 21-24: further combined DSM/CTM examinations of patient #1, #3, #13 and #16 at follow-up (after treatment of one or more CVFs); pseudonyms 25-27: three included bilateral combined DSM/CTM examinations of patient #3, #11 and #3 at follow-up.