

Supplemental Table 1.

Baseline variables	Responder (n=45)	Non-Responder (n=78)	P-value
Age	43.0 (28.0-57.0)	44.5 (35.3-53.0)	0.63
Female	37 (82.2%)	61 (78.2%)	0.59
BMI	27.9 (22.9-33.1)	25.7 (21.2-29.7)	0.14
Marijuana	4 (8.9%)	6 (7.7%)	0.82
Opioids	4 (8.9%)	10 (12.8%)	0.51
Laxatives	17 (38.8%)	27 (34.6%)	0.72
Antiemetics	7 (15.6%)	8 (10.3%)	0.39
Gastroparesis Diet	14 (31.1%)	30 (38.5%)	0.41
Delayed GES	9 (20.0%)	23 (29.8%)	0.24
Delayed GET	19 (42.2%)	33 (42.3%)	0.99
GES retention at 4 hours	3 (1-7)	3 (1-15)	0.55
GES retention at 2 hours	30.0 (15.0-48.0)	35.5 (22.3-53.0)	0.25
GET	4.4 (3.2-6.1)	4.4 (3.5-6.8)	0.65
Small bowel transit time	4.7 (3.4-6.1)	4.7 (3.8-5.8)	0.7
Colonic transit time	42.7 (21.0-70.8)	35.9 (16.1-63.2)	0.32
Number of antral contractions	41.0 (24.0-88.0)	46 (25.0-86.5)	0.65
Antral motility index	11.1 (10.0-12.6)	11.3 (10.1-12.6)	0.97
Number of duodenal contractions	108.0 (61.0-166.0)	91.5 (42.8-168.3)	0.92
Duodenal motility index	12.5 (11.2-13.8)	12.3 (10.6-13.4)	0.39
Infectious prodrome	2 (4.4%)	13 (16.7%)	0.06
Duration	36 (12-72)	36 (24-108)	0.4
Diabetes	17 (37.8%)	16 (20.5%)	0.04
BSS	4 (3-5)	4 (2-6)	0.87
Functional Bowel Disorder	36 (81.8%)	60 (80.0%)	0.81
Constipation	15 (34.9%)	25 (33.8%)	0.9
Functional Dyspepsia	34 (77.3%)	55 (74.3%)	0.72
Nausea Vomiting and Belching	31 (70.5%)	44 (59.5%)	0.23
GCSI			
Baseline GCSI	2.7 (2.0-3.1)	3.1 (2.5-3.5)	0.005
GCSI at 6 months	1.2 (0.8-1.8)	2.8 (2.0-3.6)	<0.0001

Supplemental Table 1. Baseline variables which were evaluated in order to be included in the predictive model and GSCI at baseline and 6 months. The initial model included all variables with $p < 0.25$ which are BMI, Delayed GES, Infectious prodrome, Diabetes, Nausea, Vomiting, and Belching.