


Study

- Omitting Chiurlia 2005 Modena, Italy
- Omitting Javaroni 2011 Brazil
- Omitting Kaiser 2004 USA
- Omitting Kaya 2006 Istanbul, Turkey
- Omitting Murata 2012 Saitama, Japan
- Omitting Stuckey 2006 Nedlands, Australia
- Omitting Uslu 2005 Istanbul, Turkey
- Omitting Vlachopoulos 2008 Athens, Greece
- Omitting Yavuzgil 2005 Izmir, Turkey

Random effects model



	MD	95% CI	p-value	Tau ²	Tau	I ²
Omitting Chiurlia 2005 Modena, Italy	-2.54	[-3.78; -1.29]	<0.01	2.4399	1.5620	94%
Omitting Javaroni 2011 Brazil	-2.31	[-3.60; -1.02]	<0.01	2.6811	1.6374	94%
Omitting Kaiser 2004 USA	-1.83	[-2.68; -0.98]	<0.01	0.8581	0.9263	77%
Omitting Kaya 2006 Istanbul, Turkey	-2.11	[-3.24; -0.98]	<0.01	2.0233	1.4224	94%
Omitting Murata 2012 Saitama, Japan	-2.48	[-3.78; -1.18]	<0.01	2.6355	1.6234	94%
Omitting Stuckey 2006 Nedlands, Australia	-2.62	[-3.81; -1.43]	<0.01	2.1185	1.4555	91%
Omitting Uslu 2005 Istanbul, Turkey	-2.35	[-3.62; -1.08]	<0.01	2.6322	1.6224	94%
Omitting Vlachopoulos 2008 Athens, Greece	-2.54	[-3.70; -1.39]	<0.01	2.2000	1.4832	94%
Omitting Yavuzgil 2005 Izmir, Turkey	-2.29	[-3.50; -1.07]	<0.01	2.4667	1.5706	94%
Random effects model	-2.35	[-3.49; -1.22]	<0.01	2.2709	1.5069	93%

Supplement Fig. 6. Sensitivity analysis of studies on the relationship between erectile dysfunction and nitrate-mediated dilation. CI: confidence interval.