# Impact and utility of follicular lymphoma GELF criteria in routine care: an Australasian Lymphoma Alliance study

Allison Barraclough,<sup>1,2</sup> Shivam Agrawal,<sup>1,3</sup> Dipti Talaulikar,<sup>4,5</sup> Geoffrey Chong,<sup>1,6</sup> Edward Yoo,<sup>2,7</sup> Chan Y. Cheah,<sup>7,8</sup> Nunzio Franco,<sup>4,5</sup> Bianca Nguyen,<sup>2</sup> Howard Mutsando,<sup>9,10</sup> Fatima Tahir,<sup>11</sup> Judith Trotman,<sup>11</sup> Jing Huang,<sup>12</sup> Colm Keane,<sup>13</sup> Mitchel Lincoln,<sup>14</sup> Tara Cochrane,<sup>15,16</sup> Anna M. Johnston,<sup>17</sup> Michael Dickinson,<sup>18</sup> Stephen Opat,<sup>12,19</sup> Zoe K. McQuilten,<sup>12,19</sup> Erica M. Wood,<sup>12,19</sup> Gayathri St George<sup>19</sup> and Eliza A. Hawkes<sup>1,19</sup>

<sup>1</sup>Olivia Newton John Cancer Research & Wellness Center, Austin Health, Melbourne, Victoria; <sup>2</sup>Fiona Stanley Hospital, Murdoch, Western Australia; <sup>3</sup>Prince of Wales Hospital, Sydney, New South Wales; <sup>4</sup>Canberra Health Services, Canberra, Australian Capital Territory; <sup>5</sup>College of Health and Medicine, Australian National University, Canberra, Australian Capital Territory; <sup>6</sup>Ballarat Regional Integrated Cancer Center, Ballarat Health Services, Ballarat, Victoria; <sup>7</sup>Sir Charles Gairdner Hospital, Nedlands, Western Australia; <sup>8</sup>Medical School, University of Western Australia, Perth, Western Australia; <sup>9</sup>Toowoomba Hospital, Toowoomba, Queensland; <sup>10</sup>University of Queensland Rural Clinical School, Brisbane, Queensland; <sup>11</sup>Concord Repatriation General Hospital, University of Sydney, Sydney, New South Wales; <sup>12</sup>School of Clinical Sciences at Monash Health, Monash University, Melbourne, Victoria; <sup>13</sup>Princess Alexandra Hospital, Brisbane, Queensland; <sup>14</sup>Alfred Hospital, Melbourne, Victoria; <sup>15</sup>Gold Coast University Hospital, Southport, Queensland; <sup>16</sup>School of Medicine, Griffith University, Griffith, Queensland; <sup>17</sup>Royal Hobart Hospital, Hobart, Tasmania; <sup>18</sup>Peter MacCallum Cancer Center and The Royal Melbourne Hospital, Melbourne, Victoria and <sup>19</sup>School of Public Health and Preventive Medicine, Monash University, Melbourne, Victoria, Australia

**Correspondence:** E. Hawkes eliza.hawkes@onjcri.org.au

Received: October 25, 2023.
Accepted: February 28, 2024.
Early view: March 7, 2024.

https://doi.org/10.3324/haematol.2023.284538

©2024 Ferrata Storti Foundation
Published under a CC BY-NC license

# **Supplementary Tables**

# Supplementary Table S1. Modified GELF criteria by treatment group

		mber of m	mGELF criteria						
Management	0	1	2	3	4	5	6	7	Total
Watch & Wait	20	15	5	2	0	0	1	0	43
Systemic Therapy +/- Radiation	28	34	19	21	11	6	1	1	121
Radiation Alone	3	4	1	1	0	0	0	0	9
Total	51	53	25	24	11	6	2	1	173

### Supplementary Table S1. Modified GELF criteria by treatment group

GELF Groupe d'Etude des Lymphomes Folliculaires

Supplementary Table S2. Distribution of modified GELFc by management group

SPECIFIC GELF CRITERION										
MANAGEMENT STRATEGY	Mass >7cms	≥ 3 sites each >3cm diameter	B Symptoms	Splenomegaly	Compression	Effusion	Leukaemic phase	Cytopenias	LDH>ULN	B2MG>ULN
	(n=47)	(n=30)	(n=29)	(n=19)	(n=22)	(n=12)	(n=7)	(n=3)	(n=43)	(n=56)
Watch & Wait (%)	3 (6)	4 (13)	7 (24)	1 (5)	1 (5)	1 (8)	1 (14)	1 (33)	8 (19)	10 (18)
Systemic Therapy +/- RT (%)	43 (92)	26 (87)	18 (62)	17 (89)	20 (90)	11 (92)	6 (86)	2 (67)	34 (79)	45 (80)
Radiotherapy Alone (%)	1 (2)	0 (0)	4 (14)	1 (5)	1 (5)	0 (0)	0 (0)	0 (0)	1 (2)	1 (2)

## Supplementary Table S2. Distribution of modified GELFc by management group

GELFc=Groupe d'Etude des Lymphomes Folliculaires criteria, n=number, cm=centimetres, RT=Radiotherapy, LDH=lactate dehydrogenase, ULN=upper limit of normal