Antibacterial activity of the structurally novel C-2 amine-substituted

analogues based on quinoxaline

Yuting Liu^{a,b}, Pengju Yang^{a,b}, Yunyun Zhou^c, Zhiwen Zhou^{a,b,*}

^aThird-grade Pharmacological Laboratory on Traditional Chinese Medicine, State Administration of Traditional Chinese Medicine, China Three Gorges University, Yichang, 443002China ^bDepartment of Pharmacy, College of Medicine and Health Sciences, China Three Gorges University, Yichang, 443002China ^cDepartment of Quality Control, China Resources Sanjiu (Huangshi) Medical & Pharmaceutical Co., Ltd. , Huangshi, China

*Corresponding author: E-Mail: zhiwen.zhou@hotmail.com; Tel.: +86-717-6397328 (Z.W. Zhou)



Figure S1. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5a



Figure S2. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5b

Figure S3. ¹H-NMR of compound 5c



Figure S4. ¹³C-NMR spectrum of compound 5d





Figure S5. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5e



Figure S6. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5f



Figure S7. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5g



Figure S8. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5h







Figure S10. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5j



Figure S11. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5k



Figure S12. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5l



Figure S13. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5m



Figure S14. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5n



Figure S15. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 50



Figure S16. ¹H-NMR, ¹³C-NMR and HPLC spectrum of compound 5p