

Table S1. Comparison of the Survived and Deceased Patients in our literature review.

Variables	No. (%) of cases		<i>P</i> value
	Survived group (n=62)	Deceased group (n=6)	
Clinical symptoms on the day of admission			
Cough	32(51.6)	5(83.3)	0.209
Fever	57(91.9)	5(83.3)	0.438
Dyspnea	43(69.4)	5(83.3)	0.662
Headache	16(25.8)	2(33.3)	0.652
Diarrhea	4(6.5)	0(0.0)	1.000
Backache	1(1.6)	0(0.0)	1.000
Myalgia	10(16.1)	1(16.7)	1.000
Chest pain	2(3.2)	0(0.0)	1.000
Confusion	2(3.2)	0(0.0)	1.000
Hemoptysis	4(6.5)	0(0.0)	1.000
Chills	1(1.6)	0(0.0)	1.000
Shiver	5(8.1)	1(16.7)	0.438
Weakness	10(16.1)	0(0.0)	0.581
Poor appetite	10(16.1)	0(0.0)	0.581
Complications			
Hepatic dysfunction	52(83.9)	6(100.0)	0.581
Renal dysfunction	8(12.9)	5(83.3)	0.001
Neurological symptoms	9(14.5)	3(50.0)	0.063
Cardiac damage	17(27.4)	2(33.3)	1.000
Gastrointestinal symptoms	3(4.8)	0(0.0)	1.000
Pulmonary embolism	8(12.9)	0(0.0)	1.000
ECMO	5(8.1)	0(0.0)	1.000