

Supplemental Online Content

Wang L, Zhao Y, Han L, et al. Pharmacist-led management model and medication adherence among patients with chronic heart failure: a randomized clinical trial. *JAMA Netw Open*. 2024;7(12):e2453976. doi:10.1001/jamanetworkopen.2024.53976

eTable 1. The Outcomes of Morisky Scores and MLHFQ Scores

eTable 2. Safety Outcomes

eTable 3. All-Cause Deaths and Unplanned Cardiovascular Hospitalizations During 52-Week Follow-Up

This supplemental material has been provided by the authors to give readers additional information about their work.

eTable 1. The outcomes of Morisky scores and MLHFQ scores

Outcome	Intervention group (n = 223)	Control group (n =222)	P value
Morisky scores, Median (IQR)	7.8 (6.8-8.0)	7.3 (4.5-7.8)	0.001
MLHFQ scores, Median (IQR)	25.0 (12.0-40.0)	35.0 (20.0-44.0)	0.005

Abbreviations:

MLHFQ, Minnesota Living with Heart Failure Questionnaire; IQR, interquartile range.

eTable 2. Safety outcomes

NO.	Intervention group (n = 223)	Control group (n =222)	P value
AEs	19	15	0.48
No AE	204	207	

Abbreviations:AEs, adverse events.

eTable 3. All-cause deaths and unplanned cardiovascular hospitalizations during 52-week follow-up

Events	Intervention group (n = 223)	Control group (n =222)	P value
All-cause deaths, n	4	7	0.36
Unplanned CV hospitalizations, No. %	59 (26.5)	65 (29.2)	0.51
Days lost due to unplanned CV hospitalizations and death, mean (95% CI)	27.3 (13.6-40.9)	33.6 (15.8-51.5)	0.58

Abbreviations: CI, confidence interval.