

Supplemental Table 1. Characteristics of ineligible, pre-ICH depression, and deceased patients

	Ineligible Survivors	Pre-ICH Depression	Deceased Patients
	n= 211	n= 84	n= 2,071
Demographics			
Age at ICH event (median, IQR)	67, 57-77	74, 67-83	77, 69-84
Sex (male, %)	114, 54.0	32, 38.1	1,052, 53.2
Race & Ethnicity (%)			
White	163, 68.2	76, 90.5	1,739, 87.9
Black	17, 8.1	2, 2.4	85, 4.3
Asian	15, 7.1	3, 3.6	84, 4.2
Multiple, Other, and Unknown	16, 7.6	3, 3.6	70, 3.5
Hispanic	24, 11.4	2, 2.4	63, 3.2
Education			
Less than high school	23, 10.9	4, 4.8	71, 3.6
Completed high school	144, 68.2	59, 70.2	1,726, 87.3
Completed college or more	44, 20.9	21, 25.0	180, 9.1
Medical History			
Hypertension	171, 82.2	65, 78.3	1,573, 80.1
Diabetes	35, 16.7	19, 22.9	445, 22.8
Prior ICH	12, 5.7	1, 1.2	129, 6.6
Atrial Fibrillation	32, 15.3	19, 22.9	492, 25.2
Coronary Artery Disease	33, 15.9	20, 24.1	515, 26.4
Pre-ICH Dementia	6, 2.9	12, 14.5	324, 16.7
Neuroimaging Characteristics (median, IQR)			
Lobar Hematoma Location (%)	62, 29.4	54, 64.3	884, 44.7
Hematoma volume	9.8, 3.0-20.0	9.4, 2.6-28.9	23.0, 6.6-58.8
Intraventricular volume	0-0-2.3	0, 0-1.2	0.6, 0-11.7
Acute ICH Characteristics (median, IQR)			
Admission GCS Score	15, 14-15	15, 12-15	12, 6-15
ICH Score	1, 0-2	1, 0-2	2, 2-3
Discharge mRS	4, 3-4	4, 3-5	6, 4-6
Patient Location One Week Post-ICH			
Home	56, 26.5	20, 23.8	133, 6.9
Acute Rehab	66, 31.3	22, 26.2	238, 12.3
Nursing Home	3, 1.4	2, 2.4	34, 1.8
Still in Hospital	86, 40.8	30, 35.7	746, 39.2
Structural Determinants (median, IQR)			
AQI: Median Score	38, 32-39	38, 32-39	38, 32-39
SVI: Overall Percentile Rank	0.23, 0.23-0.46	0.36, 0.27-0.63	0.23, 0.23-0.46