- Enhancing antibody levels and T cell activity of quadrivalent
- 2 influenza vaccine by combining it with CpG HP021
- 3 Jia Ji<sup>1,2</sup>, Lei Chen<sup>3</sup>, Zhigang Wu<sup>1</sup>, Taoming Tang<sup>1</sup>, Linwei Zhu<sup>1</sup>, Miaojin Zhu<sup>1</sup>,
- 4 Yan Chen<sup>4</sup>, Xiangyun Lu<sup>1</sup>, Hangping Yao<sup>1,2</sup>\*
- <sup>5</sup> State Key Laboratory for Diagnosis and Treatment of Infectious Diseases, National
- 6 Clinical Research Center for Infectious Diseases, Collaborative Innovation Center for
- 7 Diagnosis and Treatment of Infectious Diseases, The First Affiliated Hospital, School
- 8 of Medicine, Zhejiang University, Hangzhou, 310003, China.
- <sup>9</sup> Jinan Microecological Biomedicine Shandong Laboratory, Jinan, 250000, China.
- <sup>3</sup> Zhejiang Toyouvax Bio-pharmaceutical Co., Ltd, 311100, Hangzhou, China.
- <sup>4</sup> Jiangsu Taipurui Biotechnology Co., Ltd, 225300, Taizhou, China.
- 13 Jia Ji and Lei Chen contributed equally to this work.

12

\* Correspondence to Hangping Yao, yaohangping@zju.edu.cn

# **Supporting Information**

## Supplementary Table 1. The seroconversion rates of the antibodies after

#### vaccination in mice.


The seroconversion rates of the antibodies (%)							
Groups (No.)		Day 14	<b>Day 21</b>	<b>Day 28</b>	<b>Day 56</b>		
IIV4 antigen (Non-adjuvant)	H1N1	0.0%	75.0%	100.0%	100.0%		
	H3N2	0.0%	50.0%	58.3%	66.7%		
	BV	0.0%	41.7%	91.7%	83.3%		
	BY	0.0%	0.0%	8.3%	33.3%		
IIV4 + 10 μg CpG	H1N1	0.0%	100.0%	100.0%	100.0%		
	H3N2	0.0%	66.7%	77.8%	100.0%		
	BV	83.3%	100.0%	100.0%	100.0%		
	BY	0.0%	55.6%	66.7%	33.3%		
IIV4 + 20 µg CpG	H1N1	16.7%	100.0%	100.0%	100.0%		
	H3N2	0.0%	75.0%	91.7%	66.7%		
	BV	83.3%	100.0%	100.0%	100.0%		
	BY	0.0%	66.7%	91.7%	83.3%		
IIV4 + 40 μg CpG	H1N1	75.0%	100.0%	100.0%	100.0%		
	H3N2	8.3%	91.7%	100.0%	100.0%		
	BV	100.0%	100.0%	100.0%	100.0%		
	BY	0.0%	100.0%	100.0%	100.0%		
IIV4 + 80 μg CpG	H1N1	41.7%	91.7%	100.0%	100.0%		
	H3N2	0.0%	83.3%	91.7%	100.0%		
	BV	91.7%	100.0%	100.0%	100.0%		
	BY	0.0%	83.3%	91.7%	100.0%		
IIV4 + 160 μg CpG	H1N1	75.0%	100.0%	100.0%	100.0%		
	H3N2	0.0%	83.3%	91.7%	100.0%		
	BV	91.7%	100.0%	100.0%	100.0%		
	BY	0.0%	91.7%	91.7%	100.0%		

## 19 Supplementary Table 2. The seroconversion rates of antibodies after the

### 20 vaccination in SD rats.

The seroconversion rates of antibodies (%)						
Groups (No.)		Day 14	<b>Day 21</b>	<b>Day 28</b>		
	H1N1	100%	100%	100%		
IIV4 antigen	H3N2	0%	50%	33%		
(Non-adjuvant)	BV	100%	100%	100%		
	BY	33%	83%	67%		
	H1N1	100%	100%	100%		
6 IIV.4 + 50 CC	H3N2	50%	100%	100%		
6 μg IIV4 + 50 μg CpG	BV	100%	100%	100%		
	BY	100%	100%	100%		
	H1N1	100%	100%	100%		
6 IIVA - 100 G G	H3N2	67%	100%	100%		
6 μg IIV4 + 100 μg CpG	BV	100%	100%	100%		
	BY	100%	100%	100%		
	H1N1	100%	100%	100%		
6 HV4 + 200 G G	H3N2	83%	100%	100%		
6 μg IIV4 + 200 μg CpG	BV	100%	100%	100%		
	BY	100%	100%	100%		
	H1N1	100%	100%	100%		
6 11114 - 400 - 6 6	H3N2	100%	100%	100%		
6 μg IIV4 + 400 μg CpG	BV	100%	100%	100%		
	BY	100%	100%	100%		

21



23 Supplementary Fig 1. Analysis of serum cytokine levels in mice at day 28.