

## CORRECTION

The structure of a glycopeptide purified  
porcine thyroglobulin

By M. FUKUDA and F. EGAMI

Volume 123 (1971)

Scheme 2: Page 419, for  $\alpha$ -NeuNAc-(1 $\rightarrow$ 6)- $\beta$ -D-Gal and  
 $\alpha$ -NeuNGc-(1 $\rightarrow$ 6)- $\beta$ -D-Gal read  $\alpha$ -NeuNAc-(2 $\rightarrow$ 6)- $\beta$ -D-Gal  
and  $\alpha$ -NeuNGc-(2 $\rightarrow$ 6)- $\beta$ -D-Gal respectively

Turnover of protein-bound phosphorylserine in membrane  
preparations from ox brain catalysed by intrinsic kinase and  
phosphatase activity

By M. WELLER and R. RODNIGHT

Volume 124 (1971)

p. 394, col. 2, l. 6: for  $5 \times 10^5$ - $30 \times 10^5$  c.p.m./mol), 1 $\mu$ M-MgSO<sub>4</sub>  
read  $5 \times 10^5$ - $30 \times 10^5$  c.p.m./ $\mu$ mol), 1mM-MgSO<sub>4</sub>

p. 400, Fig. 5 legend, l. 6: for Open symbols represent Na<sup>+</sup>  
and closed symbols K<sup>+</sup> read Open symbols represent K<sup>+</sup> and  
closed symbols Na<sup>+</sup>