

A Munc13/RIM/Rab3 Tripartite Complex: From Priming to Plasticity?

Irina Dulubova¹, Xuelin Lou², Jun Lu¹, Iryna Huryeva¹, Amer Alam¹, Ralf Schneggenburger²,
Thomas C. Südhof³ and Josep Rizo^{1,4}

¹*Departments of Biochemistry and Pharmacology, ³Center for Basic Neuroscience, Department of Molecular Genetics, and Howard Hughes Medical Institute, University of Texas Southwestern Medical Center, 5323 Harry Hines Blvd., Dallas, Texas 75390, USA*

²*Max Planck Institute for Biophysical Chemistry, AG Synaptic Dynamics & Modulation and Dept. of Membrane Biophysics, Am Fassberg 11, 37077 Germany*

⁴ Corresponding author. E-mail: jose@arnie.swmed.edu. Phone: 214-645-6360. FAX: 214-645-6353

Supplementary Fig. 2. Mutations in the RIM2 α ZF domain do not perturb the overall structure of the domain. (A) ¹H-¹⁵N HSQC spectra of the wild type RIM2 α ZF domain (residues 82-142). (B-F) Superpositions of ¹H-¹⁵N HSQC spectrum of the wild type RIM2 α ZF domain (in black) with ¹H-¹⁵N HSQC spectra of the: (B) K₉₇E mutant (in red), (C) K₉₇E / K₉₉E double-point mutant (in green), (D) D₁₀₂K mutant (in yellow), (E) R₁₂₆E mutant (in red), and (F) K₁₂₉D mutant RIM2 α ZF domain (in yellow).

