



# Preventive health care, 1999 update:

## 2. Echocardiography for the detection of a cardiac source of embolus in patients with stroke

Moira K. Kapral,\*† MD; Frank L. Silver,\*‡ MD; with the Canadian Task Force on Preventive Health Care§

### Abstract

**Objective:** To develop guidelines for the use of echocardiography in the investigation of patients with stroke.

**Options:** (1) Routine transthoracic echocardiography (TTE); (2) routine transesophageal echocardiography (TEE); (3) routine TTE followed by TEE if the TTE findings are noncontributory; (4) selective TTE or TEE in patients with cardiac disease who would not otherwise receive anticoagulant therapy.

**Outcomes:** This article reviews the available evidence on the yield of TTE and TEE in detecting cardiac sources of cerebral emboli in patients with stroke, the effectiveness of treatment for cardiac sources of emboli and the effectiveness of screening echocardiography for secondary stroke prevention.

**Evidence:** MEDLINE was searched for relevant articles published from January 1966 to April 1998; also reviewed were additional articles identified from the bibliographies and citations obtained from experts.

**Benefits, harms and costs:** Echocardiography can detect intracardiac masses (thrombus, vegetation or tumour) in about 4% (with TTE) to 11% (with TEE) of stroke patients. The yield is lower among patients without clinical evidence of cardiac disease by history, physical examination, electrocardiography or chest radiography (less than 2%) than among patients with clinical evidence of cardiac disease (less than 19%). The risks of echocardiography to patients are small. TTE has virtually no risks, and TEE is associated with cardiac, pulmonary and bleeding complications in 0.18%. Patients with an identified intracardiac thrombus are at increased risk for embolic events (absolute risk uncertain, range 0%–38%), and this appears to be reduced with anticoagulant therapy (absolute risk reduction uncertain). Anticoagulant therapy carries a risk of major hemorrhage of 1% to 3% per year. The overall effectiveness of echocardiography in the prevention of recurrent stroke is unknown.

**Values:** The strength of evidence was evaluated using the methods of the Canadian Task Force on Preventive Health Care.

**Recommendations:** There is fair evidence to recommend echocardiography in patients with stroke and clinical evidence of cardiac disease by history, physical examination, electrocardiography or chest radiography (grade B recommendation). There is insufficient evidence to recommend for or against TEE in patients with normal results of TTE (grade C recommendation). There is insufficient evidence to recommend for or against routine echocardiography in patients (including young patients) without clinical cardiac disease (grade C recommendation). Routine echocardiography is not recommended for patients with clinical cardiac disease who have independent indications for or contraindications to anticoagulant therapy (grade D recommendation). There is fair evidence to recommend anticoagulant therapy in patients with stroke and intracardiac thrombus (grade B recommendation). There is insufficient

### Education

### Éducation

From \*the Department of Medicine, University of Toronto, Toronto, Ont., and the Divisions of †General Internal Medicine and Clinical Epidemiology and of ‡Neurology, University Health Network, Toronto, Ont.

§The list of task force members appears at the end of the article.

This article has been peer reviewed.

CMAJ 1999;161(8):989-96

‡ See related article page 981



(no) evidence to recommend for or against any specific therapy for patent foramen ovale (grade C recommendation).

**Validation:** The findings of this analysis were reviewed through an iterative process by the members of the Canadian Task Force on Preventive Health Care.

**Sponsors:** The Canadian Task Force on Preventive Health Care is funded through a partnership between the Provincial and Territorial Ministries of Health and Health Canada.

**E**chocardiography uses ultrasonography to provide images of cardiac and great vessel anatomy and blood flow and is commonly used in the investigation of patients with stroke. The purpose of performing echocardiography is to detect cardiac sources of cerebral emboli, thereby allowing the provision of appropriate therapy and preventing recurrent stroke. In this paper we review the available evidence on the yield of echocardiography in patients with stroke, the effectiveness of treatment for cardiac sources of emboli and the effectiveness of screening echocardiography for secondary stroke prevention.

Stroke is the third leading cause of death in Canada. About 200 000 Canadians have experienced stroke, and each year an estimated 50 000 new cases occur in Canadians over 65 years of age.<sup>1</sup> At least 20% of stroke survivors require institutional care, and another 15% require assistance with activities of daily living.<sup>2</sup>

Cardiogenic embolism accounts for 15% to 30% of ischemic strokes.<sup>2,3</sup> Cardioembolic stroke, defined as “the presence of a potential cardioembolic source in the absence of cerebrovascular disease in a patient with nonlacunar stroke,”<sup>3</sup> is a heterogeneous entity, since a variety of cardiac conditions can predispose to cerebral embolism. No “gold standard” exists for the diagnosis, which leads to limitations in accurately determining the frequency and causes of cardioembolic stroke.

Cardiac conditions that are considered potential cardioembolic sources may be classified as major risk, minor risk or uncertain risk (Table 1). Echocardiography is not required for the diagnosis of many of these conditions. For

example, atrial fibrillation, recent myocardial infarction and valvular heart disease are often established diagnoses before the cardioembolic event or may be diagnosed by means of other investigative techniques. For other conditions, the results of echocardiography do not lead to a change in management. For example, patients with stroke and atrial fibrillation or mechanical cardiac valves will usually receive anticoagulant therapy regardless of the findings on echocardiography, whereas those with conditions for which there is no proven effective therapy (e.g., atrial septal aneurysm and aortic arch atheromata) will usually be treated with standard antiplatelet therapy.<sup>7</sup>

In contrast, the detection of a previously undiagnosed intracardiac thrombus, vegetation or tumour will often lead to a change in management. Therefore, this guideline article focuses on the use of screening echocardiography for the detection of intracardiac masses. Although the evidence for the risk of recurrent stroke is less definitive, the detection of patent foramen ovale may also result in a change in therapy, and the role of echocardiography in the management of this condition is also reviewed.

## Methods

We identified relevant articles by a computerized search of MEDLINE (January 1966 to April 1998) using the MeSH terms “cerebrovascular disorders,” “heart diseases,” “echocardiography,” “thromboembolic disorders,” “intracardiac thrombus,” “diagnosis,” “prevention” and “therapy.” A professional librarian assisted with the search. We also performed a manual review of references

**Table 1: Potential cardioembolic sources**

Major risk sources	Embolic rate, %	Minor or uncertain risk sources	Embolic rate, %
Atrial fibrillation	1–12/yr <sup>4-6</sup>	Mitral valve prolapse	< 0.02/yr <sup>3,7</sup>
Intracardiac thrombus	0–35 <sup>3,8-10</sup>	Mitral annular calcification	?
Atrial myxoma	30–40 <sup>11</sup>	Spontaneous echo contrast	?
Mitral stenosis	Normal sinus rhythm: 8–14 <sup>12,13</sup> Atrial fibrillation: 31 Prior embolism: 31–65	Left ventricular aneurysm	< 1 <sup>3,14</sup>
Mechanical valves (anticoagulated)	Aortic: 1.5/yr <sup>3</sup> Mitral: 3/yr	Atrial septal aneurysm	?
Recent myocardial infarction	1–2/yr <sup>15-17</sup>	Calcific aortic stenosis	?
Infective endocarditis	12–40 <sup>18</sup>	Patent foramen ovale	?2/yr <sup>7,19</sup>
Marantic endocarditis	14–90 <sup>20</sup>		
Dilated cardiomyopathy	4/yr <sup>3</sup>		
Aortic arch atheromatous plaques	4–16/yr <sup>7,21-23</sup>		



and obtained citations from experts. Studies were included if they were published in manuscript form in peer-reviewed journals. For the evaluation of the yield of echocardiography, studies involving patients referred for echocardiography (rather than consecutive patients with stroke) were excluded from this review because of the potential for selection bias, as were studies that included patients with systemic (peripheral) emboli (references available from the authors on request).

We systematically reviewed the evidence using the method of the Canadian Task Force on Preventive Health Care (formerly the Canadian Task Force on the Periodic Health Examination).<sup>24</sup> The task force, comprising expert clinician/methodologists from a variety of medical specialties, used a standardized evidence-based method for evaluating the effectiveness of this intervention. A manuscript providing critical appraisal of the evidence was prepared by the lead authors. This included identification and critical appraisal of key studies, and ratings of the quality of this evidence using the task force's established methodologic hierarchy (Appendix 1), resulting in a summary of proposed conclusions and recommendations for consideration by the task force. This manuscript was circulated to the task force members in December 1997, and evidence for this topic was presented by the lead author(s) and deliberated on at a meeting in January 1998.

At the meeting the expert panellists addressed critical issues, clarified ambiguous concepts and analysed the synthesis of the evidence. At the end of this process, the specific clinical recommendations proposed by the lead authors were discussed, as were issues related to clarification of the recommendations for clinical application and any gaps in evidence. The results of this process are reflected in the description of the decision criteria presented with the specific recommendations. The final decisions on recommendations were arrived at unanimously by the group and lead authors.

After the meeting, the lead authors revised the manuscript accordingly. After final revision, the manuscript was sent by the task force to 2 experts in the field (identified by task force members at the meeting). Feedback from these experts was incorporated into a subsequent draft of the manuscript, which was then submitted to the *Canadian Medical Association Journal*.

## Manoeuvres

Transthoracic echocardiography (TTE) is a noninvasive procedure that poses little or no risk to the patient. The diagnostic accuracy of TTE has been evaluated in patients undergoing cardiac surgery or autopsy (level II evidence). For the detection of left ventricular thrombi, TTE has a sensitivity of 86% to 95% and a specificity of

86% to 95%.<sup>25-27</sup> TTE does not reliably visualize the left atrium or left atrial appendage, and it has a sensitivity of only 39% to 63% for the detection of left atrial thrombi and a sensitivity of less than 50% for the diagnosis of patent foramen ovale.<sup>28-32</sup>

Transesophageal echocardiography (TEE) involves the insertion of an ultrasound transducer into the esophagus. The patient is required to fast for at least 4 hours before the procedure. Sedation may be used. The safety of TEE has been evaluated in a multicentre study of 10 419 attempted examinations.<sup>33</sup> Studies were interrupted in 0.6% of cases because of patient intolerance of the TEE probe and in 0.18% of cases because of reversible pulmonary, cardiac or bleeding complications. One death (0.0098%) occurred, in a patient with malignant esophageal infiltration. The diagnostic accuracy of TEE has been evaluated in patients undergoing surgery or autopsy. For the diagnosis of left ventricular thrombus, TEE has a sensitivity and specificity similar to those of TTE.<sup>34</sup> For the diagnosis of left atrial thrombi, TEE has a sensitivity of 100% and a specificity of 99%.<sup>35</sup> For the diagnosis of patent foramen ovale, contrast TEE (involving intravenous administration of agitated saline or other solutions) has a sensitivity of at least 89% and a specificity of 100%.<sup>32,36</sup>

More than 30 cross-sectional studies (level III evidence) have evaluated the yield of TTE or TEE, or both, in detecting intracardiac masses in patients with stroke. In consecutive patients, the yield of echocardiography for the detection of intracardiac masses ranged from 0% to 21%.<sup>37-55</sup> Pooled data from these studies suggest an overall yield of 4% for TTE and 11% for TEE (Table 2). Patent foramen ovale may be detected by contrast TEE in up to 55% of patients with stroke.<sup>56-58</sup>

In some studies the patients were stratified by the presence or absence of clinically apparent cardiac disease, which is typically defined as the presence of myocardial infarction, angina, arrhythmia, valvular heart disease, congestive heart failure or hypertension by history, physical examination, electrocardiography or chest radiography.<sup>37,40-48,51,52,54</sup> Pooled data from these studies suggest that TTE can detect intracardiac masses in 13% and 0.7% of patients with and without clinical cardiac disease respectively; the corresponding values for TEE are 19% and 1.6% (Table 2). Although the prevalence of intracardiac masses in patients without clinical cardiac disease appears to be very low, it should be

**Table 2: Yield of echocardiography in the detection of intracardiac masses in patients with stroke (pooled data)**

Manoeuvre	Yield (and range), %		
	Overall	Patients with cardiac disease*	Patients without cardiac disease
Transthoracic echocardiography (TTE)	4 (0-16)	13 (0-40)	0.7 (0-6)
Transesophageal echocardiography (TEE)	11 (0-21)	19 (10-34)	1.6 (0-4)

\*Defined as myocardial infarction, angina, arrhythmia, valvular disease, congestive heart failure or hypertension by history, physical examination, electrocardiography or chest radiography.



noted that these estimates are based on a secondary analysis of cross-sectional and case-control studies and may be inaccurate.

Three studies, involving a total of 350 patients, addressed the yield of TEE in detecting intracardiac masses in patients with a negative result of TTE.<sup>45,53,54</sup> Inclusion in these studies was also limited to patients without significant carotid stenosis. The pooled data suggest an incremental yield of 17%, predominantly left atrial thrombi. Of note, in the 2 studies that provided data on the presence or absence of known cardiac disease, the yield of TEE in patients with a negative result of TTE and no clinical cardiac disease was 0%, as compared with 18% among patients with cardiac disease.<sup>45,53</sup>

Some authorities suggest that echocardiography is of greater utility in the investigation of younger patients.<sup>59,60</sup> Several investigators have evaluated the yield of echocardiography in patients under the age of 60 years.<sup>41,42,55,61</sup> Potential cardioembolic sources, particularly patent foramen ovale and mitral valve prolapse, were common in this population (prevalence up to 50%), but intracardiac masses were detected less frequently (0%–11%).

## Effectiveness of early detection

Several investigators have assessed the effect of echocardiography on patient management, with inconsistent findings. Two studies (one of TTE only) showed that the results of echocardiography did not alter patient management significantly.<sup>51,62</sup> In one study the authors reported a change in management in 4 (10%) of 39 patients;<sup>63</sup> however, 3 of the 4 had atrial fibrillation and would therefore receive anticoagulant therapy under current standards of care. Two further studies documented a change in management in 5%–10% of patients.<sup>53,64</sup>

A decision analysis of cardiac imaging strategies after stroke showed that echocardiography resulted in improved patient outcomes and that TEE was more cost-effective than TTE.<sup>65</sup> The analysis showed that TEE performed in all patients with stroke cost US\$13 000 per quality-adjusted life-year, whereas TEE performed only in patients with cardiac disease cost US\$9000 per quality-adjusted life-year.

## Effectiveness of treatment

The effectiveness of anticoagulant therapy for intracardiac thrombus has not been properly studied. Observational studies of left ventricular thrombus suggest that the risk of embolization without therapy ranges from 0% to 38% (overall risk 14%).<sup>8,9,66–87</sup> Two small nonblinded randomized trials of warfarin for left ventricular thrombus yielded inconsistent results in the reduction of systemic emboli with anticoagulant therapy, although both documented increased resolution of thrombi.<sup>88,89</sup> Although not specifically designed to address the effectiveness of therapy for intracardiac thrombus, one well-designed randomized trial (level I evidence) of low-dose versus high-dose heparin in patients

with myocardial infarction demonstrated a reduction in thrombus formation of 21% and a subsequent absolute reduction in the risk of systemic emboli of 2.8% with high-dose warfarin.<sup>90</sup> Three well-designed randomized trials (level I evidence) of anticoagulant therapy after myocardial infarction did not evaluate thrombus formation but did document an absolute reduction in the risk of stroke of 1.5% to 3.7% with systemic anticoagulant therapy.<sup>15–17</sup>

Many nonrandomized observational studies (level II/III evidence) have suggested a reduction in the risk of systemic emboli with anticoagulant therapy for left ventricular thrombus.<sup>8,9,66–87</sup> A meta-analysis of 7 studies showed the odds of embolization with anticoagulant therapy to be 0.14 (95% confidence interval 0.04–0.52) compared with no anticoagulant therapy.<sup>10</sup> Of note, most of these studies included patients with recent myocardial infarction, and the results may not be generalizable to all patients with intracardiac thrombi. In addition, therapy for left atrial thrombi has not been studied systematically. Anticoagulant therapy carries a risk of major bleeding of about 1% to 3% per year.<sup>4,91,92</sup>

Treatment options for patent foramen ovale include surgical closure, transcatheter closure, systemic anticoagulant therapy (particularly in the setting of deep vein thrombosis and suspected paradoxical embolus) and antiplatelet therapy.<sup>93–95</sup> However, the relative effectiveness of these therapeutic options has not been studied systematically.

## Recommendations

### *Of the Canadian Task Force on Preventive Health Care (Table 3)*

- There is fair evidence to recommend echocardiography in patients with stroke and clinical evidence of cardiac disease by history, physical examination, electrocardiography or chest radiography (grade B recommendation). TEE is recommended as the preferred initial screening test, based on sensitivity and cost-effectiveness data. There is insufficient evidence to recommend for or against TEE in patients with normal results of TTE (grade C recommendation).
- There is insufficient evidence to recommend for or against routine echocardiography in patients (including young patients) without clinical cardiac disease (grade C recommendation).
- Routine echocardiography is not recommended for patients with clinical cardiac disease who have independent indications for anticoagulant therapy or contraindications to anticoagulant therapy (grade D recommendation). It should be noted, however, that such patients often have non-stroke-related indications for echocardiography.
- There is fair evidence to recommend anticoagulant therapy in patients with stroke and documented intracardiac thrombus (grade B recommendation). There is insufficient (no) evidence to recommend for or against



any specific therapy for patent foramen ovale (grade C recommendation).

### Of other bodies

The American College of Cardiology and the American Heart Association recommend echocardiography in patients with stroke who have clinical evidence of heart disease or who are less than 45 years of age.<sup>59</sup> Routine echocardiography is not recommended in patients over the age of 45 without clinical cardiac disease.

The Cerebral Embolism Task Force recommends “liberal” echocardiography in patients with stroke in whom cerebrovascular mechanisms are deemed unlikely.<sup>3</sup>

The Ad Hoc Committee on Guidelines for the Manage-

ment of Transient Ischemic Attacks for the Stroke Council of the American Heart Association recommends TTE only in patients with clinical evidence of cardiac disease or young subgroups without major risk factors for primary cerebrovascular disease, and possibly for those in whom no source of transient ischemic attack has been identified after other tests are completed.<sup>96</sup> TEE is not recommended in unselected patients.

### Research agenda

- More data are required on the yield of echocardiography in subgroups of patients at low risk.
- The incremental yield of TEE in patients with normal findings on TTE needs further study.

**Table 3: Recommendations of the Canadian Task Force on Preventive Health Care — screening and therapy for potential cardioembolic sources in patients with stroke**

Manoeuvre	Effectiveness	Level of evidence*	Recommendation*
<b>Echocardiography (TTE or TEE for detection of intracardiac masses)</b>	TTE is safe and has a sensitivity and specificity of 86%–95% for the detection of left ventricular thrombus. Sensitivity for detection of left atrial thrombus is 39%–63%. TEE has a 0.2% risk of adverse effects, and sensitivity of 100% and specificity of 99% for the detection of left atrial thrombi		
Patients with clinical cardiac disease† and no pre-existing indications for anticoagulation	Screening yield is about 13% (TTE) or 19% (TEE). The effect of screening on patient management and outcomes is unknown	Case-control (II-2) <sup>42,51,54</sup> and cross-sectional (III) studies <sup>37,40,41,43–48,52</sup>	Fair evidence to recommend echocardiography for the detection of intracardiac masses in patients with clinical cardiac disease and no pre-existing indications for anticoagulation (grade B).‡ TEE is the preferred initial screening test
Patients with pre-existing indications for anticoagulation or contraindications to anticoagulation§	Screening rarely results in a change in management	Case-control (II-2) and cross-sectional (III) studies <sup>51,62,63</sup>	Fair evidence to recommend against routine echocardiography for patients with pre-existing indications for anticoagulation or contraindications to anticoagulation (grade D)
Patients without clinical cardiac disease	Screening yield is less than 2% in patients aged 45 years or more, and unknown in patients aged less than 45 without clinical cardiac disease	Case-control (II-2) and cross-sectional (III) studies <sup>37,40–48,51,52,54,55,61</sup>	Insufficient evidence to recommend for or against echocardiography in patients without clinical cardiac disease (grade C)
<b>Anticoagulation (warfarin) for intracardiac thrombus to prevent systemic emboli</b>	Anticoagulation for intracardiac thrombus decreases the risk of systemic emboli and stroke, although the magnitude of treatment benefit is unknown. The annual risk of major hemorrhage with systemic anticoagulation is between 1% and 3%	Small flawed randomized controlled trials (II-1), <sup>88,89</sup> case-control studies (II-2) and case series (III). <sup>8,9,66–87</sup> Randomized controlled trials of anticoagulation after myocardial infarction (I) <sup>15–17,90</sup>	Fair evidence to recommend warfarin therapy in patients with stroke and intracardiac thrombus (grade B)
<b>Treatment for patent foramen ovale</b>	Treatment options include surgical closure, transcatheter closure, systemic anticoagulation and antiplatelet therapy	Treatment options have not been systematically evaluated	Insufficient (no) evidence to recommend for or against any specific therapy for patent foramen ovale (grade C)

\*See Appendix 1 for definitions of the levels of evidence and grades of recommendations.

†As defined in footnote to Table 2.

‡This recommendation is based on positive test characteristics, prevalence and the availability of effective treatment.

§Such patients may have other indications for echocardiography.



- Further research is needed to determine whether screening echocardiography results in decreased risk of recurrent stroke and improved patient outcomes.
- Further research is required into the optimal management of conditions such as patent foramen ovale and aortic arch atheromata. As this information becomes available, the benefit of screening echocardiography will need to be re-evaluated.

We thank Dr. Henry J.M. Barnett, Robarts Research Institute, University of Western Ontario, London, Ont., and Dr. Jay P. Mohr, Neurological Institute, Columbia University, New York, for reviewing a draft form of this report. The views expressed in this report are those of the authors and the Canadian Task Force on Preventive Health Care and do not necessarily reflect the positions of reviewers.

Dr. Kapral held a Research Fellowship from the Heart and Stroke Foundation of Canada at the time of authorship.

Competing interests: None declared.

## References

1. Statistics Canada, Health Canada, Heart and Stroke Foundation of Canada. *Heart disease and stroke in Canada*. Toronto: Heart and Stroke Foundation of Canada; 1997.
2. Bonita R. Epidemiology of stroke. *Lancet* 1992;339:342-4.
3. Cerebral Embolism Task Force. Cardiogenic brain embolism. The second report of the Cerebral Embolism Task Force. *Arch Neurol* 1989;46:727-43.
4. Atrial Fibrillation Investigators. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation: analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994;154:1449-57.
5. EAFT Study Group. European Atrial Fibrillation Trial: secondary prevention of vascular events in patients with nonrheumatic atrial fibrillation and a recent transient ischaemic attack or minor ischaemic stroke. *Lancet* 1993;342:1255-62.
6. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke* 1991;22:983-8.
7. Manning WJ. Role of transesophageal echocardiography in the management of thromboembolic stroke. *Am J Cardiol* 1997;80:19D-28D.
8. Visser CA, Kan G, Meltzer RS, Lie KI, Durrer D. Long-term follow-up of left ventricular thrombus after acute myocardial infarction. A two-dimensional echocardiographic study in 96 patients. *Chest* 1984;86:532-6.
9. Stratton J, Resnick AD. Increased embolic risk in patients with left ventricular thrombi. *Circulation* 1987;75:1004-11.
10. Vaitkus P, Barnathan E. Embolic potential, prevention and management of mural thrombus complicating anterior myocardial infarction: a meta-analysis. *J Am Coll Cardiol* 1993;22:1004-9.
11. Reynen K. Cardiac myxomas [review]. *N Engl J Med* 1995;333:1610-7.
12. Szekeley P. Systemic embolism and anticoagulant prophylaxis in rheumatic heart disease. *BMJ* 1964;1:209-12.
13. Deverall PB, Olley PM, Smith DR, Watson DA, Whitaker W. Incidence of systemic embolism before and after mitral valvotomy. *Thorax* 1968;23:530-6.
14. Lapeyre AC III, Steele PM, Kazmier FJ, Chesebro JH, Vlietstra RE, Fuster V. Systemic embolism in chronic left ventricular aneurysm: incidence and the role of anticoagulation. *J Am Coll Cardiol* 1985;6:534-8.
15. Effect of long-term oral anticoagulant treatment on mortality and cardiovascular morbidity after myocardial infarction. Anticoagulants in the Secondary Prevention of Events in Coronary Thrombosis (ASPECT) Research Group. *Lancet* 1994;343:499-503.
16. Smith P, Arnesen H, Holme I. The effect of warfarin on mortality and reinfarction after myocardial infarction. *N Engl J Med* 1990;323:147-52.
17. Loh E, Sutton MS, Wun CC, Rouleau JL, Flaker GC, Gottlieb SS, et al. Ventricular dysfunction and the risk of stroke after myocardial infarction. *N Engl J Med* 1997;336:251-7.
18. Sherman DG, Dyken ML, Gent M, Harrison MJG, Hart RG, Mohr JP. Antithrombotic therapy for cerebrovascular disorders. An update. *Chest* 1995; 108:444S-456S.
19. Bogousslavsky J, Garazi S, Jeanrenaud X, Aebischer N, Van Melle G. Stroke recurrence in patients with patent foramen ovale: the Lausanne Study. Lausanne Stroke with Paradoxical Embolism Study Group. *Neurology* 1996;46:1301-5.
20. Lopez JA, Ross RS, Fishbein MC, Siegel RJ. Nonbacterial thrombotic endocarditis: a review. *Am Heart J* 1987;113:773-84.
21. French Study of Aortic Plaques in Stroke Group. Atherosclerotic disease of the aortic arch as a risk factor for recurrent ischemic stroke. *N Engl J Med* 1996;334:1216-21.
22. Mitusch R, Doherty C, Wucherpfennig H, Memmesheimer C, Tepe C, Stierle U, et al. Vascular events during follow-up in patients with aortic arch atherosclerosis. *Stroke* 1997;28:36-9.
23. Witteman JC, Kannel WB, Wolf PA, Grobbee DE, Hofman A, D'Agostino RB, et al. Aortic calcified plaques and cardiovascular disease (the Framingham Study). *Am J Cardiol* 1990;66:1060-4.
24. Woolf SH, Battista RN, Anderson GM, Logan AG, Wang E, and other members of the Canadian Task Force on the Periodic Health Examination. Assessing the clinical effectiveness of preventive maneuvers: analytic principles and systematic methods in reviewing evidence and developing clinical practice recommendations. *J Clin Epidemiol* 1990;43:891-905.
25. Visser CA, Kan G, David GK, Lie KI, Durrer D. Two dimensional echocardiography in the diagnosis of left ventricular thrombus. A prospective study of 67 patients with anatomic validation. *Chest* 1983;83:228-32.
26. Stratton JR, Lighty GW, Pearlman AS, Ritchie JL. Detection of left ventricular thrombus by two-dimensional echocardiography: sensitivity, specificity, and causes of uncertainty. *Circulation* 1982;66:156-66.
27. Starling MR, Crawford MH, Sorensen GS, Grover FL. Comparative value of invasive and noninvasive techniques for identifying left ventricular mural thrombi. *Am Heart J* 1983;106:1143-9.
28. Schweizer P, Bardos P, Erbel R. Detection of left atrial thrombi by echocardiography. *Br Heart J* 1981;45:148-56.
29. Shrestha NK, Moreno FL, Narciso FV, Torres L, Calleja HB. Two-dimensional echocardiographic diagnosis of left-atrial thrombus in rheumatic heart disease. A clinicopathological study. *Circulation* 1983;67:341-7.
30. DePace NL, Soulen RL, Kotler MN, Mintz GS. Two dimensional echocardiographic detection of intraatrial masses. *Am J Cardiol* 1981;48:954-60.
31. Di Tullio MD, Sacco RL, Venketasubramanian N, Sherman D, Mohr JP, Homma S. Comparison of diagnostic techniques for the detection of a patent foramen ovale in stroke patients. *Stroke* 1993;24:1020-4.
32. Chen WJ, Kuan P, Lien WP, Lin FY. Detection of patent foramen ovale by contrast transesophageal echocardiography. *Chest* 1992;101:1515-20.
33. Daniel WG, Erbel R, Kasper W, Visser CA, Engberding R, Sutherland GR, et al. Safety of transesophageal echocardiography. A multicenter survey of 10,419 examinations. *Circulation* 1991;83:817-21.
34. Daniel WG, Mugge A. Transesophageal echocardiography. *N Engl J Med* 1995;332:1268-78.
35. Manning WJ, Weintraub RM, Waksmonski CA, Haering JM, Rooney PS, Maslow AD, et al. Accuracy of transesophageal echocardiography for identifying left atrial thrombi. A prospective, intraoperative study. *Ann Intern Med* 1995;123:817-22.
36. Schneider B, Zienkiewicz T, Jansen V, Hofmann T, Noltenius H, Meinertz T. Diagnosis of patent foramen ovale by transesophageal echocardiography and correlation with autopsy findings. *Am J Cardiol* 1996;77:1202-9.
37. Greenland P, Knopman DS, Mikell FL, Asinger RW, Anderson DC, Good DC. Echocardiography in diagnostic assessment of stroke. *Ann Intern Med* 1981;95:51-3.
38. Nishide M, Irino T, Gotoh M, Naka M, Tsuji K. Cardiac abnormalities in ischemic cerebrovascular disease studied by two-dimensional echocardiography. *Stroke* 1983;14:541-5.
39. Rem JA, Hachinski VC, Boughner DR, Barnett HJM. Value of cardiac monitoring and echocardiography in TIA and stroke patients. *Stroke* 1985;16: 950-6.
40. Good D, Frank S, Verhulst S, Sharma B. Cardiac abnormalities in patients with negative arteriograms. *Stroke* 1986;17:6-11.
41. Fogelholm R, Melin J. Echocardiography in ischemic cerebrovascular disease. *BMJ* 1987;295:305-6.
42. Zenker G, Erbel R, Kramer G, Mohr-Kahaly S, Drexler M, Harmoncourt K, et al. Transesophageal two-dimensional echocardiography in young patients with cerebral ischemic events. *Stroke* 1988;19:345-8.
43. Pop G, Sutherland GR, Koudstaal PJ, Sit TW, de Jong G, Roelandt JR. Transesophageal echocardiography in the detection of intracardiac embolic sources in patients with transient ischemic attacks. *Stroke* 1990;21:560-5.
44. Pearson AC, Labovitz AJ, Tatineni S, Gomez CR. Superiority of transesophageal echocardiography in detecting cardiac source of embolism in patients with cerebral ischemia of uncertain etiology. *J Am Coll Cardiol* 1991;17:66-72.
45. Lee RJ, Bartzokis T, Yeoh TK, Grogin HR, Choi D, Schnittger I. Enhanced detection of intracardiac sources of cerebral emboli by transesophageal echocardiography. *Stroke* 1991;22:734-9.
46. Voyce SJ, Aurigemma GP, Dahlberg S, Orsinelli D, Pape LA, Sweeney A, et al. A comparison of two-dimensional echocardiography vs carotid duplex scanning in older patients with cerebral ischemia. *Arch Intern Med* 1992;152: 2089-93.
47. Comess KA, DeRook FA, Beach KW, Lytle NJ, Golby AJ, Albers GW. Transesophageal echocardiography and carotid ultrasound in patients with cerebral ischemia: prevalence of findings and recurrent stroke risk. *J Am Coll Cardiol* 1994;23:1598-603.



48. Albers GW, Comess KA, DeRook FA, Bracci P, Atwood JE, Bolger A, et al. Transesophageal echocardiographic findings in stroke subtypes. *Stroke* 1994;25:23-8.
49. Shyu KG, Chen JJ, Huang ZS, Hwang JJ, Lee TK, Kuan P, et al. Role of transesophageal echocardiography in the diagnostic assessment of cardiac sources of embolism in patients with acute ischemic stroke. *Cardiology* 1994;85:53-60.
50. Lindgren A, Roijer A, Norrving B, Wallin L, Eskilsson J, Johansson BB. Carotid artery and heart disease in subtypes of cerebral infarction. *Stroke* 1994;25:2356-62.
51. Sansoy V, Abbott RD, Jayaweera AR, Kaul S. Low yield of transthoracic echocardiography for cardiac source of embolism. *Am J Cardiol* 1995;75:166-9.
52. Stollberger C, Brainin M, Abzieher F, Slany J. Embolic stroke and transesophageal echocardiography: Can clinical parameters predict the diagnostic yield? *J Neurol* 1995;242:437-42.
53. Rauh G, Fischereder M, Spengel FA. Transesophageal echocardiography in patients with focal cerebral ischemia of unknown cause. *Stroke* 1996;27:691-4.
54. Labovitz AJ, Camp A, Castello R, Martin TJ, Ofili EO, Rickmeyer N, et al. Usefulness of transesophageal echocardiography in unexplained cerebral ischemia. *Am J Cardiol* 1993;72:1448-52.
55. Nighoghossian N, Perinetti M, Barthelet M, Adeleine P, Trouillas P. Transesophageal echocardiography in patients less than 60 years of age without obvious cardiac source of embolism. *Neurol Res* 1995;17:368-72.
56. Nighoghossian N, Perinetti M, Barthelet M, Adeleine P, Trouillas P. Potential cardioembolic sources of stroke in patients less than 60 years of age. *Eur Heart J* 1996;17:590-4.
57. Lechat P, Mas JL, Lascault G, Loron P, Theard M, Klimczak M, et al. Prevalence of patent foramen ovale in patients with stroke. *N Engl J Med* 1988;318:1148-52.
58. Ranoux D, Cohen A, Cabanes L, Amarenco P, Bousser MG, Mas JL. Patent foramen ovale: Is stroke due to paradoxical embolism? *Stroke* 1993;24:31-4.
59. American College of Cardiology/American Heart Association Task Force on Assessment of Diagnostic and Therapeutic Cardiovascular Procedures. ACC/AHA guidelines for the clinical application of echocardiography. *Circulation* 1990;82:2323-45.
60. Chambers JB, de Belder MA, Moore D. Echocardiography in stroke and transient ischemic attack. *Heart* 1997;78(Suppl 1):2-6.
61. Biller J, Johnson MR, Adams HP Jr, Kerber RE, Toffol GJ, Butler MJ. Echocardiographic evaluation of young adults with nonhemorrhagic cerebral infarction. *Stroke* 1986;17:608-12.
62. Robbins JA, Sagar KB, French M, Smith PJ. Influence of echocardiography on management of patients with systemic emboli. *Stroke* 1983;14:546-9.
63. Hata JS, Ayres RW, Biller J, Adams HP Jr, Stuhlmuller JE, Burns TL, et al. Impact of transesophageal echocardiography on the anticoagulation management of patients admitted with focal cerebral ischemia. *Am J Cardiol* 1993;72:707-10.
64. Cujec B, Polasek P, Voll C, Shuaib A. Transesophageal echocardiography in the detection of potential cardiac sources of embolism in stroke patients. *Stroke* 1991;22:727-33.
65. McNamara RL, Lima JAC, Whelton PK, Powe NR. Echocardiographic identification of cardiovascular sources of emboli to guide clinical management of stroke: a cost-effectiveness analysis. *Ann Intern Med* 1997;127:775-87.
66. Simpson MT, Oberman A, Kouchoukos NT, Rogers WJ. Prevalence of mural thrombi and systemic embolization with left ventricular aneurysm. Effect of anticoagulant therapy. *Chest* 1980;77:463-9.
67. Reeder GS, Lengyel M, Tajik AJ, Seward JB, Smith HC, Danielson GK. Mural thrombus in left ventricular aneurysm: incidence, role of angiography, and relation between anticoagulation and embolization. *Mayo Clin Proc* 1981;56:77-81.
68. Asinger RW, Mikell FL, Elsparger J, Hodges M. Incidence of left-ventricular thrombus after acute transmural myocardial infarction. *N Engl J Med* 1981;305:297-302.
69. Keating EC, Gross SA, Schlamowitz RA, Glassman J, Mazur JH, Pitt WA, et al. Mural thrombi in myocardial infarctions. Prospective evaluation by two-dimensional echocardiography. *Am J Med* 1983;74:989-95.
70. Gottdiener JS, Gay JA, VanVoorhees L, DiBianco R, Fletcher RD. Frequency and embolic potential of left ventricular thrombus in dilated cardiomyopathy: assessment by 2-dimensional echocardiography. *Am J Cardiol* 1983;52:1281-5.
71. Stratton JR, Ritchie JL. The effects of antithrombotic drugs in patients with left ventricular thrombi: assessment with indium-111 platelet imaging and two-dimensional echocardiography. *Circulation* 1984;69:561-8.
72. Weinrich DJ, Burke JF, Pauletto FJ. Left ventricular mural thrombi complicating acute myocardial infarction. Long-term follow-up with serial echocardiography. *Ann Intern Med* 1984;100:789-94.
73. Haugland JM, Asinger RW, Mikell FL, Elsparger J, Hodges M. Embolic potential of left ventricular thrombi detected by two-dimensional echocardiography. *Circulation* 1984;70:588-98.
74. Visser CA, Kan G, Meltzer RS, Dunning AJ, Roelandt J. Embolic potential of left ventricular thrombus after myocardial infarction: a two-dimensional echocardiographic study of 119 patients. *J Am Coll Cardiol* 1985;5:1276-80.
75. Spirito P, Bellotti P, Chiarella F, Domenicucci S, Sementa A, Vecchio C. Prognostic significance and natural history of left ventricular thrombi in patients with acute anterior myocardial infarction: a two-dimensional echocardiographic study. *Circulation* 1985;72:774-80.
76. Davis MJE, Ireland MA. Effect of early anticoagulation on the frequency of left ventricular thrombi after anterior wall acute myocardial infarction. *Am J Cardiol* 1986;57:1244-7.
77. Johannessen KA. Peripheral emboli from left ventricular thrombi of different echocardiographic appearance in acute myocardial infarction. *Arch Intern Med* 1987;147:641-4.
78. Stratton JR, Nemanich JW, Johannessen KA, Resnick AD. Fate of left ventricular thrombi in patients with remote myocardial infarction or idiopathic cardiomyopathy. *Circulation* 1988;78:1388-93.
79. Johannessen KA, Nordrehaug JE, von der Lippe G, Vollset SE. Risk factors for embolisation in patients with left ventricular thrombi and acute myocardial infarction. *Br Heart J* 1988;60:104-10.
80. Nihoyannopoulos P, Smith GC, Maseri A, Foale RA. The natural history of left ventricular thrombus in myocardial infarction: a rationale in support of masterly inactivity. *J Am Coll Cardiol* 1989;14:903-11.
81. Jugdutt BI, Sivaram CA. Prospective two-dimensional echocardiographic evaluation of left ventricular thrombus and embolism after acute myocardial infarction. *J Am Coll Cardiol* 1989;13:554-64.
82. Kupper AJ, Verheugt FW, Peels CH, Galema TW, Roos JP. Left ventricular thrombus incidence and behavior studied by serial two-dimensional echocardiography in acute anterior myocardial infarction: left ventricular wall motion, systemic embolism and oral anticoagulation. *J Am Coll Cardiol* 1989;13:1514-20.
83. Ciaccheri M, Castelli G, Cecchi F, Nannini M, Santoro G, Troiani V, et al. Lack of correlation between intracavitary thrombosis detected by cross sectional echocardiography and systemic emboli in patients with dilated cardiomyopathy. *Br Heart J* 1989;62:26-9.
84. Keren A, Goldberg S, Gottlieb S, Klein J, Schuger C, Medina A, et al. Natural history of left ventricular thrombi: their appearance and resolution in the posthospitalization period of acute myocardial infarction. *J Am Coll Cardiol* 1990;15:790-800.
85. Falk RH, Foster E, Coats MH. Ventricular thrombi and thromboembolism in dilated cardiomyopathy: a prospective follow-up study. *Am Heart J* 1992;123:136-42.
86. Katz SD, Marantz PR, Biasucci L, Jondeau G, Lee K, Brennan C, et al. Low incidence of stroke in ambulatory patients with heart failure: a prospective study. *Am Heart J* 1993;126:141-6.
87. Cregler LL. Antithrombotic therapy in left ventricular thrombosis and systemic embolism. *Am Heart J* 1992;123:1110-4.
88. Kouvaras G, Chronopoulos G, Soufras G, Sofronas G, Solomos D, Bakirtzis A, et al. The effects of long-term antithrombotic treatment on left ventricular thrombi in patients after an acute myocardial infarction. *Am Heart J* 1990;119:73-8.
89. Tramarin R, Pozzoli M, Febo O, Opasich C, Colombo E, Cobelli F, et al. Two-dimensional echocardiographic assessment of anticoagulant therapy in left ventricular thrombosis early after acute myocardial infarction. *Eur Heart J* 1986;7:482-92.
90. Turpie AG, Robinson JG, Doyle DJ, Mulji AS, Mishkel GJ, Sealey BJ, et al. Comparison of high-dose with low-dose subcutaneous heparin to prevent left ventricular mural thrombosis in patients with acute transmural anterior myocardial infarction. *N Engl J Med* 1989;320:352-7.
91. Albers GW. Atrial fibrillation in stroke: three new studies, three remaining questions. *Arch Intern Med* 1994;154:1443-8.
92. European Atrial Fibrillation Study Group. Optimal oral anticoagulant therapy in patients with nonrheumatic atrial fibrillation and recent cerebral ischemia. *N Engl J Med* 1995;333:5-10.
93. Devuyst G, Bogousslavsky J, Ruchat P, Jeanrenaud X, Despland PA, Regli F, et al. Prognosis after stroke followed by surgical closure of patent foramen ovale: a prospective follow-up study with brain MRI and simultaneous transesophageal and transcranial Doppler ultrasound. *Neurology* 1996;47:1162-6.
94. Homma S, Di Tullio MR, Sacco RL, Sciaccia RR, Smith C, Mohr JP. Surgical closure of patent foramen ovale in cryptogenic stroke patients. *Stroke* 1997;28:2376-81.
95. Bogousslavsky J, Devuyst G, Nendaz M, Yamamoto H, Sarasin F. Prevention of stroke recurrence with presumed paradoxical embolism. *J Neurol* 1997;244:71-5.
96. Guidelines for the management of transient ischemic attacks. From the Ad Hoc Committee on Guidelines for the Management of Transient Ischemic Attacks of the Stroke Council of the American Heart Association. *Stroke* 1994;25:1320-35.

**Reprint requests to:** Canadian Task Force on Preventive Health Care, Parkwood Hospital, 801 Commissioners Rd. E, London ON N6C 5J1; [ctf@ctfphc.org](mailto:ctf@ctfphc.org)



**Members of the Canadian Task Force on Preventive Health Care**

**Chairman:** Dr. John W. Feightner, Professor, Department of Family Medicine, University of Western Ontario, London, Ont.  
**Past Chairman:** Dr. Richard Goldbloom, Professor, Department of Pediatrics, Dalhousie University, Halifax, NS. **Members:** Drs. R. Wayne Elford, Professor and Chair of Research, Department of Family Medicine, University of Calgary, Calgary, Alta.; Michel Labrecque, Associate Professor and Director of Research, Department of Family Medicine and Centre hospitalier universitaire de Québec, Laval University, Sainte-Foy, Que.; Robin S. McLeod, Professor, Department of Surgery, Mount Sinai Hospital and University of Toronto, Toronto; Harriet MacMillan, Associate Professor, Departments of Psychiatry and Pediatrics and Centre for Studies of Children at Risk, McMaster University, Hamilton, Ont.; Jean-Marie Moutquin, Professor, Department of Obstetrics and Gynecology and Saint-François d'Assise Research Centre, Laval University, Sainte-Foy, Que.; Christopher Patterson, Professor and Head, Division of Geriatric Medicine, Department of Medicine, McMaster University, Hamilton, Ont.; and Elaine E.L. Wang, Associate Professor, Departments of Pediatrics and Public Health Sciences, Faculty of Medicine, University of Toronto, Toronto. **Resource people:** Nadine Wathen, Coordinator, and Tim Pauley, Research Assistant, Canadian Task Force on Preventive Health Care, Department of Family Medicine, University of Western Ontario, London, Ont.

**Appendix 1: Canadian Task Force on Preventive Health Care levels of evidence and grades of recommendations**

**Levels of evidence**

- I Evidence from at least one well-designed randomized controlled trial
- II-1 Evidence from well-designed controlled trials without randomization
- II-2 Evidence from well-designed cohort or case-control analytic studies, preferably from more than one centre or research group
- II-3 Evidence from comparisons between times or places with or without the intervention; dramatic results from uncontrolled studies could be included here
- III Opinions of respected authorities, based on clinical experience; descriptive studies or reports of expert committees

**Grades of recommendations**

- A Good evidence to support the recommendation that the condition or manoeuvre be specifically considered in a periodic health examination (PHE)
- B Fair evidence to support the recommendation that the condition or manoeuvre be specifically considered in a PHE
- C Insufficient evidence regarding inclusion or exclusion of the condition or manoeuvre in a PHE, but recommendations may be made on other grounds
- D Fair evidence to support the recommendation that the condition or manoeuvre be specifically excluded from a PHE
- E Good evidence to support the recommendation that the condition or manoeuvre be specifically excluded from a PHE

# Manuscripts invited


The *Canadian Medical Association Journal* invites manuscript submissions from authors interested in wide exposure among Canadian physicians and around the world.

- Most widely circulated Canadian medical journal — almost 60 000 copies
- Highest readership of any scholarly journal in Canada
- Quality peer review
- Prompt editorial decision-making
- Fast-track review for articles of public health importance
- Indexed in MEDLINE, EMBASE/Excerpta Medica, *Current Contents: Life Sciences*, *Current Contents: Clinical Medicine* and other abstracting/indexing services
- Table of contents, abstracts and selected articles on *CMA Online*: [www.cma.ca/cmaj](http://www.cma.ca/cmaj)
- Full text available through numerous electronic document delivery services and CD-ROM

• Biweekly media coverage coordinated by CMA  
Guidelines for authors are available on *CMA Online*  
([www.cma.ca/cmaj/author.htm](http://www.cma.ca/cmaj/author.htm)).

**To discuss submission of a manuscript, contact:**

John Hoey, MD  
Editor-in-Chief, *CMAJ*  
hoeyj@cma.ca  
tel 800 663-7336 x2118  
(in North America)  
tel 613 731-8610 x2118  
(from other points)  
fax 613 565-2382



ASSOCIATION  
MÉDICALE  
CANADIENNE



CANADIAN  
MEDICAL  
ASSOCIATION