

His · · · Asp Catalytic Dyad of Ribonuclease A: Histidine pK_a Values in the Wild-Type, D121N, and D121A Enzymes

David J. Quirk and Ronald T. Raines

Departments of Biochemistry and Chemistry, University of Wisconsin–Madison, Madison, Wisconsin 53706 USA

ABSTRACT Bovine pancreatic ribonuclease A (RNase A) has a conserved His · · · Asp catalytic dyad in its active site. Structural analyses had indicated that Asp¹²¹ forms a hydrogen bond with His¹¹⁹, which serves as an acid during catalysis of RNA cleavage. The enzyme contains three other histidine residues including His¹², which is also in the active site. Here, ¹H-NMR spectra of wild-type RNase A and the D121N and D121A variants were analyzed thoroughly as a function of pH. The effect of replacing Asp¹²¹ on the microscopic pK_a values of the histidine residues is modest: none change by more than 0.2 units. There is no evidence for the formation of a low-barrier hydrogen bond between His¹¹⁹ and either an aspartate or an asparagine residue at position 121. In the presence of the reaction product, uridine 3'-phosphate (3'-UMP), protonation of one active-site histidine residue favors protonation of the other. This finding is consistent with the phosphoryl group of 3'-UMP interacting more strongly with the two active-site histidine residues when both are protonated. Comparison of the titration curves of the unliganded enzyme with that obtained in the presence of different concentrations of 3'-UMP shows that a second molecule of 3'-UMP can bind to the enzyme. Together, the data indicate that the aspartate residue in the His · · · Asp catalytic dyad of RNase A has a measurable but modest effect on the ionization of the adjacent histidine residue.

INTRODUCTION

RNase A (EC 3.1.27.5) has been the object of much landmark work in biochemistry and biophysics (Raines, 1998). For example, RNase A was the first enzyme and second protein (after insulin) for which a complete amino acid sequence was determined (Smythe et al., 1963), and the third enzyme and fourth protein (after myoglobin, lysozyme, and carboxypeptidase) whose structure was solved by x-ray diffraction analysis (Kartha et al., 1967). The use of NMR spectroscopy in elaborating protein structure (Saunders et al., 1957), histidine pK_a values [for a compilation, see Antosiewicz et al. (1996)], and protein folding pathways (Udgaonkar and Baldwin, 1988) was developed with RNase A. The ¹H-NMR resonances of the enzyme have been assigned, and the structure of the enzyme in solution has been determined (Rico et al., 1989; Robertson et al., 1989). This wealth of information has made RNase A an ideal model system for detailed biophysical analyses of protein structure–function relationships.

RNase A is a small protein (124 amino acid residues; 13.7 kDa) that catalyzes the hydrolysis of RNA in two distinct steps. In the first step, the side chain of His¹² acts as a base to abstract a proton from the 2'-hydroxyl group of a substrate molecule and thereby facilitate attack on the phosphorus atom. The side chain of His¹¹⁹ acts as an acid to

protonate the 5'-oxygen and facilitate its displacement (Findlay et al., 1961; Thompson and Raines, 1994). Both products are then released to solvent. The slow hydrolysis of the 2',3'-cyclic phosphodiester occurs in a separate step that resembles the reverse of transphosphorylation (Cuchillo et al., 1993; Thompson et al., 1994).

RNase A may have converged upon a catalytic dyad that is similar to the catalytic triad of serine proteases and other hydrolases. Joint x-ray/neutron diffraction analysis indicates that His¹¹⁹ forms a hydrogen bond with Asp¹²¹ (Fig. 1). The importance of this hydrogen bond can be inferred from the conservation of His¹¹⁹ and Asp¹²¹ in all of the over 40 homologous pancreatic ribonucleases of known sequence (Beintema, 1987; Beintema et al., 1988).

We are interested in structure–function relationships that involve the His · · · Asp catalytic dyad of RNase A. Previously, we reported that replacing Asp¹²¹ with an asparagine or alanine residue has no effect on the overall three-dimensional structure of the enzyme, and a significant but not substantial effect on catalysis (Schultz et al., 1998). In addition, replacing Asp¹²¹ with an asparagine or alanine residue results in a loss of conformational stability at pH 6.0 of $\Delta\Delta G^\circ = 2.0$ kcal/mol, from a total of 9.0 kcal/mol (Quirk et al., 1998). The magnitude of this loss is similar to that to transition state binding during catalysis.

Here, we have used site-directed mutagenesis and ¹H-NMR spectroscopy to gain additional insight into the role of Asp¹²¹ in the structure and function of RNase A. Specifically, we have determined how the pK_a of His¹¹⁹ changes when Asp¹²¹ is replaced with an asparagine or alanine residue. We have extended this analysis to the pK_a of His¹² and the two other histidine residues in RNase A. Finally, we have assessed the effect of two different concentrations of the hydrolysis product 3'-UMP on the histidine pK_a values in all three enzymes.

Received for publication 17 July 1998 and in final form 17 November 1998.

Address reprint requests to Dr. Ronald T. Raines, Department of Biochemistry, University of Wisconsin–Madison, 433 Babcock Drive, Madison, WI 53706-1544. Tel.: 608-262-8588; Fax: 608-262-3453; E-mail: raines@biochem.wisc.edu.

Abbreviations used: RNase A, bovine pancreatic ribonuclease A; A, absorbance; DSS, 2,2-dimethyl-2-silapentane-5-sulfonate; 3'-UMP, uridine 3'-phosphate (otherwise Up).

© 1999 by the Biophysical Society

0006-3495/99/03/1571/09 \$2.00




FIGURE 1 Crystalline structure of the active site of RNase A bound to uridine 2',3'-cyclic vanadate (U > v), which is an analog of the transition state during catalysis. The structure was refined at 2.0 Å from x-ray and neutron diffraction data collected from crystals grown at pH 5.3 (Wlodawer et al., 1983). The structure has also been refined at 1.3 Å from x-ray diffraction data alone (Wladkowski et al., 1998). The His ··· Asp catalytic dyad is highlighted. The side chain of Phe¹²⁰ and the uracil base are not shown.

MATERIALS AND METHODS

Materials

RNase A (Type XII-A, lyophilized, salt free) was from Sigma Chemical (St. Louis, MO). 3'-UMP (98% pure; Sigma Chemical) was used without further purification. D121N RNase A and D121A RNase A were produced in *Escherichia coli* and purified as described by Schultz et al. (1998). D₂O (99.9%), DCl solution (35% v/v in D₂O), and NaOD solution (40% w/v in D₂O) were from Isotec (Miamisburg, OH). DSS was from Cambridge Isotope Laboratories (Andover, MA).

Procedures

To facilitate the pH titrations by ¹H-NMR spectroscopy, exchangeable enzymic hydrogens were replaced with deuteriums. To exchange hydrogen for deuterium, lyophilized enzyme was dissolved in D₂O, lyophilized again, and redissolved in D₂O. DCl solution was used to adjust the pH* to 3.0, where pH* represents a direct reading of pH that is not corrected for a deuterium isotope effect. The resulting solution was heated to 60°C for 1 h to exchange the amide protons (Markley, 1975b). NaCl and DSS were then added to final concentrations of 0.20 M and 0.50 mM, respectively. 3'-UMP was added from a stock solution of 0.27 M.

¹H-NMR spectra were recorded on a Bruker DMX 400 MHz spectrometer at 25°C using 16K data points and an acquisition time of 1.5 s with 128, 256, or 512 scans. Values of chemical shift (δ_{obs}) were recorded with

respect to that of DSS. pH* was adjusted to values between 3.0 and 9.0 by adding commercial solutions of DCl or NaOD that had been diluted 10-fold with D₂O.

Both the deuterium exchange procedure and the pH titrations were performed in solutions with an enzyme concentration of 50 mg/mL. Enzyme concentrations were determined by assuming that $A = 0.72$ at 277.5 nm for a 1.0 mg/mL solution (Sela et al., 1957). 3'-UMP quantities were determined by weight.

Values of δ_{obs} for the imidazolyl C-2 proton (otherwise H^{e1}; Fig. 1) at different values of [H⁺] were fitted to Eqs. 1, 2, or 3 (Fisher et al., 1998b):

$$\delta_{\text{obs}} = \delta_{\text{A}} \delta_{\text{AH}^+} \left(\frac{1 + K_{105}[\text{H}^+]}{\delta_{\text{A}} + \delta_{\text{AH}^+}(K_{105}[\text{H}^+])} \right) \quad (1)$$

$$\delta_{\text{obs}} = \delta_{\text{A}} \delta_{\text{AH}^+} \left(\frac{1 + \frac{K_{12a}K_{119a} + K_{119a}[\text{H}^+]}{[\text{H}^+](K_{12a} + [\text{H}^+])}}{\delta_{\text{A}} + \delta_{\text{AH}^+} \frac{K_{12a}K_{119a} + K_{119a}[\text{H}^+]}{[\text{H}^+](K_{12a} + [\text{H}^+])}} \right) \quad (2)$$

$$\delta_{\text{obs}} = \delta_{\text{A}} \delta_{\text{AH}^+} \left(\frac{1 + \frac{K_{12a}K_{119a} + K_{12a}[\text{H}^+]}{[\text{H}^+](K_{119a} + [\text{H}^+])}}{\delta_{\text{A}} + \delta_{\text{AH}^+} \frac{K_{12a}K_{119a} + K_{12a}[\text{H}^+]}{[\text{H}^+](K_{119a} + [\text{H}^+])}} \right) - \delta_{\text{o}} \left(\frac{\delta_{\text{o}} - \delta_{\text{oH}^+}}{\delta_{\text{o}} + \delta_{\text{oH}^+}(K_{\text{o}}[\text{H}^+])} \right) \quad (3)$$

Data for His¹⁰⁵ were fitted to Eq. 1, which describes the pH titration of a group with one pK_a. The apparent interaction between the two active-site histidine residues—His¹² and His¹¹⁹—necessitates a fit involving the microscopic pK_a values of these two residues. Data for His¹¹⁹, which has two microscopic pK_a values, were fitted to Eq. 2. Data for His¹², which has an acidic inflection in the pH* titration in addition to two microscopic pK_a values (Karpeisky and Yakovlev, 1981), were fitted to Eq. 3. The difference in the two microscopic pK_a values for each residue must be identical (Wyman and Gill, 1990). To accommodate this constraint, data for His¹² and His¹¹⁹ were fitted simultaneously to Eqs. 2 and 3. All data were fitted by using the program MATHEMATICA 3.0 from Wolfram Research (Champaign, IL).

In Eqs. 2 and 3, an "a" subscript refers to the microscopic pK_a of one active-site histidine residue when the other active-site histidine residue is protonated, and a "b" subscript refers to the pK_a of this histidine when the other histidine residue is unprotonated. In Eq. 3, the "o" subscript refers to the pK_a for the acidic inflection observed in the His¹² titration curve. Equations 1–3 are analogous to those used by Schechter and co-workers (Shrager et al., 1972) and by us (Chivers et al., 1997; Fisher et al., 1998b).

RESULTS

Titration of the C(2)-H chemical shifts of unliganded enzymes

RNase A has four histidine residues. The pK_a values for His¹², His¹¹⁹, and His¹⁰⁵ of RNase A can be determined by analyzing the chemical shift of the imidazolyl C-2 proton (Fig. 1) upon changing pH* (Markley, 1975b). The other histidine residue in RNase A, His⁴⁸, is inaccessible to solvent and its titration curve shows anomalous behavior with pH*, preventing a determination of its pK_a value (Markley, 1975a). Results from the pH titration of the unliganded ribonucleases are shown in Fig. 2, and the calculated pK_a values and limiting chemical shifts are listed in Table 1. The




FIGURE 2 pH dependence of the histidine $^1\text{H}^{\text{N}}$ signals of wild-type ribonuclease A and the D121N and D121A variants in D_2O . (A) Wild-type ribonuclease A (B) D121N ribonuclease A. (C) D121A ribonuclease A. Chemical shifts are shown for all four histidine residues: His 12 (●), His 119 (○), His 105 (■), and His 48 (×). Titrations were carried out at 25°C in buffer containing enzyme (3.5 mM) and NaCl (0.20 M). The pK_a values determined from fitting the data to Eqs. 1–3 are listed in Table 1.

pK_a values of His 12 , His 119 , and His 105 in wild-type RNase A and the D121N and D121A variants were well defined; the largest contribution to error appears to lie in the measurement of pH itself. Such systematic errors largely disappear when differences in pK_a values are calculated for any given titration.

Lys 41 , like His 12 and His 119 , resides in the active site of RNase A. The pK_a of the side chain of Lys 41 is near 9.0 (Jentoft et al., 1981), a value much greater than that of any histidine residue in RNase A. Accordingly, a perturbation to the pK_a of Lys 41 upon replacing Asp 121 is unlikely to have a substantial effect on the titrations and interpretations reported herein.

The microscopic pK_a values indicate that protonation of the active-site histidine residues has negative cooperativity (that is, protonation of one histidine residue disfavors protonation of the other) as had been observed previously (Markley and Finkenstadt, 1975). The negative cooperativity observed here is slightly greater than that in the previous study, perhaps because the previous observations were made in a solution of higher salt concentration (0.30 M versus 0.20 M NaCl), which should reduce Coulombic interactions.

Replacing Asp 121 with an asparagine or alanine residue decreases the pK_a of His 119 by 0.11 units (which is the average of the microscopic values), and increases the pK_a of His 12 by ~ 0.15 units. For all three enzymes, the titration of His 12 displayed an inflection in the acidic region (Karpeisky and Yakovlev, 1981). The proximity of this inflection to the pK_a of His 12 and the lack of data below $\text{pH}^* 3$ made for a large error in its determination. For all three enzymes, the titration of His 48 also displayed an inflection in the acidic region, as had been seen previously (Markley, 1975b). The effect of the substitutions on the titration of His 48 was minor. No effect was observed on the titration of His 105 . The limiting chemical shifts of each histidine residue were almost identical for all three enzymes.

Titration of the C(2)-H chemical shifts of enzyme · product complexes

pH titrations of wild-type RNase A and the D121N and D121A variants were performed in the presence of the reaction product 3'-UMP. In these titrations, the concentration of enzyme was 3.5 mM and that of 3'-UMP was either 1.5 mM or 11 mM. The results are shown in Figs. 3 and 4, and the calculated pK_a values and limiting chemical shifts are listed in Table 1. The data were fitted without regard to the association of 3'-UMP.

In the wild-type enzyme, His 119 is more perturbed by the presence of 3'-UMP than is His 12 , as had been observed previously (Eftink and Biltonen, 1983). In the presence of 11 mM 3'-UMP, replacing Asp 121 with an asparagine residue reduces the pK_a of His 119 by 0.36 and increases that of His 12 by 0.21. Replacing Asp 121 with an alanine reduces the pK_a of His 119 by 0.97 and increases that of His 12 by 0.35.

TABLE 1 Microscopic pK_a values and limiting C(2)-H chemical shifts of histidine residues in wild-type ribonuclease A and the D121N and D121A variants in the absence and presence of uridine 3'-phosphate

Residue	[3'-UMP] (mM)	Wild-Type Ribonuclease A			D121N Ribonuclease A			D121A Ribonuclease A		
		pK_a^*	δ_{low}	$\delta_{high}^{\#}$	pK_a	δ_{low}	δ_{high}	pK_a	δ_{low}	δ_{high}
His ¹²	0	5.74 ± 0.03	8.89 ± 0.03	7.67 ± 0.01	5.84 ± 0.02	8.96 ± 0.02	7.67 ± 0.01	5.88 ± 0.02	8.98 ± 0.01	7.67 ± 0.01
		6.12 ± 0.04			6.28 ± 0.06			6.32 ± 0.04		
His ¹¹⁹	0	3.99 ± 0.28	9.01 ± 0.03	8.86	3.19 ± 1.35	8.96 ± 0.11	8.86	2.92 ± 1.87	8.97 ± 0.31	8.86
		5.94 ± 0.02	8.82 ± 0.01	7.76 ± 0.01	5.80 ± 0.03	8.82 ± 0.01	7.74 ± 0.01	5.80 ± 0.02	8.84 ± 0.01	7.74 ± 0.00
His ¹⁰⁵	0	6.78 ± 0.01	8.75 ± 0.00	7.70 ± 0.01	6.78 ± 0.01	8.76 ± 0.00	7.70 ± 0.00	6.73 ± 0.01	8.76 ± 0.00	7.71 ± 0.01
		6.33 ± 0.02	8.75 ± 0.01	7.66 ± 0.01	6.27 ± 0.02	8.78 ± 0.02	7.68 ± 0.05	6.39 ± 0.01	8.86 ± 0.01	7.68 ± 0.00
His ¹¹⁹	1.5	6.62 ± 0.07			6.49 ± 0.06			6.47 ± 0.06		
		4.14 ± 0.26	8.98 ± 0.02	8.86	4.26 ± 0.15	9.05 ± 0.02	8.86	3.78 ± 0.15	9.02 ± 0.02	8.86
His ¹¹⁹	1.5	6.74 ± 0.06	8.75 ± 0.02	7.78 ± 0.01	6.73 ± 0.06	8.67 ± 0.02	7.74 ± 0.01	6.37 ± 0.04	8.73 ± 0.01	7.75 ± 0.01
		7.03 ± 0.02	8.91 ± 0.02	8.86	6.95 ± 0.02	8.97 ± 0.02	8.86	6.44 ± 0.03	9.00 ± 0.02	8.86
His ¹⁰⁵	1.5	4.34 ± 0.59			4.63 ± 0.18			3.89 ± 0.26		
		6.86 ± 0.01	8.76 ± 0.00	7.70 ± 0.01	6.80 ± 0.01	8.78 ± 0.00	7.71 ± 0.00	6.78 ± 0.01	8.77 ± 0.00	7.70 ± 0.01
His ¹²	11	6.45 ± 0.03	8.72 ± 0.02	7.69 ± 0.01	6.66 ± 0.01	8.75 ± 0.04	7.71 ± 0.00	6.80 ± 0.01	8.78 ± 0.00	7.71 ± 0.00
		6.35 ± 0.24			6.30 ± 0.13			6.50 ± 0.05		
His ¹¹⁹	11	3.77 ± 0.36	8.99 ± 0.02	8.86	2.63 ± 1.01	9.19 ± 1.53	8.86	2.78 ± 0.25	9.26 ± 0.15	8.86
		7.95 ± 0.29	8.69 ± 0.03	7.76 ± 0.03	7.86 ± 0.15	8.65 ± 0.01	7.74 ± 0.01	7.18 ± 0.05	8.64 ± 0.01	7.75 ± 0.00
His ¹⁰⁵	11	7.85 ± 0.07	8.96 ± 0.04	8.86	7.49 ± 0.02	9.04 ± 0.02	8.86	6.88 ± 0.02	9.09 ± 0.01	8.86
		4.06 ± 0.75			3.88 ± 0.13			3.89 ± 0.08		
His ¹⁰⁵	11	6.86 ± 0.01	8.76 ± 0.00	7.70 ± 0.00	6.87 ± 0.01	8.78 ± 0.00	7.71 ± 0.00	6.79 ± 0.00	8.77 ± 0.00	7.71 ± 0.00

Data were obtained at 25°C in buffers containing 3'-UMP (0, 1.5 mM, or 11 mM) and NaCl (0.20 M). Values were determined by fitting the experimental data in Figs. 2–4 to Eq. 1 for His¹⁰⁵, Eq. 2 for His¹¹⁹, and Eq. 3 for His¹². Errors were determined by a nonlinear least-squares fit of the data to Eqs. 1–3. *For His¹² and His¹¹⁹, the first pK_a value is that observed when the other active-site histidine residue is protonated (corresponding to K_{12a} and K_{119a} in Eqs. 2–4); the second, when it is unprotonated (K_{12b} and K_{119b}).

[#]The δ_{high} values for additional pK_a values are defined to be 8.86.

These comparisons are based upon the microscopic pK_a values of His¹¹⁹ (in which His¹² is not protonated) and the microscopic pK_a values of His¹² (in which His¹¹⁹ is protonated), as this combination is the predominant one. The presence of 3'-UMP produces a new pK_a of 3.9 during the titration of His¹¹⁹. The effect of 3'-UMP on the cooperativity of the titrations is discussed below.

In the course of titrating D121N RNase A in the presence of 3'-UMP, we observed that the chemical shift of the C-4 proton (otherwise H^{δ2}) of His¹¹⁹ had shifted downfield at acidic pH by ~0.3 ppm, to a spectral region in which it is no longer obscured by signals from phenylalanine and tyrosine residues. A comparison of spectra of D121N RNase A with that for the wild-type enzyme led us to conclude that our assignment of the C-4 proton of His¹¹⁹ and that of Kaptein and co-workers are correct (Lenstra et al., 1979).

DISCUSSION

Unliganded enzymes

We find that upon loss of Asp¹²¹, the pK_a of His¹¹⁹ drops by only 0.1 unit: a value much smaller than that expected for the loss of a hydrogen bond between functional groups of opposite charge (Fersht et al., 1985). Why is the pK_a of His¹¹⁹ not perturbed more by Asp¹²¹? The side chain of His¹¹⁹ is flexible, occupying two distinct positions in wild-type RNase A (Borkakoti et al., 1982). In position A, N^{ε2} of His¹¹⁹ forms a hydrogen bond with O^{δ1} of Asp¹²¹ (Fig. 1),

with the side chain of His¹¹⁹ having torsion angles of $\chi_1 = 149^\circ$ and $\chi_2 = -101^\circ$. In position B, the imidazolyl group is removed from Asp¹²¹ by 7 Å, with $\chi_1 = -69^\circ$ and $\chi_2 = -63^\circ$. In analyses of the crystalline and solution structure of RNase A, His¹¹⁹ has been found in position A, position B, or both [for reviews, see Gilliland (1997) and González et al. (1997)]. The pH-dependencies of the thermodynamic stabilities of wild-type RNase A and the D121N, D121A, and H119A variants suggest that His¹¹⁹ of the wild-type enzyme resides predominantly in position A when its side chain is protonated (Quirk, 1996; Quirk et al., 1998). These studies also revealed that the pK_a of Asp¹²¹ is perturbed by one unit upon the folding of RNase A. One possibility, then, is that the deprotonated state of His¹¹⁹ favors position B, which is removed from Asp¹²¹. A cation (e.g., Na⁺) could then replace the neutral His¹¹⁹ side chain in position A. A second possibility is that another residue buffers the effect of removing Asp¹²¹. This residue could be Glu¹¹¹, which is only 5 Å from the imidazolyl group of His¹¹⁹ when it resides in position B.

Replacing Asp¹²¹ with an asparagine or alanine residue increases the pK_a of His¹² by 0.15, an effect larger and in the opposite direction to that on His¹¹⁹. A simple Coulombic calculation indicates that removing the anionic side chain of Asp¹²¹ is expected to *decrease* the pK_a of His¹² by ~0.2 (= $0.4 \times 7^2/10^2$ by Coulomb's law). This calculation is based on the observations that protonating His¹² decreases the pK_a of His¹¹⁹ by 0.4 units (and vice versa) and that His¹² is




FIGURE 3 pH dependence of the histidine $^1H^{\epsilon 1}$ signals of wild-type ribonuclease A and the D121N and D121A variants in D_2O containing a low concentration of 3'-UMP. (A) Wild-type ribonuclease A. (B) D121N ribonuclease A. (C) D121A ribonuclease A. Chemical shifts are shown for all four histidine residues: His¹² (●), His¹¹⁹ (○), His¹⁰⁵ (■), and His⁴⁸ (×). Titrations were carried out at 25°C in buffer containing enzyme (3.5 mM), 3'-UMP (1.5 mM), and NaCl (0.20 M). The pK_a values determined from fitting the data to Eqs. 1–3 are listed in Table 1.




FIGURE 4 pH dependence of the histidine $^1H^{\epsilon 1}$ signals of wild-type ribonuclease A and the D121N and D121A variants in D_2O containing a high concentration of 3'-UMP. (A) Wild-type ribonuclease A. (B) D121N ribonuclease A. (C) D121A ribonuclease A. Chemical shifts are shown for all four histidine residues: His¹² (●), His¹¹⁹ (○), His¹⁰⁵ (■), and His⁴⁸ (×). Titrations were carried out at 25°C in buffer containing enzyme (3.5 mM), 3'-UMP (11 mM), and NaCl (0.20 M). The pK_a values determined from fitting the data to Eqs. 1–3 are listed in Table 1.

separated from His¹¹⁹ and Asp¹²¹ by 7 and 10 Å, respectively, and on the assumption that the dielectric constant is similar for both interactions. A role for His¹¹⁹—through its conformational flexibility—in this curious perturbation of His¹² is difficult to justify because the His¹²–His¹¹⁹ distance is similar in the A and B positions, and the change in the microscopic pK_a values is similar when His¹¹⁹ is in the A or B position. The replacement of Asp¹²¹ therefore appears to have effects on a more global scale that quite possibly involves some of the many basic residues in RNase A.

Our results conflict with those from a study that used a semisynthetic RNase A (Cederholm et al., 1991). That enzyme, RNase-(1–118) · (111–124), consists of a noncovalent complex between residues 1–118 of RNase A and a synthetic 14-residue peptide consisting of residues 111–124 of RNase A. Surprisingly, replacing Asp¹²¹ of the semisynthetic enzyme with an asparagine residue increases the macroscopic pK_a of His¹¹⁹ by 0.05 and decreases the macroscopic pK_a of His¹² by 0.09; effects that are in opposition to ours [Cederholm et al. (1991) did not report any microscopic pK_a values]. Most notably, the difference between the pK_a values of His¹¹⁹ and His¹² ($\Delta pK_a = pK_a^{\text{His}119} - pK_a^{\text{His}12}$) increases from 0.3 for the parent semisynthetic enzyme to 0.45 for the D121N variant, but ΔpK_a decreases from 0.18 for authentic wild-type RNase A to –0.04 for D121N RNase A. This deviation is consistent with the semisynthesis having introduced unpredictable structural perturbations. Indeed, x-ray diffractions analyses of the two semisynthetic enzymes revealed many small changes in their structures (de Mel et al., 1992). In contrast, the structure of D121N RNase A is essentially identical to that of the authentic wild-type enzyme (Schultz et al., 1998).

Ribonuclease A · 3'-UMP complexes

Both NMR spectroscopy and x-ray diffraction analysis have been used to determine the three-dimensional structures of complexes of RNase A with various mononucleotides, including 3'-UMP (Bruix et al., 1991). These structures have been the subject of recent reviews (Gilliland, 1997; González et al., 1997). No significant perturbation to enzymic structure is observed upon formation of the complexes, which have active sites analogous to that shown in Fig. 1.

Our pH titrations reveal new insight on the interaction between RNase A and 3'-UMP. The relationship between the microscopic pK_a values of the active-site histidine residues can be defined by the parameter *c*, with *c* > 1 indicating positive cooperativity and *c* < 1 indicating negative cooperativity (Wyman and Gill, 1990). For titrations in the active site of RNase A, $c = K_{12b}/K_{12a} = K_{119b}/K_{119a}$ (Fisher et al., 1998b). The values for *c* for wild-type RNase A and the D121N and D121A variants in the absence and presence of 3'-UMP are listed in Table 2. In the absence of 3'-UMP, *c* < 1 for all three enzymes. The presence of 3'-UMP increases the value of *c*. Moreover, in the presence of a high concentration of 3'-UMP, *c* > 1 for all three

TABLE 2 Cooperativity (*c*) values from the analysis of the microscopic pK_a values of the active-site histidine residues of wild-type ribonuclease A and the D121N and D121A variants in the absence and presence of uridine 3'-phosphate

[3'-UMP] (mM)	Ribonuclease A		
	Wild-Type	D121N	D121A
0	0.42	0.36	0.36
1.5	0.51	0.60	0.84
11	1.3	2.3	2.0

Values were calculated from the data in Table 1 and the equation: $c = [(K_{12b}/K_{12a}) + (K_{119b}/K_{119a})]/2$.

enzymes. In other words, protonation of one active-site histidine residue then *favors* protonation of the other. Such positive cooperativity is unusual for the titration of two proximal acids of like charge. This result is consistent with the anionic phosphoryl group of 3'-UMP binding more strongly to the active site when both histidine residues are protonated and thus cationic.

To extract as much information as possible from the ¹H-NMR data, we attempted to fit the microscopic pK_a values in Table 1 to Eq. 4, which describes the pH-dependence of the dissociation constant (*K_d*) of the enzyme · 3'-UMP complex:

$$\begin{aligned}
 K_d = & \frac{K_d^{\text{int}} \left(\frac{K_{12b}K_{119a}}{K_{12b}^1 K_{119a}^1} \right)}{1 + \frac{[H^+]}{K_{119a}^1} + \frac{[H^+]K_{119b}^1 + [H^+]^2}{K_{12b}^1 K_{119a}^1} + \frac{[H^+]^3}{K_{12b}^1 K_{119a}^1 K_p^1}} \\
 & + \frac{K_d^{\text{int}} \left(\frac{K_{12b}}{K_{12b}^1} \right)}{1 + \frac{K_{119a}^1}{[H^+]} + \frac{K_{119b}^1 + [H^+]}{K_{12b}^1} + \frac{[H^+]^2}{K_{12b}^1 K_p^1}} \\
 & + \frac{K_d^{\text{int}} \left(\frac{K_{119b}}{K_{119b}^1} \right)}{1 + \frac{K_{12b}^1 + [H^+]}{K_{119b}^1} + \frac{[H^+]^2}{K_{119b}^1 K_p^1} + \frac{K_{12b}^1 K_{119a}^1}{K_{119b}^1 [H^+]}} \\
 & + \frac{K_d^{\text{int}}}{1 + \frac{[H^+]}{K_p^1} + \frac{K_{12b}^1 K_{119a}^1 + K_{12b}^1 K_{119b}^1}{[H^+]^2}} \\
 & + \frac{K_d^{\text{int}} \left(\frac{K_p^1}{K_p^1} \right)}{1 + \frac{K_p^1}{[H^+]} + \frac{K_{119b}^1 K_p^1}{[H^+]^2} + \frac{K_{12b}^1 K_{119a}^1 K_p^1}{[H^+]^3}} \quad (4)
 \end{aligned}$$

where *K_{12a}* refers to His¹² when His¹¹⁹ is protonated, *K_{119a}* refers to His¹¹⁹ when His¹² is protonated, *K_{12b}* refers to His¹² when His¹¹⁹ is not protonated; *K_{119b}* refers to His¹¹⁹ when His¹² is not protonated, and *K_p* refers to the phosphoryl group of 3'-UMP as obtained from steady-state kinetic analyses (Schultz et al., 1998). The *K_d*^{int} is the intrinsic

dissociation constant of the enzyme · 3'-UMP complex when the enzymic active site is fully protonated and the inhibitor is fully deprotonated (Fig. 5). Terms containing an "I" superscript refer to pK_a values in the presence of 3'-UMP. The equilibrium constants in Eq. 4 are related by the thermodynamic cube shown in Fig. 5. The concentration of enzyme · 3'-UMP complex ([E · I]) at any pH can be determined from the values of K_d and the equation:

$$K_d = \frac{([E]_{\text{total}} - [E \cdot I])([I]_{\text{total}} + [E \cdot I])}{[E \cdot I]} \quad (5)$$

where [E]_{total} is the total concentration of RNase A and [I]_{total} is the total concentration of 3'-UMP. The calculated chemical shift (δ_{calc}) is then simply the weighted sum of the chemical shifts for the unliganded (δ_{free}) and bound (δ_{bound}) enzyme, as in the equation:

$$\delta_{\text{calc}} = \delta_{\text{free}} \left(1 - \frac{[E \cdot I]}{[E]_{\text{total}}} \right) + \delta_{\text{bound}} \frac{[E \cdot I]}{[E]_{\text{total}}} \quad (6)$$

Favorable Coulombic interactions contribute to the affinity of RNase A for 3'-UMP. Isothermal titration calorimetry indicates that the RNase A · 3'-UMP complex has K_d = 0.054 mM at 25°C in 0.10 M MES-NaOH buffer (pH 6.0) containing NaCl (0.10 M) (Fisher et al., 1998b). This solution has a [Na⁺] of 0.142 M. Because the pH titrations described herein were performed in solutions with a [Na⁺] of 0.20 M, the RNase A · 3'-UMP complex is likely to have K_d > 0.054 M. To attempt to fit the microscopic pK_a values to Eqs. 4–6, we assumed that only the active site is occupied when 3'-UMP is present at a low concentration, such as 1.5 mM.

Our attempts to fit the experimental data to Eqs. 4–6 revealed a problem. Chemical shifts observed in the presence of 1.5 mM 3'-UMP should fall between that of the unliganded enzyme and that in the presence of 11 mM 3'-UMP. Yet, the chemical shifts calculated with Eq. 6 do




FIGURE 5 Thermodynamic cube showing the relationships between the titration of the imidazolyl groups of His¹² and His¹¹⁹ of ribonuclease A (E), the phosphoryl group of 3'-UMP (I), and a residue that could be Asp⁸³, in both the unliganded enzyme and the enzyme · 3'-UMP complex. The nomenclature used is as in Eq. 4.

not obey this constraint. The perturbation to these chemical shifts by a low concentration of 3'-UMP was equal to or greater than that from a high concentration. We conclude that the RNase A plus 3'-UMP system is at least a three-body one when 3'-UMP is present at high concentration. A second molecule of RNase A would have resulted in significant spectral line broadening, which was not observed. Instead, we conclude that a second molecule of 3'-UMP binds to the enzyme, affecting the binding of the first. Our conclusion is in accord with kinetic data suggesting that more than one molecule of cytidine 2',3'-cyclic phosphate (C > p) binds to RNase A when [C > p] > 10 mM (Moussaoui et al., 1998).

Evidence for subsites in RNase A comes from many workers [for reviews, see Parés et al. (1991); Nogués et al. (1995); Raines (1998)]. That RNase A binds and cleaves polymeric substrates is itself suggestive of enzymic subsites that bind to monomeric units. A number of x-ray diffraction analyses on complexes with substrate analogs have revealed the presence of subsites (Eftink and Biltonen, 1983; McPherson et al., 1986; Fontecilla-Camps et al., 1994). In addition, Irie and co-workers used ³¹P-NMR spectroscopy to provide evidence of binding sites for three phosphoryl groups and two bases (Irie et al., 1984). Recently, a fourth phosphoryl-group subsite has been discovered (Fisher et al., 1998a), and the existence of a fifth nucleotide subsite has been inferred (Moussaoui et al., 1998). Perhaps most germane to this work is the finding that adenine, adenosine, adenosine 3'-phosphate, and adenosine 5'-phosphate increase the rate of the RNase A-catalyzed hydrolysis of C > p (Wieker and Witzel, 1967; Haffner and Wang, 1973; Moussaoui et al., 1998).

Acidic inflections

The inflection with a pK_a of 3.9 observed in the titration of His¹¹⁹ in the presence of 3'-UMP almost certainly arises from the deprotonation of the phosphoryl group. The decrease from a pK_a of 5.8 for free 3'-UMP is consistent with its interaction with the two active-site histidine residues (Tanokura, 1983). A thermodynamic determination of K_p (Flogel and Biltonen, 1975) as a function of pH supports this view, as do studies that relied on ³¹P-NMR spectroscopy (Gorenstein and Wyrwicz, 1973).

The inflection seen in the titration of His¹² has been reported widely, but its assignment less so. Perhaps the most thoughtful analysis of the titration behavior of His¹² comes from Karpeisky and Yakovlev (1981). These workers concluded from much evidence that the functional group responsible for the inflection is the side chain of Asp⁸³, which mediates its effect through Thr⁴⁵ and its interaction with both the uracil base and His¹². The importance of hydrogen bonding between Thr⁴⁵ and Asp⁸³ to the binding of a uracil base has been demonstrated by site-directed mutagenesis (deICardayré and Raines, 1995).

Low-barrier hydrogen bonds?

The small contribution of Asp¹²¹ to catalysis belies the importance of the hydrogen bond between His¹¹⁹ and Asp¹²¹ (Schultz et al., 1998). Nonetheless, a similar His ··· Asp dyad has been proposed to form a low-barrier hydrogen bond of extraordinary strength during catalysis by the protease chymotrypsin (Frey et al., 1994; Cassidy et al., 1997), though this interpretation is controversial (Ash et al., 1997). One criterion for such a bond is an ¹H chemical shift of 17–20 ppm (Cleland and Kreevoy, 1994). The ¹H chemical shift of N_{ε2}H of His¹¹⁹ appears at a much higher field, <12 ppm (J. L. Markley, personal communication). By this criterion, the His ··· Asp dyad of RNase A does not have a low-barrier hydrogen bond. In theory, the His ··· Asn dyad of D121N RNase A could also form a low-barrier hydrogen bond. Another criterion for such a bond is that it arise from acids with matched pK_a values (Cleland, 1992). The pK_a values of imidazole [14.2 (Walba and Isensee, 1955)] and acetamide [15.1 (Bordwell, 1988)], which are reasonable models for the side chains of His¹¹⁹ and Asn¹²¹, respectively, are indeed similar. The existence of a low-barrier hydrogen bond would, however, require that the side chain of His¹¹⁹ be in its neutral imidazole form. ¹H-NMR titrations of His¹¹⁹ in the D121N variant (Figs. 2–4) and pH-rate profiles for catalysis by D121N RNase A (Schultz et al., 1998) indicate that His¹¹⁹ titrates between its imidazole and imidazolium forms with a pK_a value similar to that of His¹¹⁹ in wild-type RNase A. These data deny the existence of a low-barrier hydrogen bond in the His ··· Asn dyad.

CONCLUSIONS

Though conserved during evolution, Asp¹²¹ of RNase A has little effect on the ionization of His¹¹⁹. ¹H-NMR spectroscopy has shown that neither the microscopic pK_a values of His¹¹⁹ nor those of any other histidine residue is perturbed by more than 0.2 units when Asp¹²¹ is replaced by an asparagine or alanine residue. Microscopic pK_a values of His¹² and His¹¹⁹ in the presence of 3'-UMP exhibit positive rather than negative cooperativity. Finally, ¹H-NMR data provide evidence that RNase A can bind a second molecule of the 3'-UMP product within its active site. As in previous studies (Quirk et al., 1998; Schultz et al., 1998), we conclude that the mere presence of a His ··· Asp catalytic dyad in an active site is not a mandate for its playing a crucial role in the structure or function of the enzyme.

This work was supported by National Institutes of Health Grant GM44783. D.J.Q. was supported by Cellular and Molecular Biology Training Grant GM07215 from the National Institutes of Health. NMR spectroscopy was carried out at the National Magnetic Resonance Facility at Madison, which is funded by National Institutes of Health Grant RR02301; equipment in the facility was purchased with funds from the University of Wisconsin, the NSF Biological Instrumentation Program (Grant DMB-8415048), the National Biomedical Research Technology Program (Grant RR02301), the National Institutes of Health Shared Instrumentation Program (Grant RR02781), and the U.S. Department of Agriculture.

REFERENCES

- Antosiewicz, J., J. A. McCammon, and M. K. Gilson. 1996. The determinants of pK_as in proteins. *Biochemistry*. 35:7819–7833.
- Ash, E. L., J. L. Sudmeier, E. C. De Fabo, and W. W. Bachovchin. 1997. A low-barrier hydrogen bond in the catalytic triad of serine proteases? Theory versus experiment. *Science*. 278:1128–1132.
- Beintema, J. J. 1987. Structure, properties and molecular evolution of pancreatic-type ribonucleases. *Life Chem. Rep.* 4:333–389.
- Beintema, J. J., C. Schüller, M. Irie, and A. Carsana. 1988. Molecular evolution of the ribonuclease superfamily. *Prog. Biophys. Mol. Biol.* 51:165–192.
- Bordwell, F. G. 1988. Equilibrium acidities in dimethyl sulfoxide solution. *Acc. Chem. Res.* 21:456–463.
- Borkakoti, N., D. S. Moss, and R. A. Palmer. 1982. Ribonuclease-A: least-squares refinement of the structure at 1.45 Å resolution. *Acta Crystallogr. B.* 38:2210–2217.
- Bruix, M., M. Rico, C. González, J. L. Neira, J. Santoro, J. L. Nieto, and H. Rüterjans. 1991. Two dimensional ¹H-NMR studies of the solution structure of RNase A–pyrimidine-nucleotide complexes. In *Structure, Mechanism and Function of Ribonucleases*. R. de Llorens, C. M. Cuchillo, M. V. Nogués, and X. Parés, editors. Universitat Autònoma de Barcelona, Bellaterra, Spain. 15–20.
- Cassidy, C. S., J. Lin, and P. A. Frey. 1997. A new concept for the mechanism of action of chymotrypsin: the role of the low-barrier hydrogen bond. *Biochemistry*. 36:4576–4584.
- Cederholm, M. T., J. A. Stuckey, M. S. Doscher, and L. Lee. 1991. Histidine pK_a shifts accompanying the inactivating Asp¹²¹ → Asn substitution in a semisynthetic bovine pancreatic ribonuclease. *Proc. Natl. Acad. Sci. U.S.A.* 88:8116–8120.
- Chivers, P. T., K. E. Prehoda, B. Volkman, B.-M. Kim, J. L. Markley, and R. T. Raines. 1997. Microscopic pK_a values of *Escherichia coli* thioredoxin. *Biochemistry*. 36:14985–14991.
- Cleland, W. W. 1992. Low-barrier hydrogen bonds and low fractionation factor bases in enzymatic reactions. *Biochemistry*. 31:317–319.
- Cleland, W. W., and M. M. Kreevoy. 1994. Low-barrier hydrogen bonds and enzymic catalysis. *Science*. 264:1887–1890.
- Cuchillo, C. M., X. Parés, A. Guasch, T. Barman, F. Travers, and M. V. Nogués. 1993. The role of 2',3'-cyclic phosphodiester in the bovine pancreatic ribonuclease A catalysed cleavage of RNA: intermediates or products? *FEBS Lett.* 333:207–210.
- delCardayré, S. B., and R. T. Raines. 1995. A residue to residue hydrogen bond mediates the nucleotide specificity of ribonuclease A. *J. Mol. Biol.* 252:328–336.
- de Mel, V. S. J., P. D. Martin, M. S. Doscher, and B. F. P. Edwards. 1992. Structural changes that accompany the reduced catalytic efficiency of two semisynthetic ribonuclease analogs. *J. Biol. Chem.* 267:247–256.
- Eftink, M. R., and R. L. Biltonen. 1983. Energetics of ribonuclease A catalysis. 1. pH, ionic strength, and solvent isotope dependence of the hydrolysis of cytidine cyclic 2',3'-phosphate. *Biochemistry*. 22: 5123–5134.
- Fersht, A. R., J.-P. Shi, J. Knill-Jones, D. M. Lowe, A. J. Wilkinson, D. M. Blow, P. Brick, P. Arter, M. M. Y. Waye, and G. Winter. 1985. Hydrogen bonding and biological specificity analysed by protein engineering. *Nature*. 314:235–238.
- Findlay, D., D. G. Herries, A. P. Mathias, B. R. Rabin, and C. A. Ross. 1961. The active site and mechanism of action of bovine pancreatic ribonuclease. *Nature*. 190:781–784.
- Fisher, B. M., J. E. Grilley, and R. T. Raines. 1998a. A new remote subsite in ribonuclease A. *J. Biol. Chem.* 273:34134–34138.
- Fisher, B. M., L. W. Schultz, and R. T. Raines. 1998b. Coulombic effects of remote subsites on the active site of ribonuclease A. *Biochemistry*. 37:17386–17401.
- Flogel, M., and R. L. Biltonen. 1975. The pH dependence of the thermodynamics of the interaction of 3'-cytidine monophosphate with ribonuclease A. *Biochemistry*. 14:2610–2615.
- Fontecilla-Camps, J. C., R. de Llorens, M. H. le Du, and C. M. Cuchillo. 1994. Crystal structure of ribonuclease A · d(ApTpApApG) complex. *J. Biol. Chem.* 269:21526–21531.

- Frey, P. A., S. A. Whitt, and J. B. Tobin. 1994. A low-barrier hydrogen bond in the catalytic triad of serine proteases. *Science*. 264:1927–1930.
- Gilliland, G. L. 1997. Crystallographic studies of ribonuclease complexes. In *Ribonucleases: Structures and Functions*. G. D'Alessio and J. F. Riordan, editors. Academic Press, New York. 306–341.
- González, C., J. Santoro, and M. Rico. 1997. NMR solution structures of ribonuclease A and its complexes with mono and dinucleotides. In *Ribonucleases: Structures and Functions*. G. D'Alessio and J. F. Riordan, editors. Academic Press, New York. 343–381.
- Gorenstein, D. G., and A. Wyrwicz. 1973. ³¹P-NMR study on the binding of 3'-cytidine monophosphate to ribonuclease A. I. *Biochem. Biophys. Res. Commun.* 54:976–982.
- Haffner, P. H., and J. H. Wang. 1973. Chemical kinetic and proton magnetic resonance studies of 5'-adenosine monophosphate binding to ribonuclease A. *Biochemistry*. 12:1608–1618.
- Irie, M., H. Watanabe, K. Ohgi, M. Tobe, G. Matsumura, Y. Arata, T. Hirose, and S. Inayama. 1984. Some evidence suggesting the existence of P2 and B3 sites in the active site of bovine pancreatic ribonuclease A. *J. Biochem.* 95:751–759.
- Jentoft, J. E., T. A. Gerken, N. Jentoft, and D. G. Dearborn. 1981. [¹³C]Methylated ribonuclease A. ¹³C-NMR studies of the interaction of lysine 41 with active site ligands. *J. Biol. Chem.* 256:231–236.
- Karpeisky, M. Y., and G. I. Yakovlev. 1981. Topochemical principles of the substrate specificity of nucleases. *Sov. Sci. Rev., Sect. D*. 2:145–257.
- Kartha, G., J. Bello, and D. Harker. 1967. Tertiary structure of ribonuclease. *Nature*. 213:862–865.
- Lenstra, J. A., B. G. J. M. Bolscher, S. Stob, J. J. Beintema, and R. Kaptein. 1979. The aromatic residues of bovine pancreatic ribonuclease studied by proton nuclear magnetic resonance. *Eur. J. Biochem.* 98:385–397.
- Markley, J. L. 1975a. Correlation proton magnetic resonance studies at 250 MHz of bovine pancreatic ribonuclease. II. The pH and inhibitor-induced conformational transitions affecting histidine-48 and one tyrosine residue of ribonuclease A. *Biochemistry*. 14:3554–3561.
- Markley, J. L. 1975b. Correlation proton magnetic resonance studies at 250 MHz of bovine pancreatic ribonuclease. I. Reinvestigation of the histidine peak assignment. *Biochemistry*. 14:3546–3553.
- Markley, J. L., and W. R. Finkstadt. 1975. Correlation proton magnetic resonance studies at 250 MHz of bovine pancreatic ribonuclease. III. Mutual electrostatic interaction between histidine residues 12 and 119. *Biochemistry*. 14:3562–3566.
- McPherson, A., G. Brayer, D. Cascio, and R. Williams. 1986. The mechanism of binding of a polynucleotide chain to pancreatic ribonuclease. *Science*. 232:765–768.
- Moussaoui, M., M. V. Nogués, A. Guasch, T. Barman, F. Travers, and C. M. Cuchillo. 1998. The subsite structure of bovine pancreatic ribonuclease A account for the abnormal kinetic behavior with cytidine 2',3'-cyclic phosphate. *J. Biol. Chem.* 273:25562–25572.
- Nogués, M. V., M. Vilanova, and C. M. Cuchillo. 1995. Bovine pancreatic ribonuclease A as a model of an enzyme with multiple substrate binding sites. *Biochim. Biophys. Acta*. 1253:16–24.
- Parés, X., M. V. Nogués, R. de Llorens, and C. M. Cuchillo. 1991. Structure and function of ribonuclease A binding subsites. *Essays Biochem.* 26:89–103.
- Quirk, D. J. 1996. Role of the His ··· Asp catalytic dyad of ribonuclease A. Ph.D. thesis, University of Wisconsin–Madison.
- Quirk, D. J., C. Park, J. E. Thompson, and R. T. Raines. 1998. His ··· Asp catalytic dyad of ribonuclease A: conformational stability of the wild-type, D121N, D121A, and H119A enzymes. *Biochemistry*. 37:17958–17964.
- Raines, R. T. 1998. Ribonuclease A. *Chem. Rev.* 98:1045–1065.
- Rico, M., M. Bruix, J. Santoro, C. González, J. L. Neira, J. L. Nieto, and J. Herranz. 1989. Sequential 1H-NMR assignment and solution structure of bovine pancreatic ribonuclease A. *Eur. J. Biochem.* 183:623–638.
- Robertson, A. D., E. O. Purisima, M. A. Eastman, and H. A. Scheraga. 1989. Proton NMR assignments and regular backbone structure of bovine pancreatic ribonuclease A in aqueous solution. *Biochemistry*. 1989:5930–5938.
- Saunders, M., A. Wishnia, and J. Kirkwood. 1957. The nuclear magnetic resonance spectrum of ribonuclease. *J. Am. Chem. Soc.* 79:3289.
- Schultz, L. W., D. J. Quirk, and R. T. Raines. 1998. His ··· Asp catalytic dyad of ribonuclease A: structure and function of the wild-type, D121N, and D121A enzymes. *Biochemistry*. 37:8886–8898.
- Sela, M., C. B. Anfinsen, and W. F. Harrington. 1957. The correlation of ribonuclease activity with specific aspects of tertiary structure. *Biochim. Biophys. Acta*. 26:502–512.
- Shrager, R. I., J. S. Cohen, S. R. Heller, D. H. Sachs, and A. N. Schechter. 1972. Mathematical models for interacting groups in nuclear magnetic resonance titration curves. *Biochemistry*. 11:541–547.
- Smythe, D. G., W. H. Stein, and S. Moore. 1963. The sequence of amino acid residues in bovine pancreatic ribonuclease: revisions and confirmations. *J. Biol. Chem.* 238:227.
- Tanokura, M. 1983. Proton NMR study on the tautomerism of the imidazole ring of histidine residues. II. Microenvironments of histidine-12 and histidine-119 of bovine pancreatic ribonuclease A. *Biochim. Biophys. Acta*. 742:586–596.
- Thompson, J. E., and R. T. Raines. 1994. Value of general acid-base catalysis to ribonuclease A. *J. Am. Chem. Soc.* 116:5467–5468.
- Thompson, J. E., F. D. Venegas, and R. T. Raines. 1994. Energetics of catalysis by ribonucleases: fate of the 2',3'-cyclic intermediate. *Biochemistry*. 33:7408–7414.
- Udgaonkar, J., and R. Baldwin. 1988. NMR evidence for an early framework intermediate on the folding pathway of ribonuclease A. *Science*. 335:694–699.
- Walba, H., and R. W. Isensee. 1955. Spectrophotometric study of the hydrolysis constants of the negative ions of some aryl imidazoles. *J. Am. Chem. Soc.* 77:5488–5492.
- Wieker, H.-J., and H. Witzel. 1967. Zum mechanismus der ribonuclease-reaktion. 3. Zuordnung der kinetischen parameter k_{+1} , k_{-1} , k_{+2} and interpretation of K_m . *Eur. J. Biochem.* 1:251–258.
- Wladkowski, B. D., L. A. Svensson, L. Sjölin, J. E. Ladner, and G. L. Gilliland. 1998. Structure (1.3 Å) and charge states of a ribonuclease A–uridine vanadate complex: implications for the phosphate ester hydrolysis mechanism. *J. Am. Chem. Soc.* 120:5488–5498.
- Wlodawer, A., M. Miller, and L. Sjölin. 1983. Active site of RNase: neutron diffraction study of a complex with uridine vanadate, a transition-state analog. *Proc. Natl. Acad. Sci. U.S.A.* 80:3628–3631.
- Wyman, J., and S. J. Gill. 1990. *Binding and Linkage*. University Science Books, Mill Valley, CA.