

# A Kinetic Assessment of the Sequence of Electron Transfer from $F_X$ to $F_A$ and Further to $F_B$ in Photosystem I: The Value of the Equilibrium Constant between $F_X$ and $F_A$

Vladimir P. Shinkarev,\* Ilya R. Vassiliev,<sup>†</sup> and John H. Golbeck<sup>†</sup>

\*Department of Plant Biology, University of Illinois, Urbana, Illinois 61801, and <sup>†</sup>Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, Pennsylvania 16802 USA

**ABSTRACT** The x-ray structure analysis of photosystem I (PS I) crystals at 4-Å resolution (Schubert et al., 1997, *J. Mol. Biol.* 272:741–769) has revealed the distances between the three iron-sulfur clusters, labeled  $F_X$ ,  $F_1$ , and  $F_2$ , which function on the acceptor side of PS I. There is a general consensus concerning the assignment of the  $F_X$  cluster, which is bound to the PsaA and PsaB polypeptides that constitute the PS I core heterodimer. However, the correspondence between the acceptors labeled  $F_1$  and  $F_2$  on the electron density map and the  $F_A$  and  $F_B$  clusters defined by electron paramagnetic resonance (EPR) spectroscopy remains controversial. Two recent studies (Diaz-Quintana et al., 1998, *Biochemistry*. 37:3429–3439; Vassiliev et al., 1998, *Biophys. J.* 74:2029–2035) provided evidence that  $F_A$  is the cluster proximal to  $F_X$ , and  $F_B$  is the cluster that donates electrons to ferredoxin. In this work, we provide a kinetic argument to support this assignment by estimating the rates of electron transfer between the iron-sulfur clusters  $F_X$ ,  $F_A$ , and  $F_B$ . The experimentally determined kinetics of P700<sup>+</sup> dark relaxation in PS I complexes (both  $F_A$  and  $F_B$  are present), HgCl<sub>2</sub>-treated PS I complexes (devoid of  $F_B$ ), and P700- $F_X$  cores (devoid of both  $F_A$  and  $F_B$ ) from *Synechococcus* sp. PCC 6301 are compared with the expected dependencies on the rate of electron transfer, based on the x-ray distances between the cofactors. The analysis, which takes into consideration the asymmetrical position of iron-sulfur clusters  $F_1$  and  $F_2$  relative to  $F_X$ , supports the  $F_X \rightarrow F_A \rightarrow F_B \rightarrow \text{Fd}$  sequence of electron transfer on the acceptor side of PS I. Based on this sequence of electron transfer and on the observed kinetics of P700<sup>+</sup> reduction and  $F_X^-$  oxidation, we estimate the equilibrium constant of electron transfer between  $F_X$  and  $F_A$  at room temperature to be  $\sim 47$ . The value of this equilibrium constant is discussed in the context of the midpoint potentials of  $F_X$  and  $F_A$ , as determined by low-temperature EPR spectroscopy.

## INTRODUCTION

Photosystem I (PS I) of oxygenic photosynthesis is a membrane-bound protein-cofactor complex that functions as a light-dependent plastocyanin (or cytochrome  $c_6$ ):ferredoxin (or flavodoxin) oxidoreductase. Light-induced electron transfer takes place in a series of reactions between neighboring electron carriers that are embedded in this protein complex. The electron carriers include a dimeric chlorophyll (Chl), which functions as the primary electron donor (P700); a monomeric chlorophyll, which functions as the primary electron acceptor ( $A_0$ ); an intermediate quinone electron carrier ( $A_1$ ); and three [4Fe-4S] clusters ( $F_X$ ,  $F_A$ , and  $F_B$ ), which operate as the terminal electron acceptors. The cofactors P700,  $A_0$ ,  $A_1$ , and  $F_X$  are bound to the two main polypeptides, PsaA and PsaB, and the terminal electron acceptors  $F_A$  and  $F_B$  are bound to the small PsaC subunit (reviewed in Golbeck, 1995; Brettel, 1997; Fromme, 1999; Manna and Chitnis, 1999).

X-ray structure analysis of PS I crystals at 4 Å resolution (Schubert et al., 1997; Klukas et al., 1999) has revealed the distances between the three iron-sulfur clusters  $F_X$ ,  $F_1$ , and  $F_2$ , which function on the acceptor side of PS I. There is a general consensus on the assignment of the  $F_X$  cluster that is bound to the PsaA and PsaB polypeptides that constitute the PS I core heterodimer. However, the correspondence of the  $F_1$  and  $F_2$  clusters defined by the x-ray data to the  $F_A$  and  $F_B$  clusters defined by their electron paramagnetic resonance (EPR) spectra and cysteine ligands on PsaC remains controversial (reviewed in Brettel, 1997; Kamlowski et al., 1997; Vassiliev et al., 1998). In particular, the definite assignment of  $F_A$  and  $F_B$  to electron density depends on nonsymmetry elements in the polypeptide backbone of PsaC, which are difficult to resolve on the 4-Å map.

The following are the main kinetically based findings concerning the function of the iron-sulfur clusters  $F_A$  and  $F_B$  in PS I:

1. Removal of PsaC by treatment with chaotropic agents leads to the loss of NADP<sup>+</sup> photoreduction. Rebinding of PsaC to isolated P700- $F_X$  core restores NADP<sup>+</sup> photoreduction (Hanley et al., 1992).

2. Treatment of chloroplasts and PS I complexes with HgCl<sub>2</sub>, which leads to selective inactivation of  $F_B$ , inhibits NADP<sup>+</sup> (Fujii et al., 1990), ferredoxin, and flavodoxin photoreduction (Jung et al., 1995; Diaz-Quintana et al., 1998; Vassiliev et al., 1998).

3. Cluster  $F_B$  is the main site of electron donation to methyl viologen (Fujii et al., 1990).

Received for publication 6 July 1999 and in final form 19 October 1999.

Address reprint requests to Dr. John H. Golbeck, Department of Biochemistry and Molecular Biology, Pennsylvania State University, S-310 Frear Laboratory, University Park, PA 16802-4500. Tel.: 1-814-864-1163; Fax: 814-863-7405; E-mail: jhg5@psu.edu.

Dr. Vassiliev is on leave from the Department of Biophysics, Faculty of Biology, M. V. Lomonosov Moscow State University, Moscow 119899, Russia.

© 2000 by the Biophysical Society

0006-3495/00/01/363/10 \$2.00

4. Cluster  $F_A$  can still be photochemically reduced despite the absence of  $F_B$  (Golbeck and Warden, 1982; Fujii et al., 1990) and can donate electrons to methyl viologen added in high concentrations (Fujii et al., 1990; Vassiliev et al., 1998).

5. Both the amplitude and the decay rate of  $A_{430}$  of  $HgCl_2$ -treated particles are very similar to that of untreated particles, with lifetimes in both cases of  $\sim 40$ – $45$  ms (He and Malkin, 1994);

6. The amplitude of the photovoltage change is lower in  $HgCl_2$ -treated PS I complexes than in the presence of  $F_B$  (Mamedov et al., 1998; Diaz-Quintana et al., 1998).

7. In spinach, inactivation of cluster  $F_B$  by  $HgCl_2$  inhibits electron transfer to the water-soluble electron acceptor ferredoxin. However, the restoration of  $F_B$  in PS I does not lead to the restoration of ferredoxin and  $NADP^+$  reduction (He and Malkin, 1994).

8. In cyanobacteria, reconstitution of iron-sulfur cluster  $F_B$  with  $\beta$ -mercaptoethanol, inorganic iron, and sulfide results in restoration of  $NADP^+$  (Jung et al., 1995), ferredoxin, and flavodoxin (Jung et al., 1995; Diaz-Quintana et al., 1998; Vassiliev et al., 1998) photoreduction.

These findings imply that  $F_1 \equiv F_A$  and  $F_2 \equiv F_B$ . A similar conclusion regarding the positions of  $F_A$  and  $F_B$  relative to  $F_X$  has been reached using site-directed mutants with substitutions of the ligands to  $F_A$  and  $F_B$  (Golbeck, 1999) or charged amino acids surrounding  $F_A$  and  $F_B$  (Fischer et al., 1997, 1999).

Although arguments have been put forward which state that  $F_A$  is distal to  $F_X$  (values of redox potentials of  $F_A$  and  $F_B$ , the absence of photoreduction of  $F_A$  in chloroplasts with  $F_B$  modified by diazonium benzene sulfonate; Malkin, 1984, and others; see Brettel, 1997, and Scheller et al., 1997, for discussions), the kinetic data are largely consistent with the assignment of  $F_B$  as the terminal electron acceptor (particularly the requirement of the presence of  $F_B$  for  $NADP^+$  photoreduction and the accessibility of  $F_B$  to exogenous ( $O_2$ , MV) and endogenous (ferredoxin, flavodoxin) acceptors).

A general consideration of the effect of distance on the rate of electron transfer has been applied by Brettel (1997) to the analysis of structure-function relationships in PS I and, in particular, to electron transfer between the iron-sulfur clusters. Here we further elaborate on these considerations to assign the  $F_1$  and  $F_2$  iron-sulfur clusters in PS I to  $F_A$  and  $F_B$ . It is shown that the kinetics of  $P700^+$  dark relaxation in  $Hg$ -treated (only  $F_A$  is present) PS I complexes from *Synechococcus* sp. are consistent with the identification  $F_1 = F_A$ . Thus the analysis of flash-induced kinetics of  $P700^+$  based on rate versus distance relationships strongly supports the following sequence of electron transfer on the acceptor side of PS I:  $F_X \rightarrow F_A \rightarrow F_B \rightarrow Fd$ .

## MATERIALS AND METHODS

### Isolation of PS I complexes

PS I complexes from *Synechococcus* sp. PCC 6301 (TX-PS I) were isolated using Triton X-100 and sucrose gradient ultracentrifugation (Golbeck,

1995). Isolated PS I preparations were resuspended in 50 mM Tris buffer, pH 8.3, with 15% glycerol, frozen as small aliquots in liquid nitrogen and stored at  $-95^\circ C$  before use. The preparation of  $F_B$ -less,  $Hg$ -treated TX-PS I complexes and the reinsertion of the  $F_B$  iron-sulfur cluster were performed as described previously (Jung et al., 1995) by adaptation of the original protocol developed for higher plants (Sakurai et al., 1991) to cyanobacteria. To obtain kinetic confirmation for the removal of a single iron-sulfur cluster, we employed  $\Delta A_{832}$  measurements and a multiple flash excitation protocol developed by Sauer and co-workers (1978). An independent estimation of  $F_A$  and  $F_B$  content is provided by low-temperature EPR spectroscopy. Both approaches showed that treatment of the cyanobacterial PS I complexes with  $HgCl_2$  resulted in 90% destruction of the  $F_B$  iron-sulfur cluster and in the retention of 80% of the  $F_A$  iron-sulfur cluster (data not shown). Damage to the  $F_X$  cluster during the PS I complex isolation and subsequent  $HgCl_2$  treatment can be best assessed by estimating the contribution of the  $F_X$  back-reaction to  $P700^+$  reduction in the presence of methyl viologen. As follows from the kinetics measured in previous work (Vassiliev et al., 1998) for the same material as used in the present work, less than 8% of  $F_A$  and  $F_B$  is missing in the control, and less than 10% of  $F_X$  is damaged in the  $Hg$ -treated sample.

### Time-resolved absorbance spectroscopy

Samples for optical experiments were suspended anaerobically in 25 mM Tris buffer (pH 8.3) in quartz cuvettes with airtight stoppers. Triton X-100 was added to a final concentration of 0.04% to reduce light scattering. 2,6-Dichlorophenol-indophenol (DCPIP), sodium ascorbate, and methyl viologen (all from Sigma, St. Louis, MO) were added where indicated. The solutions were prepared in an anaerobic chamber using oxygen-free distilled water, with air replaced in a Thunberg tube by high-purity nitrogen.

The kinetics of the absorbance changes at 832 nm ( $\Delta A_{832}$ ) and at 811 nm ( $\Delta A_{811}$ ) were measured in a 10 mm  $\times$  4 mm cuvette placed in a laboratory-built spectrophotometer described previously (Vassiliev et al., 1998). The kinetics of the absorbance changes in the visible region were measured with a custom-built single-beam spectrophotometer. A collimated beam derived from a 400-W tungsten bulb was passed through a 5-cm water filter and a Jarrel Ash monochromator (slits set to provide 5-nm bandwidths, FWHM), and the collimated beam was passed through a 10 mm  $\times$  10 mm cuvette containing the sample. An identical monochromator was placed between the sample and a negatively biased photodetector (PIN-10D; United Detector Technology, Hawthorne, CA). The photocurrent was converted to voltage with a 10-k $\Omega$  resistor and amplified 500-fold with an EG&G model 113A amplifier (bandwidth 100 kHz). The kinetics were averaged 32 times and digitized with a Nicolet 4094A oscilloscope interfaced via a NB-GPIB/TNT board (National Instruments, Austin, TX) to a Macintosh 7100/80 computer. To reduce exposure of the sample to actinic light, a Uniblitz VS25 shutter (Vincent Associates, Rochester, NY) was placed between the first monochromator and the sample. The shutter was opened 5 ms in advance of the excitation flash for a total period of 100 ms. In both NIR and visible kinetic measurements, single turnover flashes were provided by a frequency-doubled ( $\lambda$ , 532 nm), Q-switched (FWHM, 10 ns) Nd-YAG laser model DCR-11 (Spectra-Physics, Mountain View, CA) at a flash energy of 10 mJ. The intervals between the flashes were 9 s for the  $P700$ - $F_X$  core and 50 s in all other preparations. The multiexponential fits of  $\Delta A_{832}$  kinetics were performed by the Marquardt algorithm in Igor Pro, version 3.14 (Wavemetrics, Lake Oswego, OR).

## RESULTS

### Kinetics of $P700^+$ dark relaxation in control ( $F_A$ and $F_B$ present) PS I complexes in the presence of a slow donor (DCPIP)

Fig. 1 shows typical kinetic traces of the flash-induced absorbance change at 832 nm in an integral ( $F_A/F_B$ ) TX-PS I




FIGURE 1 Kinetics of absorbance changes at 832 nm in integral ( $F_A/F_B$ ) TX-PS I complexes from *Synechococcus* sp. PCC6301. Reaction medium (anaerobic): 25 mM Tris buffer (pH 8.3), 0.04% Triton X-100, 4  $\mu$ M DCPIP, and 10 mM Na ascorbate. Chl *a* concentration, 50  $\mu$ g/ml. Each component of the multiexponential fit is plotted with a vertical offset relative to the next component (with a longer lifetime) or the baseline; the offset is equal to the amplitude of the latter component.

complex from *Synechococcus* sp. PCC 6301 in the presence of DCPIP and ascorbate. The  $\Delta A_{832}$  kinetics, which reflects the  $P700^+$  dark relaxation, is presented on a logarithmic time scale so that the charge recombination from  $A_1^-$  through  $F_A^-$  can be visualized. The multiexponential fit of these kinetics shows at least five different components with characteristic lifetimes of  $\sim 13 \mu$ s, 642  $\mu$ s, 14 ms, 98 ms, and 3.2 s and amplitudes of  $\sim 14$ , 5.6, 14.5, 40.3, and 25.1%, respectively.

The different components of  $P700^+$  dark relaxation can be identified using preparations missing some or all of the iron-sulfur clusters.  $P700^+$  reduction in  $P700-F_X$  cores (without  $F_A/F_B$ ) is dominated by kinetic components with lifetimes of  $\sim 0.2$  ms and 2 ms, which represent, either directly or indirectly (through  $A_1$ ), the back-reaction from  $F_X^-$  (Vassiliev et al., 1997; see also Figs. 3 and 4 below). Similarly,  $P700^+$  reduction in  $P700-A_1$  cores (without  $F_X$  and  $F_A/F_B$ ) is dominated by kinetic components with lifetimes of  $\sim 10$  and 70  $\mu$ s, which represents, either directly or indirectly (through  $A_0$ ), the back-reaction from  $A_1^-$  (Brettel and Golbeck, 1995). In addition, the decay of the triplet state of chlorophyll (Brettel and Golbeck, 1995; Vassiliev et al., 1997) contributes to the tens-of-microseconds kinetic phase, but this can be differentiated from the  $A_1$  back-reaction by a flash saturation study and a wavelength dependence in the near-IR. Note that the above kinetic assignments only apply to reaction centers with missing iron-sulfur clusters; they do not pertain to reaction clusters with prerduced iron-sulfur clusters, where electrostatic interactions between iron-sulfur clusters may affect the kinetics

and/or midpoint potentials of the acceptors (see Brettel, 1997). Because freshly isolated membranes show flash-induced kinetics with almost exclusively the millisecond and second components present (Vassiliev et al., 1997), the faster phases in TX-PS I complexes most likely result from differential damage to the iron-sulfur clusters in a minority of detergent-isolated PS I complexes. The slower millisecond components ( $\sim 10$  and 200 ms) are due to charge recombination between  $[F_A/F_B]^-$  and  $P700^+$ . The slowest (seconds) component was assigned to the reduction of  $P700^+$  by DCPIP (for discussion see Vassiliev et al., 1997).

### Kinetics of $P700$ dark relaxation in $HgCl_2$ -treated ( $F_B$ -less) PS I complexes in the presence of a slow electron donor for $P700^+$

Fig. 2 shows the kinetic trace of the absorbance change at 832 nm in a TX-PS I complex treated with  $HgCl_2$  to remove the  $F_B$  iron-sulfur cluster. The multiexponential fit of these kinetics shows components with characteristic lifetimes of  $\sim 12 \mu$ s, 208  $\mu$ s, 3 ms, 18 ms, and 112 ms and amplitudes of  $\sim 19.3$ , 8.3, 7.8, 44.2, and 18.2%, respectively. The slowest component is approximated by a baseline that contributes 2.2% to the overall amplitude. Thus the main effect of the  $HgCl_2$  treatment is an increase of the contribution from the millisecond components of the  $P700^+$  dark relaxation at the expense of the slow (seconds) component due to  $P700^+$  reduction by DCPIP. The fraction of these millisecond components is increased from  $\sim 55\%$  in control TX-PS I complexes to  $\sim 70\%$  in a TX-PS I complex treated with  $HgCl_2$ .

The result of the effect of  $HgCl_2$  on the kinetics of  $P700^+$  dark relaxation agrees well with similar results obtained for




FIGURE 2 Kinetics of absorbance change at 832 nm in  $HgCl_2$ -treated PS I complexes from *Synechococcus* sp. PCC6301. The reaction medium is as in Fig. 1.

spinach (He and Malkin, 1994) and *Synechocystis* sp. PCC 6803 (Diaz-Quintana et al., 1998). In previous experiments on  $\text{HgCl}_2$ -treated PS I complexes from *Synechococcus* sp. PCC 6301 (Vassiliev et al., 1997), we reported a significant amplitude of the submillisecond component due to electron transfer from  $\text{F}_X^-$  to  $\text{P700}^+$  in RCs with a damaged  $\text{F}_A$  cluster. This component is practically absent (<10%) in preparations used in the present (this work) and previous (Vassiliev et al., 1998) studies, attesting to the low amount of damage to  $\text{F}_A$  in these preparations.

### Kinetics of dark relaxation in PS I core ( $\text{F}_A/\text{F}_B$ -less) complexes

To monitor the kinetics of the back-reaction between  $\text{F}_X^-$  and  $\text{P700}^+$  we studied core PS I complexes where both  $\text{F}_A$  and  $\text{F}_B$  are absent. Fig. 3 shows the multiexponential fit of the kinetics of  $\text{P700}^+$  dark relaxation in  $\text{P700-F}_X$  core complexes measured at 811 nm. The observed absorbance change has components with characteristic times of  $\sim 9 \mu\text{s}$ ,  $103 \mu\text{s}$ ,  $576 \mu\text{s}$ , and  $3.6 \text{ ms}$  and amplitudes  $\sim 9.2$ ,  $7.6$ ,  $71.2$ , and  $7.2\%$ , respectively, plus a baseline contributing  $\sim 4.8\%$ . To definitely assign components of the  $\text{P700}^+$  dark relaxation to its reaction with  $\text{F}_X^-$  and to exclude  $\text{A}_1^-$  as a possible contributor to the observed kinetics, we measured the flash-induced changes in the blue region where the  $\text{F}_X$  iron-sulfur cluster absorbs (Parrett et al., 1989; Franke et al., 1995).

Both  $\text{P700}$  and  $\text{F}_X$  contribute to absorbance changes in the blue region. Their individual differential spectra can be obtained by using methyl viologen, which functions as an electron acceptor, preventing the back-reaction between  $\text{F}_X^-$  and  $\text{P700}^+$  (Yu et al., 1995). In the spectral region from 400 to 480 nm, we found virtually no absorbance changes at 415




FIGURE 3 Kinetics of absorbance change at 811 nm in  $\text{P700-F}_X$  core complexes from *Synechococcus* sp. PCC6301. The reaction medium is as in Fig. 1.

nm (Fig. 4) and 445 nm (not shown) in the presence of methyl viologen. Therefore these wavelengths represent the cross-over points of the  $\text{P700/P700}^+$  difference spectrum. The absorbance change measured at these points in the absence of methyl viologen should be ascribed entirely to a change in the  $\text{F}_X$  oxidation state. Fig. 4 shows a multiexponential fit of kinetics measured at 415 nm. The major decay phase has a lifetime of  $550 \mu\text{s}$  (61.4%); it is preceded by a  $153\text{-}\mu\text{s}$  phase (23.9%) and followed by a  $3\text{-ms}$  phase (10.6%) and a slower decaying component approximated with a baseline (4.1%).

The main difference between the measurement of  $\text{P700-F}_X$  core complexes at 415 nm and 811 nm is the presence of a fast  $9\text{-}\mu\text{s}$  component at 811 nm. As we have shown for both  $\text{P700-F}_X$  core and integral PS I complexes, components with lifetimes of  $\sim 10 \mu\text{s}$  are not saturated with flash energies that saturate the major (hundreds of microseconds to milliseconds) components and hence correspond to the decay of the triplet state of chlorophyll (Vassiliev et al., 1997). It follows that the major kinetic components at 811 nm ( $576 \mu\text{s}$  and  $3.6 \text{ ms}$ ) and 415 nm ( $550 \mu\text{s}$  and  $3 \text{ ms}$ ) arise from the reaction between  $\text{F}_X^-$  (directly or via  $\text{A}_1$ ) and  $\text{P700}^+$ . Note that the  $250\text{-}\mu\text{s}$  kinetic phase of NIR absorbance change in PS I complexes with  $\text{F}_A$  and  $\text{F}_B$  (but not  $\text{F}_X$ ) prereduced by dithionite before the flash was attributed to  $\text{A}_1^-$  back-reaction and was explained assuming that the  $\text{P700}^+\text{F}_X^-$  state is lower in free energy than the  $\text{P700}^+\text{A}_1$  state when  $\text{F}_A$  and  $\text{F}_B$  are prereduced (Brettel, 1989, 1997). This consideration does not apply to the experimental data presented in this paper on PS I complexes devoid of  $\text{F}_A$  and  $\text{F}_B$ .




FIGURE 4 Kinetics of absorbance change at 415 nm in  $\text{P700-F}_X$  core complexes from *Synechococcus* sp. PCC6301 in the absence (bottom curve) and in the presence (curve in the center) of 280 mM methyl viologen. The reaction medium is as in Fig. 1, but the Chl *a* concentration was  $8 \mu\text{g/ml}$ .

It is generally assumed that the monomolecular back-reaction between a particular acceptor and P700 should follow monoexponential kinetics, but in many instances (see Figs. 1–4) the experimental data are best fitted by two or more components. The nature of this heterogeneity in the dark relaxation of  $P700^+$  is not fully understood, but it may be related to the existence of different conformational (sub) states in the reaction clusters. Such a heterogeneity has been reported in PS I (Schlodder et al., 1998) and RCs from purple bacteria (McMahon et al. 1998).

In the following calculations, we use the average time determined as  $(\tau_1 A_1 + \tau_2 A_2 + \tau_3 A_3)/(A_1 + A_2 + A_3)$ , where  $\tau_i$  are the lifetimes and  $A_i$  are the amplitudes of the single exponentials, corresponding to the same reaction of  $P^+$  dark reduction. Using the multiexponential analysis data from Figs. 3 and 2, we estimate the average lifetimes to be used in further calculations:  $\tau_{XP} \approx 0.85$  ms [=  $(0.576 \cdot 71.2 + 3.6 \cdot 7.2)/(71.2 + 7.2)$ ] for the  $F_X^- \rightarrow P700^+$  back-reaction in P700- $F_X$  cores and  $\tau_{AP} \approx 40.7$  ms [=  $(3 \cdot 7.8 + 18 \cdot 44.2 + 112 \cdot 18.2)/(7.8 + 44.2 + 18.2)$ ] for the  $F_A^- \rightarrow P700^+$  back-reaction in  $HgCl_2$ -treated ( $F_B$ -less) PS I complexes.

## DISCUSSION

### Data from x-ray structural analysis of PS I and low-molecular-mass [4Fe-4S] ferredoxins

The preliminary x-ray structure analysis of crystals of PS I depicts the geometry of iron-sulfur clusters on the acceptor side of PS I (Schubert et al., 1997; Fromme, 1999; Klukas et al., 1999). The center-to-center distance is 15 Å between  $F_X$  and  $F_1$  and 12 Å between  $F_1$  and  $F_2$ . The x-ray data do not allow one to determine the identity of  $F_1$  and  $F_2$  (see Brettel, 1997, and Kamlowski et al., 1997, for a full discussion). The distance between  $F_1$  and  $F_2$  can be modeled by the bacterial ferredoxins for which precise x-ray structural analysis has been carried out. For example, in the [4Fe-4S] ferredoxin from *Peptococcus asacharolyticus* (formerly *P. aerogenes*; Adman et al., 1976), the closest edge-to-edge distance between the iron atoms in different clusters is 8 Å, while the center-to-center distance is 12 Å, which is in good agreement with the 12-Å distance determined between clusters  $F_1$  and  $F_2$  in PS I.

### Application of the relationship between electron transfer rate and distance among the electron carriers on the acceptor side of PS I

The rate of electron transfer between electron carriers decreases exponentially with distance, and this dependence has been tabulated for different reactions in proteins (see, for example, Likhtenshtein, 1988; Moser et al., 1995; Gray and Winkler, 1996). The most detailed “ruler” has been suggested by Dutton and co-workers (Moser et al., 1995), who deduced this relationship from an analysis of electron

transfer in photosynthetic proteins. According to this formulation (Moser et al., 1995), the logarithm of the rate constant of intraprotein electron transfer between two electron carriers with edge-to-edge distance  $R$  can be described by the following equation:

$$\log k = 15 - 0.6R - 3.1(\Delta G^0 + \lambda)^2/\lambda \quad (1)$$

where  $R$  is the distance in Å,  $\Delta G^0$  is the standard reaction free energy in eV, and  $\lambda$  is the reorganization energy in eV. The most essential consequence of this dependency is a one-order change in the rate of electron transfer in proteins with a change of distance of 1.7 Å.

Schlodder et al. (1998) estimated from the temperature dependence of electron transport that the reorganization energy for the reaction between  $A_1$  and  $F_X$  has a value of  $\sim 1$  eV. In the following calculations, we assume  $\lambda = 1$  eV for reactions between all iron-sulfur clusters in PS I.

By applying Eq. 1 to the acceptor side of PS I, we can estimate (Fig. 5) that the rate of electron transfer between  $F_X$  and  $F_1$  and between  $F_1$  and  $F_2$  is faster than 10  $\mu$ s (see also Brettel, 1997). The distance between  $F_1$  and  $F_2$  and P700 exceeds 35 Å, which requires the experimentally observed millisecond reduction of  $P700^+$  by these acceptors to occur indirectly via thermal repopulation of  $F_X$ . There-




FIGURE 5 Dependence of the logarithm of the rate of electron transfer in biological systems on the distance between cofactors described by Eq. 1 and the suggested rate constants of electron transfer between the acceptors of PS I, based on data of x-ray structural analysis. Values of  $\lambda = 1$  eV and  $\Delta G^0 = 0$  eV were used in calculations to draw the theoretical line. The edge-to-edge distance between iron sulfur centers (indicated by vertical lines) is assumed to be equal to the center-center distance minus 4 Å. Inset: Scheme of electron transfer in the PS I derived from application of the correlation between rate of electron transfer and x-ray distances between the cofactors to the data of x-ray structural analysis. The rates of electron transfer from  $F_1$  to P700 and from  $F_2$  to P700 are significantly slower than back-reactions via  $F_X$  (directly or via  $A_1$ ).

fore the distances between the cofactors strongly support a linear chain of electron transfer between  $F_X$ ,  $F_1$ , and  $F_2$  during the dark relaxation of  $P700^+$ :





Fig. 5 (*inset*) shows the scheme of electron transfer in PS I that takes into account these considerations.

### Identification of $F_1$ as $F_A$ , based on an analysis of the kinetics of $P700^+$ dark relaxation in PS I complexes containing only $F_A$

We now analyze in detail the kinetics of electron transfer between the iron-sulfur clusters of PS I. We make no a priori assumptions about the sequence of electron transfer; rather we consider the two possible positions of  $F_A$  and  $F_B$ . The linearity of electron transfer between  $F_X$ ,  $F_1$ , and  $F_2$  and the noncontroversial identification of  $F_X$  leads to only two alternatives, either  $F_A \equiv F_1$  or  $F_A \equiv F_2$ . Depending on the assumption about the identity of the  $F_A$  cluster, the kinetics of the dark relaxation of  $P700$  should be different because of the asymmetrical positions, and hence different distances, of  $F_1$  and  $F_2$  relative to  $F_X$ . The results of our theoretical calculations with experimental observations are compared below.

*Assumption that  $F_1 \equiv F_A$*

Let us assume, first, that  $F_A \equiv F_1$ . In this case the center-to-center distance between  $F_X$  and  $F_1 \equiv F_A$  is short,  $\sim 15 \text{ \AA}$ , and the edge-to-edge distance can be as short as  $11 \text{ \AA}$ . Applying the rate versus distance dependency to this case (Fig. 5), we find that the rate constant of electron transfer between these clusters is in the range of  $10^5$ – $10^6 \text{ s}^{-1}$ , i.e., this electron transfer is significantly faster than the back-reaction from  $F_X^-$  to  $P700^+$  ( $\tau \approx 0.85 \text{ ms}$  in  $P700$ - $F_X$  core complexes). In this case electron transfer in Hg-treated PS I complexes during dark relaxation can be represented by the following scheme (we ignore the very small minority of PS I complexes with damaged  $F_A$ ):



Here P stands for  $P700$ .  $k_{XP}$ ,  $k_{X1}$ , and  $k_{1X}$  are the respective rate constants of electron transfer. The system of differential equations describing the kinetics of electron transfer according to Scheme 3 is

$$\begin{cases} \frac{d[P^+ F_X^- F_1]}{dt} = -(k_{XP} + k_{X1})[P^+ F_X^- F_1] + k_{1X}[P^+ F_X F_1^-] \\ \frac{d[P^+ F_X F_1^-]}{dt} = k_{X1}[P^+ F_X^- F_1] - k_{1X}[P^+ F_X F_1^-] \end{cases} \quad (4)$$

Its solution, with the initial conditions  $[P^+ F_X^- F_1](0) = 1$  and  $[P^+ F_X F_1^-](0) = 0$ , can be written in the following general form:

$$[P^+ F_X^- F_1] = \frac{\eta + k_{1X}}{\eta - \mu} e^{\eta t} - \frac{\mu + k_{1X}}{\eta - \mu} e^{\mu t} \quad (5)$$

$$[P^+ F_X F_1^-] = \frac{k_{X1}}{\eta - \mu} e^{\eta t} - \frac{k_{X1}}{\eta - \mu} e^{\mu t}$$

where  $\eta = -\sigma - \sqrt{\sigma^2 - \rho}$ ,  $\mu = -\sigma + \sqrt{\sigma^2 - \rho}$ , and  $\sigma = (k_{XP} + k_{1X} + k_{X1})/2$ ,  $\rho = k_{1X}k_{XP}$ . When  $\sigma^2 \gg \rho$ , the  $\eta$  and  $\mu$  can be approximated by the following simple expressions:

$$\eta \approx -2\sigma; \quad \mu \approx -\rho/2\sigma \quad (6)$$

From Eq. 5 it follows that the normalized kinetics of the dark relaxation of flash-induced  $P^+$  ( $\equiv [P^+ F_X^- F_1] + [P^+ F_X F_1^-]$ ) according to Scheme 3 is described by two exponential components ( $F$  stands for the relative amplitude of the fast component, and  $S$  stands for the relative amplitude of the slow component):

$$[P^+] = \frac{\eta + k_{1X} + k_{X1}}{\eta - \mu} e^{\eta t} - \frac{\mu + k_{1X} + k_{X1}}{\eta - \mu} e^{\mu t} \equiv F e^{\eta t} + S e^{\mu t} \quad (7)$$

According to Eq. 7 the fraction of the fast component of  $P700^+$  dark relaxation is

$$F = (\eta + k_{X1} + k_{1X})/(\eta - \mu) \quad (8)$$

For the case considered here ( $F_A \equiv F_1$ ) the estimated rate constant of electron transfer between  $F_X$  and  $F_1$  ( $10^5$ – $10^6 \text{ s}^{-1}$ ) is significantly larger than the rate constant of the back-reaction from  $F_X^-$  to  $P700^+$  ( $10^3$ – $10^4 \text{ s}^{-1}$ ), i.e.,  $k_{X1} \gg k_{XP}$ . From Eq. 6 it follows that in this case  $\eta \approx -(k_{X1} + k_{1X} + k_{XP})$ ,  $\mu \approx -k_{XP}k_{1X}/(k_{X1} + k_{1X})$ . Using these approximations and Eq. 8, we find that the fraction of the fast component is very small:

$$F \approx k_{XP}/(k_{X1} + k_{1X}) \ll 1 \quad (9)$$

Thus the kinetics of  $P700^+$  dark relaxation according to Scheme 3 is described by a “slow” exponent with a lifetime,

$$\tau_{sl} \equiv -1/\mu \approx (k_{X1} + k_{1X})/(k_{XP}k_{1X}) \quad (10)$$

This lifetime of  $P700^+$  dark relaxation can be derived by assuming equilibrium between iron-sulfur clusters  $F_X$  and  $F_1$  during dark relaxation (see, for example, Shinkarev and Wraight, 1993).

From Eq. 10 it follows that the time of the slow millisecond component (2–200 ms) of  $P700^+$  dark relaxation,  $\tau_{sl}$ , is determined by the lifetime of electron transfer from  $F_X$  to  $P700$ ,  $\tau_{XP}$ , and by the equilibrium constant  $L_{X1}$  ( $= k_{X1}/k_{1X}$ ) responsible for redistribution of the electron


between  $F_X$  and  $F_1 \equiv F_A$ :

$$\tau_{sl} = \tau_{XP}(1 + L_{X1}) \quad (11)$$

The fast partitioning of the electron between  $F_X$  and  $F_A$  in Scheme 3 leads to the absence of any significant  $F_X$  component ( $< 2$  ms). This agrees well with measured kinetics of  $P700^+$  dark relaxation in PS I complexes treated with  $HgCl_2$  (Fig. 2), where the slow millisecond  $F_A/F_B$  component (2–200 ms) is  $\sim 70\%$ .

*Assumption that  $F_A \equiv F_2$*

Let us now assume that  $F_A \equiv F_2$ . In this case  $F_1 = F_B$  is damaged by  $HgCl_2$ , and the distance between  $F_X$  and  $F_A$  will be 22 Å (18-Å edge-to-edge distance). According to Fig. 5 this distance corresponds to an optimum rate of electron transfer in the region of  $10^1$  to  $10^2$  s $^{-1}$ , i.e., lifetimes of  $\sim 10$ –100 ms. The electron transfer between  $F_X$  and  $F_A = F_2$  in this case is essentially slower than the back-reaction between  $F_X^-$  and  $P700^+$ , and we can no longer assume a fast equilibrium between  $F_X$  and  $F_2$ . The scheme of electron transfer in PS I after a single flash will be as follows:



This scheme is analogous to Scheme 3 considered above. Therefore, the general solution for  $P700^+$  dark relaxation (Eqs. 5–8) is valid here, too, if one replaces  $k_{X1}$  with  $k_{X2}$  and  $k_{1X}$  with  $k_{2X}$ . However, in the case considered here the rate of electron transfer between  $F_X$  and  $P700$  is essentially larger than the rate of electron transfer between  $F_X$  and  $F_A$ , i.e.,  $k_{XP} \gg k_{X2}$ . In this case  $\eta \approx -(k_{XP} + k_{2X} + k_{X2})$ ,  $\mu \approx -k_{2X}$ , and the fraction of the fast component is close to 1:

$$F \approx \frac{k_{XP}}{k_{XP} + k_{X2}} \approx 1 \quad (13)$$

Thus, for the case considered here, the fast component will have a time  $\approx \tau_{XP}$  and a relative amplitude of  $\tau_{X2}/(\tau_{X2} + \tau_{XP}) \approx 1$ . Using  $\tau_{X2} \approx 10$ –100 ms (estimated from Eq. 1 or from Fig. 5) and  $\tau_{XP} \approx 0.85$  ms, we can estimate that the fraction of the fast component must be greater than 0.92 and the slow component smaller than 0.08. Thus we see that the assumption that  $F_A \equiv F_2$  leads to the conclusion that the amplitude of the slow ( $\sim 10$  and 200 ms) components should be practically zero. This contradicts the experimental data on the kinetics of  $P700^+$  dark relaxation in  $HgCl_2$ -treated PS I complexes presented in Fig. 2. The fraction of the millisecond components ( $\sim 10$  and 200 ms) of  $P700^+$  dark relaxation in these preparations is  $\sim 70\%$ . This fraction will become even higher after exclusion of the fastest component (12  $\mu$ s) that arises from either  $^3Chl$  formation or from damage to some of the PS I clusters. The presence of

a significant fraction of slow component(s) ( $> 70\%$ ) coincides well with other results (Fujii et al., 1990; He and Malkin, 1994; Vassiliev et al., 1997; Diaz-Quintana et al., 1998).

Similarly, this scheme predicts that the relative amplitude of the fast component must be greater than 92%. Our experiments (Fig. 2) show that no more than  $\sim 8.3\%$  of the submillisecond (0.2–2 ms) phase is present in the kinetics of PS I after treatment with  $HgCl_2$ . This is significantly less than 92%, which should be observed in this case.

Therefore, we must draw the conclusion that the identity  $F_A \equiv F_2$  suggested above is inconsistent with the experimental data. Only the identity  $F_A \equiv F_1$  can explain the absence of a significant fast submillisecond (0.2–2 ms)  $F_X$  component and the presence of a significant millisecond (2–200 ms)  $F_A/F_B$  component, providing a reasonable, non-contradictory description of the dark relaxation of  $P700^+$  observed experimentally in  $HgCl_2$ -treated PS I complexes. Even with the large uncertainties in the theoretical treatment (the value of  $\lambda$ , the geometry of the cubane clusters, and hence the precise edge-to-edge distance), we suggest that the correlation between the rate of electron transfer and distances between cofactors and the theoretical analysis of electron transfer in PS I strongly supports the following sequence of electron transfer:  $F_X \rightarrow F_A \rightarrow F_B \rightarrow Fd$ . The above considerations are summarized in the flow chart shown in Fig. 6.

From the chart in Fig. 6 one can see that the above conclusion for the sequence of electron transfer on the acceptor side of PS I will be valid insofar as the rate constant of electron transfer between  $F_X^-$  and  $P^+$  is larger than the rate constant for the electron transfer between  $F_X$  and  $F_2$  and less than the rate constant of electron transfer between  $F_X$  and  $F_1$ , i.e.,  $k_{X1} > k_{XP} > k_{X2}$ , or

$$\log(k_{X1}) > \log(1/\tau_{XP}) > \log(k_{X2}) \quad (14)$$

Assuming the validity of Eq. 1, we can solve the latter inequality for  $(\Delta G^\circ + \lambda)^2/\lambda$ :

$$0.36 \leq (\Delta G^\circ + \lambda)^2/\lambda \leq 1.72 \quad (15)$$

Assuming that  $\Delta G^\circ = -0.1$  eV, we have the following range for reorganization energies for which this inequality will hold (we used here the relationship  $(\Delta G^\circ + \lambda)/\lambda \approx \lambda + 2\Delta G^\circ$ , which is valid when  $|\lambda| \gg |\Delta G^\circ|$ ):  $0.56 \leq \lambda \leq 1.92$ .

Thus our conclusion about the sequence of electron transport will hold for a wide range of reorganization energies.

### Estimation of equilibrium constant of electron transfer between $F_X$ and $F_A$

The established sequence of electron transfer between the iron-sulfur clusters in PS I allows one to determine the value of the equilibrium constant of electron transfer between  $F_X$




FIGURE 6 The flow chart of the logic for determining the sequence of the electron transfer in the PS I, based on the analysis of the kinetics of P700<sup>+</sup> dark relaxation.

and F<sub>A</sub> from the kinetics of P700<sup>+</sup> dark relaxation. According to Eq. 11 the equilibrium constant of electron transfer between F<sub>X</sub> and F<sub>A</sub> ( $L_{XA}$ ) can be estimated from the values of the lifetimes of P700<sup>+</sup> dark reduction by F<sub>A</sub><sup>-</sup> ( $\tau_{sl}$ ) and by

F<sub>X</sub><sup>-</sup> ( $\tau_{XP}$ ):

$$L_{XA} = \frac{\tau_{sl} - \tau_{XP}}{\tau_{XP}} \approx \frac{\tau_{sl}}{\tau_{XP}} \quad (16)$$

Based on the kinetics of the P700-F<sub>X</sub> core complex (Fig. 3), we estimate that  $\tau_{XP} \approx 0.85$  ms. Based on the P700<sup>+</sup> reduction kinetics of the HgCl<sub>2</sub>-treated PS I complex (Fig. 2), we estimate that  $\tau_{AP} \approx 41$  ms. Thus  $L_{XA} = (41 - 0.85)/0.85 \approx 47$ . This value corresponds to a  $\sim 100$ -mV difference between the midpoint potentials of F<sub>X</sub> and F<sub>A</sub> at room temperature. Note that this calculation uses data from preparations where F<sub>B</sub> and where F<sub>A</sub>/F<sub>B</sub> are missing, thereby avoiding problems with electrostatic interaction from prerduced electron acceptors. A considerably higher value of this equilibrium constant is estimated based on the generally accepted midpoint potentials of the iron-sulfur clusters (F<sub>X</sub>, -705 mV (Chamorovsky and Cammack, 1982); F<sub>A</sub>, -540 mV (Evans and Heathcote, 1980)), measured using EPR at low temperatures:  $L_{XA} = 10^{(705 - 540)/60} \approx 562$ .

This discrepancy cannot be resolved by using the lifetime of the main (largest) component of the P<sup>+</sup> reduction by F<sub>X</sub><sup>-</sup> instead of the average time used above for the calculation of equilibrium constant  $L_{XA}$ . Indeed, according to the data in Fig. 4 this time is equal to 0.55 ms, which corresponds to an equilibrium constant  $L_{XA} = (41 - 0.55)/0.55 \approx 73.5$ .

One possible reason for the discrepancy is that the midpoint potential of F<sub>X</sub> may be affected by electrostatic interaction with F<sub>A</sub><sup>-</sup> and F<sub>B</sub><sup>-</sup>. This potential was determined by low-temperature EPR under conditions where both F<sub>A</sub> and F<sub>B</sub> were prerduced (Chamorovsky and Cammack, 1982). A proper comparison with the kinetics of P700<sup>+</sup> dark relaxation should use the measurement of midpoint potential of F<sub>X</sub> under conditions where both F<sub>A</sub> and F<sub>B</sub> are oxidized. The influence of an electrostatic interaction between the iron-sulfur clusters on the kinetics and thermodynamics of electron transfer is consistent with the fact that the F<sub>X</sub> back-reaction, measured under conditions where both F<sub>A</sub> and F<sub>B</sub> are reduced by dithionite, is consistently faster than when measured in a P700-F<sub>X</sub> core complex. In control and HgCl<sub>2</sub>-treated TX-PS I complexes in the presence of dithionite at pH 10, where F<sub>A</sub> and F<sub>B</sub> (when present) are both reduced, the  $\tau_{XP} \approx 0.35$  ms and 0.56 ms, respectively (data not shown).

This agrees well with the lower value of the midpoint potential of F<sub>X</sub> (-670 mV) estimated for the P700-F<sub>X</sub> core complex by transient optical spectroscopy (Parrett et al., 1989). The equilibrium constant estimated using the latter value of the midpoint potential of F<sub>X</sub> ( $L_{XA} = 10^{(670 - 540)/60} \approx 147$ ) is closer to the value estimated here from the kinetics of P700<sup>+</sup> dark relaxation at room temperature. The other reason for the discrepancy may be a possible temperature dependence of the equilibrium constant of electron transfer between F<sub>X</sub> and F<sub>A</sub>.



## Rationale for uphill electron transfer on the acceptor side of PS I

The values of midpoint redox potentials of iron-sulfur clusters in PS I ( $-540$  mV for  $F_A$  and  $-590$  mV for  $F_B$ ) indicate the presence of an uphill electron transfer between  $F_A$  and  $F_B$ . This needs to be examined from a functional point of view. Such an uphill electron transfer step assumes that the redox equilibrium between  $F_A$  and  $F_B$  will be shifted toward  $F_A$ . Low-potential exogenous acceptors of electrons in PS I can interact with oxygen and produce superoxide  $O_2^-$ . The interaction of  $F_A$  with oxygen is slower than with  $F_B$  (at least in the presence of MV) (Fujii et al., 1990; Vassiliev et al., 1998). Thus redistribution of the electron to the  $F_A$  cluster will decrease the rate of electron donation from  $F_B$  to oxygen when ferredoxin is not in its binding site and should thereby maintain a high quantum yield of  $NADP^+$  reduction relative to formation of  $O_2^-$ . It may also protect the reaction center from oxidative destruction by  $O_2^-$ . At the same time, the total driving force for electron transfer to ferredoxin bound to the  $F_B$  site is energetically favorable and will be sufficient for nearly irreversible transfer of the electron to ferredoxin.

Using an equation similar to Eq. 16, we can estimate the equilibrium constant of electron transfer between  $F_A$  and  $F_B$  at room temperature as  $0.8 \leq L_{AB} \leq 4.5$ . The range of values of the equilibrium constant is determined by the range of average lifetimes of the main components of  $P700^+$  dark relaxation. This estimate shows that the energy difference between  $F_A$  and  $F_B$  at room temperature is less than that predicted by the values of the redox potentials. This is likely because the redox potential of  $F_B$  was experimentally determined by EPR in the presence of reduced  $F_A$ . The redox potentials of  $F_A$  and  $F_B$  were also determined under conditions where ferredoxin (or flavodoxin) is not bound to the PS I complexes. Binding of ferredoxin (or flavodoxin) may change the equilibrium constant of electron transfer between  $F_A$  and  $F_B$ . We will report elsewhere a study of the thermodynamics of electron transport between  $F_A$  and  $F_B$  at room temperature in an attempt to further explore the issue of uphill electron transfer on the acceptor side of PS I.

## CONCLUSIONS

We show that the kinetics of  $P700^+$  dark relaxation in the presence of  $F_A$  only and in the presence of both  $F_A$  and  $F_B$  can be used to identify the sequence of electron transfer through the iron-sulfur clusters  $F_A$  and  $F_B$ . By applying the dependence of the electron transfer rate versus distance to the kinetics of electron transfer in PS I, and by taking into consideration the asymmetrical position of iron-sulfur clusters  $F_A$  and  $F_B$  relative to  $F_X$ , we are able to determine that the sequence of electron transfer is  $F_X \rightarrow F_A \rightarrow F_B (\rightarrow Fd)$ . Using this sequence of electron transfer during  $P700^+$  dark

relaxation, we estimate that the equilibrium constant between  $F_X$  and  $F_A$  at room temperature is  $\sim 47$  in  $HgCl_2$ -treated ( $F_B$ -less) PS I complexes.

Note added in proof: On the basis of a comparison between experimental and theoretical values of spin relaxation enhancement effects on  $P700^+$  in PS I particles containing and lacking the  $F_B$  cluster, Lakshmi, Jung, Golbeck, and Brudvig recently showed that iron-sulfur cluster  $F_A$  is closer to  $P700$  than the  $F_B$  cluster. This agrees with the orientation for  $F_A$  and  $F_B$  determined in the present study. *Biochemistry* 1999, 38: 13210–13215.

This work was supported by a grant to JHG from the National Science Foundation (MCB-9723661).

## REFERENCES

- Adman, E. T., L. C. Sieker, and L. H. Jensen. 1976. Structure of *Peptococcus aerogenes* ferredoxin. I. Refinement at 2 Å resolution. *J. Biol. Chem.* 251:3801–3806.
- Brettel, K. 1989. New assignment for the 250 μs kinetics of photosystem I:  $P-700^+$  recombines with  $A_1^-$  (not  $F_X^-$ ). *Biochim. Biophys. Acta.* 976:246–249.
- Brettel, K. 1997. Electron transfer and arrangement of the redox cofactors in photosystem I. *Biochim. Biophys. Acta.* 1318:322–373.
- Brettel, K., and J. H. Golbeck. 1995. Spectral and kinetic characterization of electron acceptor  $A_1$  in a photosystem I core devoid of iron-sulfur centers  $F_X$ ,  $F_B$  and  $F_A$ . *Photosynth. Res.* 45:183–193.
- Chamorovsky, S. K., and R. Cammack. 1982. Direct determination of the midpoint potential of the acceptor X in chloroplast photosystem I by electrochemical reduction and ESR spectroscopy. *Photobiophys.* 4:195–200.
- Diaz-Quintana, A., W. Leibl, H. Bottin, and P. Sétif. 1998. Electron transfer in photosystem I reaction centers follows a linear pathway in which iron-sulfur cluster  $F_B$  is the immediate electron donor to soluble ferredoxin. *Biochemistry.* 37:3429–3439.
- Evans, M. C. W., and P. Heathcote. 1980. Effects of glycerol on the redox properties of the electron acceptor complex in spinach photosystem I particles. *Biochim. Biophys. Acta.* 590:89–96.
- Fischer, N., P. Sétif, and J. D. Rochaix. 1997. Targeted mutations in the *psaC* gene of *Chlamydomonas reinhardtii*: preferential reduction of  $F_B$  at low temperature is not accompanied by altered electron flow from photosystem I ferredoxin. *Biochemistry.* 36:93–102.
- Fischer, N., P. Sétif, and J. D. Rochaix. 1999. Site-directed mutagenesis of the *PsaC* subunit of photosystem I— $F_B$  is the cluster interacting with soluble ferredoxin. *J. Biol. Chem.* 274:23333–23340.
- Franke, J. F., L. Ciesla, and J. T. Warden. 1995. Kinetics of *PsaC* reduction in photosystem I. In *Photosynthesis: From Light to Biosphere*, Vol. 2. P. Mathis, editor. Kluwer Academic Publishers, Dordrecht, the Netherlands. 75–78.
- Fromme, P. 1999. Biology of photosystem I: structural aspects. In *Concepts in Photobiology: Photosynthesis and Photomorphogenesis*. G. S. Singhal, G. Renger, S. K. Sopory, K. D. Irrgang, and Govindjee, editors. Narosa Publishing House, New Delhi. 181–220.
- Fujii, T., E. I. Yokoyama, K. Inoue, and H. Sakurai. 1990. The sites of electron donation of photosystem I to methyl viologen. *Biochim. Biophys. Acta.* 1015:41–48.
- Golbeck, J. H. 1995. Resolution and reconstitution of photosystem I. In *CRC Handbook of Organic Photochemistry and Photobiology*. P. S. Song and W. M. Horspool, editors. CRC Press, Boca Raton, FL. 1407–1419.
- Golbeck, J. H. 1999. A comparative analysis of the spin state distribution of in vitro and in vivo mutants of *PsaC*. A biochemical argument for the sequence of electron transfer as  $F_X \rightarrow F_A \rightarrow F_B \rightarrow$  ferredoxin. *Photosynth. Res.* 61:107–144.

- Golbeck, J. H., and J. Warden. 1982. Electron spin resonance studies of the bound iron-sulfur centers in photosystem I. Photoreduction of center A occurs in the absence of center B. *Biochim. Biophys. Acta.* 681:77–84.
- Gray, H. B., and J. R. Winkler. 1996. Electron transfer in proteins. *Annu. Rev. Biochem.* 65:537–561.
- Hanley, J., J. Kear, G. Bredenkamp, G. Li, P. Heathcote, and M. C. W. Evans. 1992. Biochemical evidence for the role of the bound iron-sulfur centre-B in NADP reduction by photosystem I. *Biochim. Biophys. Acta.* 1099:152–156.
- He, W. Z., and R. Malkin. 1994. Reconstitution of iron-sulfur center B of photosystem I damaged by mercuric chloride. *Photosynth. Res.* 41: 381–388.
- Jung, Y. S., L. Yu, and J. H. Golbeck. 1995. Reconstitution of iron-sulfur center F<sub>A</sub> results in complete restoration of NADP(+) photoreduction in Hg-treated Photosystem I complexes from *Synechococcus* sp. PCC 6301. *Photosynth. Res.* 46:249–255.
- Kamlowski, A., A. Van der Est, P. Fromme, N. Krauss, W. D. Schubert, O. Klukas, and D. Stehlik. 1997. The structural organization of the psaC protein in photosystem I from single crystal EPR and x-ray crystallographic studies. *Biochim. Biophys. Acta.* 1319:199–213.
- Klukas, O., W. D. Schubert, P. Jordan, N. Krauss, P. Fromme, H. T. Witt, and W. Saenger. 1999. Photosystem I, an improved model of the stromal subunits PsaC, PsaD, and PsaE. *J. Biol. Chem.* 274:7351–7360.
- Likhtenshtein, G. I. 1988. Chemical Physics of Redox Metalloenzyme Catalysis. Springer-Verlag, Berlin.
- Malkin, R. 1984. Diazonium modification of photosystem I. A specific effect on iron-sulfur center B. *Biochim. Biophys. Acta.* 764:63–69.
- Mamedov, M. D., K. N. Gourovskaya, I. R. Vassiliev, J. H. Golbeck, and A. Y. Semenov. 1998. Electrogenicity accompanies photoreduction of the iron-sulfur clusters F<sub>A</sub> and F<sub>B</sub> in photosystem I. *FEBS Lett.* 431: 219–223.
- Manna, P., and P. R. Chitnis. 1999. Function and molecular genetics of photosystem I. In *Concepts in Photobiology: Photosynthesis and Photomorphogenesis*. G. S. Singhal, G. Renger, S. K. Sopory, K. D. Irrgang, and Govindjee, editors. Narosa Publishing House, New Delhi. 221–263.
- McMahon, B. H., J. D. Müller, C. A. Wraight, and G. U. Nienhaus. 1998. Electron transfer and protein dynamics in the photosynthetic reaction center. *Biophys. J.* 74:2567–2587.
- Moser, C. C., C. C. Page, R. Farid, and P. L. Dutton. 1995. Biological electron transfer. *J. Bioenerg. Biomembr.* 27:263–274.
- Naver, H., M. P. Scott, J. H. Golbeck, B. L. Møller, and H. V. Scheller. 1996. Reconstitution of barley photosystem I with modified PSI-C allows identification of domains interacting with PSI-D and PSI-A/B. *J. Biol. Chem.* 271:8996–9001.
- Parrett, K., T. Mehari, P. G. Warren, and J. H. Golbeck. 1989. Purification and properties of the intact P700 and Fx-containing photosystem I core protein. *Biochim. Biophys. Acta.* 973:324–332.
- Sakurai, H., K. Inoue, T. Fujii, and P. Mathis. 1991. Effects of selective destruction of iron-sulfur center B on electron transfer and charge recombination in Photosystem I. *Photosynth. Res.* 27:65–21.
- Sauer, K., P. Mathis, S. Acker, and J. A. Van Best. Electron acceptors associated with P-700 in Triton solubilized Photosystem I particles from spinach chloroplasts. *Biochim. Biophys. Acta.* 503:120–134.
- Scheller, H. V., H. Naver, and B. L. Møller. 1997. Molecular aspects of photosystem I. *Physiol. Plant.* 100:842–851.
- Schlodder, E., K. Falkenberg, M. Gergeleit, and K. Brettel. 1998. Temperature dependence of forward and reverse electron transfer from A<sub>1</sub><sup>-</sup>, the reduced secondary electron acceptor in photosystem I. *Biochemistry.* 37:9466–9476.
- Schubert, W. D., O. Klukas, N. Krauss, W. Saenger, P. Fromme, and H. T. Witt. 1997. Photosystem I of *Synechococcus elongatus* at 4 Å resolution: comprehensive structure analysis. *J. Mol. Biol.* 272:741–769.
- Shinkarev, V. P., and C. A. Wraight. 1993. Electron and proton transfer in the acceptor quinone complex of reaction centers of phototrophic bacteria. In *The Photosynthetic Reaction Center*, Vol. 1. J. Deisenhofer and J. Norris, editors. Academic Press, New York. 193–255.
- Vassiliev, I. R., Y. S. Jung, M. D. Mamedov, A. Yu. Semenov, and J. H. Golbeck. 1997. Near-IR absorbance changes and electrogenic reactions in the microsecond-to-second time domain in photosystem I. *Biophys. J.* 72:301–315.
- Vassiliev, I. R., Y. S. Jung, F. Yang, and J. H. Golbeck. 1998. PsaC subunit of photosystem I is oriented with iron-sulfur cluster F<sub>B</sub> as the immediate electron donor to ferredoxin and flavodoxin. *Biophys. J.* 74:2029–2035.
- Yu, J., L. B. Smart, Y. S. Jung, J. H. Golbeck, and L. McIntosh. 1995. Absence of PsaC allows assembly of photosystem I core but prevents binding of PsaD and PsaE in *Synechocystis* sp. PCC 6803. *Plant Mol. Biol.* 29:331–342.