

Table 7. No. of secondary cases caused by three *M. tuberculosis* lineages in five different patient populations in San Francisco (1991–2001)

Patient population (born in)	<i>M. tuberculosis</i> lineage	Secondary cases					
		<i>n</i>	(%)	Rate	Rate ratio	(95% C.I.)	<i>P</i> value
Total San Francisco	East-Asian	89	(14.7)	0.17	0.32	(0.25–0.40)	<0.0001
	Indo-Oceanic	124	(20.5)	0.3	0.71	(0.58–0.86)	<0.001
	Euro-American	381	(63.1)	0.68	3.0	(2.5–3.6)	<0.0001
	Total	604	(100)				
United States	East-Asian	43	(11.8)	0.08	0.25	(0.18–0.35)	<0.0001
	Indo-Oceanic	10	(2.8)	0.02	0.08	(0.04–0.14)	<0.0001
	Euro-American	302	(83.2)	0.54	9.5	(7.1–13.0)	<0.0001
	Total	363	(100)				
China	East-Asian	31	(67.4)	0.06	3.8	(2.0–7.5)	<0.0001
	Indo-Oceanic	8	(17.4)	0.02	0.56	(0.23–1.2)	0.13
	Euro-American	7	(15.2)	0.01	0.3	(0.11–0.68)	<0.01
	Total	46	(100)				
The Philippines	East-Asian	2	(2.6)	0.004	0.05	(0.006–0.18)	<0.0001
	Indo-Oceanic	67	(88.2)	0.16	19.9	(9.9–45.5)	<0.0001
	Euro-American	7	(9.2)	0.01	0.17	(0.07–0.37)	<0.0001
	Total	76	(100)				
Central America	East-Asian	2	(5.6)	0.004	0.11	(0.01–0.42)	<0.0001
	Indo-Oceanic	2	(5.6)	0.005	0.16	(0.02–0.61)	<0.01
	Euro-American	32	(88.9)	0.06	13.4	(4.7–52.1)	<0.0001
	Total	36	(100)				
Vietnam	East-Asian	3	(9.7)	0.006	0.2	(0.04–0.63)	<0.01
	Indo-Oceanic	26	(83.9)	0.06	13.9	(5.3–46.5)	<0.0001
	Euro-American	2	(6.5)	0.004	0.12	(0.01–0.46)	<0.001
	Total	31	(100)				

C.I., confidence interval.