

*Progress report*

## Bicarbonate ( $\text{HCO}_3$ ) delivery to the gastroduodenal mucosa by the blood: its importance for mucosal integrity

Recent results from our laboratory,<sup>1-3</sup> as well as data reported by others,<sup>4-9</sup> have shown that gastroduodenal blood flow functions not only provide  $\text{O}_2$ <sup>10</sup> and substrates, but also deliver  $\text{HCO}_3$  to the mucosa. This may be of particular relevance to the pathogenesis of stress ulceration. Sepsis and haemorrhagic shock are frequently associated with systemic acidosis and reduced arterial  $\text{HCO}_3$  concentration. The recent decline in the rate of severe bleeding from acute gastroduodenal lesions may in part be explained by the more aggressive approach to promptly correct any imbalance of the acid base status in these patients.<sup>11</sup>

It is clear that reduction of blood flow increases the susceptibility of the gastric and duodenal mucosa to the injurious actions of luminal acid.<sup>12,13</sup> In many experimental models gastric mucosal ulceration after luminal acid exposure alone can only be achieved when blood flow is artificially reduced.<sup>14,17</sup> Bile salt or aspirin injury is enhanced by concomitant reduction of blood flow.<sup>18-20</sup>

The gastric mucosa appears to be relatively resistant to changes in blood flow compared with the duodenum.<sup>15</sup> Thus a 60% reduction of baseline blood flow was necessary to produce damage in a model of hypotensive shock in the rat, whereas the duodenal mucosa ulcerated after relatively minor changes in blood flow. Conversely, increasing blood flow by intra-arterial infusion of isoproterenol decreased gastric mucosal damage caused by bile salts, haemorrhagic shock or aspirin in the dog.<sup>21,22</sup> The fact that sympathectomy attenuated the decrease in blood flow during haemorrhagic shock and also reduced gastric lesion formation is consistent with the protective role of blood  $\text{HCO}_3$ .<sup>23</sup> A direct effect on other protective mechanisms such as cellular  $\text{HCO}_3$  transport mechanisms may also be involved as this has been shown to be under sympathoadrenergic control in the isolated mucosa.<sup>24,25</sup> Similar arguments apply to the action of prostaglandins (PG) which effect both the vasculature as well as having direct effects on  $\text{HCO}_3$  transport by the mucosal cells. Whittle reported an increase in blood flow and associated protection against bile salt and indomethacin induced damage after iv administration of a variety of prostaglandin analogues. Direct infusion of  $\text{PGI}_2$  into the coeliac artery supplying an *in vivo* chambered gastric mucosal flap prevented taurocholate and macroscopic acid induced damage in indomethacin pretreated dogs, and this was associated with a large increase in mucosal blood flow.<sup>26</sup> A later report from the same laboratory, however, showed that surface cell damage was induced by taurocholate and acid alone emphasising the importance of microscopic evaluation in the assessment of damage.<sup>17,14</sup> We do not know whether increased blood flow protects against damage of the surface

epithelium because all the above studies only used macroscopic assessment of lesions.

Increased  $H^+$  back diffusion stimulates blood flow to the gastric and duodenal mucosa.<sup>13 27 28</sup> In the rabbit stomach this increase seemed to occur in apparently undamaged mucosa and was proposed to prevent tissue acidification.<sup>28</sup> Careful histological studies evaluating surface damage were, however, not done in these experiments. In the duodenum histologic damage to the villous tips was not prevented and a relative decline in blood flow was only seen when damage reached the base of the villi.<sup>13</sup> These findings suggest that at least in the duodenum an increase in mucosal blood flow is not protective for the superficial layer of the mucosa. Similar results were obtained by Cheung *et al*<sup>29</sup> who found that the increase in blood flow associated with increased  $H^+$  loss from the gastric lumen was correlated with the magnitude of gastric mucosal damage after aspirin, taurocholate, or ethanol. McGreevy<sup>30</sup> also found an increase in blood flow in areas of aspirin induced erosions compared with macroscopically undamaged mucosa.

The increase in blood flow after damage has occurred may be important in limiting the degree of mucosal acidosis by buffering and diluting back diffusing  $H^+$  and in facilitating rapid repair by creating an alkaline milieu underneath the layer of fibrin, mucus, and necrotic cells. Direct evidence for such a mechanism has recently been obtained by Kivilaakso<sup>8</sup> in the rat after taurocholate damage. Using subepithelial pH sensitive micro-electrodes, surface cell damage was associated with an intense alkalisation (after a transient drop in pH) that was blocked by haemorrhagic shock, suggesting outpouring of  $HCO_3^-$  rich fluid from subepithelial blood vessels in response to mucosal injury. Increase in  $HCO_3^-$  effusion may also be important for epithelial repair after damage of the amphibian isolated gastric mucosa with hypertonic NaCl and in the rabbit isolated duodenum after acid damage<sup>3 1-33</sup> as recovery depends on the presence of  $HCO_3^-$  in the serosal bathing solution.

Experiments using pH sensitive microelectrodes have shown that in the rabbit gastric mucosa reduction of blood flow by haemorrhagic shock or vasopressin was accompanied by a reduction of pH in the lamina propria that exceeded the fall in blood pH suggesting  $H^+$  back diffusion.<sup>6 28 34</sup> When submucosal pH dropped below 6.9 mucosal ulceration occurred. Ischaemia alone in the absence of luminal acid and associated fall in mucosal pH did not lead to macroscopic damage. These results indicated that mucosal acidosis may be more important in the development of mucosal lesions than the limitation in  $O_2$  supply. This hypothesis is further supported by the experimental findings that acid secreting stomachs were more resistant to injury by comparison with non-secreting preparations and had a lesser degree of intramural acidosis, presumably because of increased serosal  $HCO_3^-$  produced by the parietal cells.<sup>35</sup> Thus  $HCO_3^-$  is extruded across the basolateral membrane and readily available for buffering back-diffusing  $H^+$ . Microvascular architecture appropriate to accomplish the transport of  $HCO_3^-$  from the gastric glands to the surface cells has been shown.<sup>36</sup> Similar results showing that the secreting stomach is more resistant to injury have been obtained *in vitro*.<sup>37 38</sup>

Recent experiments in the frog gastric mucosa *in vitro* using pH-sensitive fluorescent dye have directly shown a profound alkalisation of the lamina propria upon onset of acid secretion.<sup>39</sup> Using the same technique to measure

pH actually within cells of rabbit gastric glands showed these cells to be sensitive to alteration of extracellular pH.<sup>40</sup> When the basolateral membrane was exposed to Ringers solutions of different pH, glands seemed to tightly regulate their intracellular pH over the range between extracellular pH 7.0 and pH 7.8. Outside this range the intracellular pH varied linearly with the pH outside. It is therefore to be expected that changes in pH in the lamina propria of the intact tissue below pH 7 cause intracellular acidosis with ensuing cell damage and death.

The combined effects of acidosis and low blood flow are certainly deleterious to the gastric mucosa. Prevention of acidosis can attenuate gastric mucosal injury. As early as 1948, Cummins and Grossman<sup>41</sup> showed that  $\text{HCO}_3$  infusions can prevent ulcer formation induced by continuous acid instillation into the stomach of dogs. Studies *in vitro* and *in vivo* have further confirmed the concept that nutrient  $\text{HCO}_3$  is essential for the gastric mucosa to withstand luminal acid.<sup>2 7 17 38 42 43</sup> This buffer species cannot be substituted by other buffers because phosphate, HEPES, MES or TES *in vitro*, and TRIS and phosphate *in vivo* were without effect.<sup>2 43</sup> In a study of acute gastric lesion formation in the rat, however, it was clearly shown that preventing the fall in gastric mucosal blood flow during hemorrhagic hypotension using intra-arterial  $\text{PGI}_2$  did not prevent gross mucosal ulceration, but correction of the accompanying acidosis by infusion of  $\text{HCO}_3$  inhibited the development of haemorrhagic lesions despite low blood flow.<sup>2</sup> These results indicated that it is the availability of  $\text{HCO}_3$  that is the critical factor in the development of lesions even in states of reduced blood flow rather than the limited  $\text{O}_2$  supply or flow *per se*.

There are no data available on the dependence of gastric  $\text{HCO}_3$  secretion in the lumen on blood flow, but experiments in our laboratory have clearly shown that changes in blood flow are directly related to changes in alkaline secretion from the duodenal mucosa.<sup>1</sup> When arterial  $\text{HCO}_3$  concentration was also considered and bicarbonate availability ( $[\text{HCO}_3]$  art x blood flow) was plotted against alkaline secretion, the fitted curve demonstrated a saturable process. Thus alkaline secretion was independent of  $\text{HCO}_3$  delivery to the mucosa above 4 mmol/cm<sup>2</sup>/min, but extremely sensitive to a reduction below this value (Fig. 1). Because alkaline secretion in this model is also an important determinant of acid induced injury and directly correlated to the extent of damage, it seems reasonable to conclude from these data that at least in the duodenum,  $\text{HCO}_3$  availability is of primary importance for the mucosa to withstand luminal acid.<sup>3</sup> Similarly, when  $\text{HCO}_3$  delivery to the rat gastric fundus during haemorrhagic shock and  $\text{PGI}_2$  and/or  $\text{HCO}_3$  infusions (data calculated from ref. 2) are plotted against the percentage of stomachs ulcerated a linear correlation is found (Fig. 2), suggesting that  $\text{HCO}_3$  delivery to the gastric mucosa is also a critical factor in the development of lesions. The mechanism of action of nutrient bicarbonate in the stomach is less clear, however, than in the duodenum where luminal buffering by alkaline secretion is the single most important factor in the defence against luminal acid.<sup>3 44</sup>

$\text{HCO}_3$  secretion from the gastric mucosa is relatively small, amounting to <5% of maximal acid secretion. Under basal conditions, acidification of the lumen down to pH 2 causes a two to three fold increase of luminal  $\text{HCO}_3$  appearance which can completely account for the  $\text{H}^+$  disappearance measured under these conditions.<sup>44-47</sup> The concentration of acid used in

these experiments<sup>47</sup> was rather small, however (10 mmol/l HCl), and the rate of H<sup>+</sup> loss in this model (dog Heidenhain pouch) is linearly dependent on luminal H<sup>+</sup> concentration up to 150 mmol/l.




Fig. 1 Correlation of alkaline secretion and  $\text{HCO}_3^-$  delivery in rabbit duodenum. Mucosal blood flow determined with radioactive microspheres. Alteration of blood flow and arterial  $\text{HCO}_3^-$  concentration with iv infusion of vasopressin,  $\text{HCO}_3^-$  and  $\text{NH}_4\text{Cl}$ .




Fig. 2 Dependence of gastric mucosal damage (expressed as per cent of stomachs ulcerated) on  $\text{HCO}_3^-$  delivery by blood flow (mucosal blood flow determined with radioactive microspheres) in stomachs of rats subjected to haemorrhagic shock with and without infusion of  $\text{PGI}_2$  and  $\text{HCO}_3^-$  (points represent means of each experimental group, ref 2).

The essential remaining question is where and when  $\text{HCO}_3^-$  acts to prevent damage and facilitate repair. The pre-epithelial 'mucus bicarbonate' barrier may not be of primary importance in the stomach because the gradient measured with microelectrodes readily dissipated at luminal pH 1.5 or less,

values frequently encountered in the *in vivo* situation.<sup>40-49-52</sup> Also luminal stirring and shear forces are likely to reduce surface unstirred layers even further in the intact stomach.<sup>53-54</sup> The actual thickness of the adherent mucus gel varies enormously and may be as thin as 5  $\mu\text{m}$ .<sup>55</sup>

There are several recent data indicating that the permeability of the apical plasma membrane to  $\text{H}^+$  is low and this, together with efficient mechanisms to regulate intracellular pH, would mean that complete neutralisation of  $\text{H}^+$  at the cell surface is unnecessary.<sup>56-57</sup> Nutrient  $\text{HCO}_3$  is intimately involved in these pH regulatory mechanisms.<sup>40-56-58</sup>

Basal  $\text{HCO}_3$  secretion in the stomach is low and stimulation by luminal acid, sham feeding or prostaglandin only increases measurable alkali output by a small amount (two to three fold).<sup>59-62</sup> Increases in secretory rate in the duodenum are much greater.<sup>63-65</sup> In the stomach damage to the surface epithelium is associated with a comparatively large flux of alkali from the interstitium into the lumen. Such an increase in passive  $\text{HCO}_3$  effusion has been shown to occur after damage in variety of models.<sup>9-33-59-66-68</sup> Thus  $\text{HCO}_3$  in conjunction with a thick layer of mucus and necrotic debris probably acts to limit damage by facilitating rapid repair of minor disruption of the epithelial cell layer that occurs during everyday life.<sup>32-68-69-70</sup>

In summary,  $\text{HCO}_3$  delivery to the gastroduodenal mucosa is necessary to maintain its structural and morphological integrity. In the stomach the main function of  $\text{HCO}_3$  probably lies in its role in intracellular pH-regulation and in passive effusion after even minor injury. In the duodenum transepithelial secretion of  $\text{HCO}_3$  and intraluminal buffering is the predominant defence mechanism against luminal acid.

M STARLINGER AND R SCHIESSEL

*Ist Surgical University Clinic of Vienna,  
Vienna, Austria.*

Address for correspondence: M Starlinger, Chirurgische Klinik u Poliklinik, Calwerstr 7, D-7400 Tübingen, West Germany.

Received for publication 23 October 1987.

## References

- 1 Schiessel R, Starlinger M, Kovats E, Appel W, Feil W, Simon A. Alkaline secretion of rabbit duodenum *in vivo*: its dependence on acid base balance and mucosal blood flow. In: Allen A, Flemström G, Garner A, *et al*, eds. *Mechanisms of mucosal protection in the upper gastrointestinal tract*. New York: Raven Press, 1984: 267-71.
- 2 Starlinger M, Jakesz R, Matthews JB, Yoon Ch, Schiessel R. The relative importance of  $\text{HCO}_3^-$  and blood flow in the protection of rat gastric mucosa during shock. *Gastroenterology* 1981; **81**: 732-5.
- 3 Wenzl E, Feil W, Starlinger M, Schiessel R. Alkaline secretion. A protective mechanism against acid injury in rabbit duodenum. *Gastroenterology* 1987; **92**: 709-15.
- 4 Bushell M, O'Brien P. Acid-base imbalance and ulceration in the cold restrained rat. *Surgery* 1982; **91**: 318-21.
- 5 Cheung LY, Porterfield G. Protection of gastric mucosa against acute ulceration by intravenous infusion of sodium bicarbonate. *Am J Surg* 1979; **137**: 106-10.
- 6 Kivilaakso E, Fromm O, Silen W. Relationship between ulceration and intramural pH of gastric mucosa during hemorrhagic shock. *Surgery* 1978; **84**: 70-8.
- 7 Kivilaakso E. High plasma  $\text{HCO}_3^-$  protects gastric mucosa against acute ulceration in the rat. *Gastroenterology* 1981; **81**: 921-7.
- 8 Kiviluoto T, Voipio J, Kivilaakso E. Is  $\text{H}^+$  back diffusion following disruption of the gastric mucosal barrier in fact alkali ( $\text{HCO}_3^-$ ) efflux? *Gastroenterology* 1987; **92**: 1470.
- 9 Takeuchi K, Okabe S. Role of luminal alkalization in repair process of ethanol-induced mucosal damage in rat stomach. *Dig Dis Sci* 1983; **28**: 993-100.

- 10 Perry MA, Haedicke GJ, Bulkeley GB, Kvietyts PR, Granger ND. Relationship between acid secretion and blood flow in the canine stomach: role of oxygen consumption. *Gastroenterology* 1983; **85**: 529–34.
- 11 Groll A, Simon JB, Wigle RD, Taguchi K, Todd RJ, Depew WT. Cimetidine prophylaxis for gastrointestinal bleeding in an intensive care unit. *Gut* 1986; **27**: 135–40.
- 12 Collan Y, Kivilaakso B, Kalima TV, Lempinen M. Ultrastructural changes in the gastric mucosa following hemorrhagic shock in pigs. *Circulatory Shock* 1977; **4**: 13–25.
- 13 Starlinger M, Matthews JB, Yoon C, Wenzl E, Feil W, Schiessel R. The effect of acid perfusion on mucosal blood flow and intramural pH of rabbit duodenum. *Surgery* 1987; **101**: 433–8.
- 14 Itoh M, Paulsen G, Guth PH. Hemorrhagic shock and acid gastric injury in the rat. Comparison of gross and histologic findings. *Gastroenterology* 1986; **90**: 1103–10.
- 15 Leung FW, Itoh M, Hirabayashi K, Guth PH. Role of blood flow in gastric and duodenal mucosal injury in the rat. *Gastroenterology* 1985; **88**: 281–9.
- 16 Mersereau WA, Hinchey EJ. Interactions of gastric blood flow, barrier breaker and hydrogen ion back diffusion during ulcer formation in the rat. *Surgery* 1978; **83**: 248–52.
- 17 Carter KJ, Farley PC, Ritchie WP. Effect of topical bile acids on gastric surface epithelial cells. *Surgery* 1984; **96**: 196–203.
- 18 Ritchie WP, Shearburn EW. Influence of isoproterenol and cholestyramine on acute gastric mucosal ulcerogenesis. *Gastroenterology* 1977; **73**: 62–5.
- 19 Whittle BJR. Mechanisms underlying gastric mucosal damage induced by indomethacin and bile salts and the actions of prostaglandin *Br J Pharmacol* 1977; **60**: 455–60.
- 20 Whittle BJR, Kauffman GL, Moncada S. Vasoconstriction with thromboxane  $A_2$  induces ulcerations of the gastric mucosa. *Nature* 1981; **292**: 472–4.
- 21 Ritchie W. Acute gastric mucosal damage induced by bile salts, acid and ischemia. *Gastroenterology* 1975; **68**: 699–707.
- 22 McGreevy JM, Moody FG. Protection of gastric mucosa against aspirin-induced erosions by enhanced blood flow. *Surg Forum* 1971; **28**: 357–9.
- 23 Levine BA, Gaskill HV, Sirinek KR. Gastric mucosal cytoprotection by splanchnicectomy is based on protection of gastric mucosal blood flow. *J Trauma* 1983; **23**: 278–84.
- 24 Fändrics L. Sympatho-adrenergic inhibition of vagally induced gastric motility and gastroduodenal  $HCO_3^-$  secretion in the cat. *Acta Physiol Scand* 1986; **128**: 555–62.
- 25 Flemström G. Effect of catecholamines,  $Ca^{2+}$  and gastrin on gastric  $HCO_3^-$  secretion. *Acta Physiol Scand Suppl Gastric Ion Transport* 1978: 81–90.
- 26 Cloud WG, Ritchie WP. Evidence for cytoprotection by endogenous prostaglandins in gastric mucosa treated with bile acid. *Surg Forum* 1982; **33**: 150–2.
- 27 Bruggeman TM, Wood JG, Davenport HW. Local control of blood flow in the dog's stomach: vasodilatation caused by acid back-diffusion following topical application of salicylic acid. *Gastroenterology* 1979; **77**: 736–44.
- 28 Starlinger M, Schiessel R, Hung CK, Silen W.  $H^+$  back diffusion stimulating gastric mucosal blood flow in the rabbit fundus. *Surgery* 1981; **89**: 232–6.
- 29 Cheung LY, Moody FG, Reese RS. Effect of aspirin, bile salt and ethanol on canine gastric mucosal blood flow. *Surgery* 1975; **7**: 786–92.
- 30 McGreevy JM, Moody FG. Focal microcirculatory changes during the production of aspirin-induced gastric mucosal erosions. *Surgery* 1983; **89**: 337–41.
- 31 Svanes K, Takeuchi K, Ito S, Silen W. Effect of luminal pH and nutrient bicarbonate concentration on restitution after gastric surface cell injury. *Surgery* 1983; **94**: 494–500.
- 32 Starlinger M, Feil W, Klimesch S, Karner P, Schiessel R. The role of an alkaline micro-environment for mucosal repair in rabbit duodenum in vitro. *Gastroenterology* 1987; **92**: 1652.
- 33 Vattay P, Feil W, Starlinger M, Schiessel R. Acid-stimulated alkaline secretion of rabbit duodenum. The role of passive diffusion and mucosal damage. *Gastroenterology* 1986; **90**: 1678.
- 34 Kivilaakso E, Barzilai A, Schiessel R, Fromm O, Silen W. Experimental ulceration of rabbit antral mucosa. *Gastroenterology* 1981; **80**: 77–83.
- 35 Kivilaakso E, Fromm O, Silen W. Effect of the acid secretory state on intramural pH of rabbit gastric mucosa. *Gastroenterology* 1978; **75**: 641–8.
- 36 Gannon B, Browning J, O'Brien P, Rogers P. Mucosal microvascular architecture of the fundus and body of human stomach. *Gastroenterology* 1984; **86**: 866–75.
- 37 Barzilai A, Schiessel R, Kivilaakso E, et al. Effect of 16-16-dimethyl-prostaglandin  $E_2$  on ulceration of isolated amphibian gastric mucosa. *Gastroenterology* 1980; **78**: 1508–12.

- 38 Kivilaakso E, Barzilai A, Schiessel R, Crass R, Silen W. Ulceration of isolated amphibian gastric mucosa. *Gastroenterology* 1979; **77**: 32–7.
- 39 Saario I, Carter K, Silen W. Intracellular pH (pHi) in stimulated vs. inhibited parietal cells of frog gastric mucosa. *Gastroenterology* 1987; **92**: 1607.
- 40 Starlinger M, Paradiso A, Machen T. Steady state regulation of intracellular pH in isolated rabbit gastric glands. Roles for Na/H and Cl/OH ( $\text{HCO}_3$ ) exchange. *Gastroenterology* 1987; **92**: 957–65.
- 41 Cummins GM, Grossman MI, Ivy AC. An experimental study of the acid factor in ulceration of the gastrointestinal tract in dogs. *Gastroenterology* 1948; **10**: 714–26.
- 42 Rowe PH, Lange R, Marrone G, Matthews JB, Kasdon E, Silen W. In vitro protection of amphibian gastric mucosa by nutrient  $\text{HCO}_3^-$  against aspirin injury. *Gastroenterology* 1985; **89**: 769–78.
- 43 Schiessel R, Merhav A, Matthews JB, Fleischer L, Barzilai A, Silen W. Role of nutrient  $\text{HCO}_3^-$  in protection of amphibian gastric mucosa. *Am J Physiol* 1980; **239**: G536–42.
- 44 Flemström G, Garner A. Gastroduodenal  $\text{HCO}_3^-$  transport: characteristics and proposed role in acidity regulation and mucosal protection. *Am J Physiol* 1982; **242**: G183–93.
- 45 Smeaton LA, Hirst BH, Allen A. Gastric bicarbonate output in the cat. In: Case RM, Garner A, Turnberg LA, *et al*, eds. *Electrolyte and water transport across gastrointestinal epithelial*. New York: Raven Press, 1982: 257–62.
- 46 Takeuchi K, Merhav A, Silen W. Mechanism of luminal alkalinization by bullfrog fundic mucosa. *Am J Physiol* 1982; **243**: G377–88.
- 47 Garner A, Hurst BC, Heylings JR, Flemstrom G. Role of gastroduodenal  $\text{HCO}_3^-$  transport in acid disposal and mucosal protection. In: Case RM, Garner A, Turnberg LA, *et al*, eds. *Electrolyte and water transport across gastrointestinal epithelia*. New York: Raven Press, 1982: 239–52.
- 48 Chung R, Field M, Silen W. Effects of methylprednisolone on hydrogen ion absorption in the canine stomach. *J Chir Invest* 1978; **62**: 267–70.
- 49 Allen A, Hutton OA, Leonard AJ, Peakson JP, Sellers LA. The role of mucus in the protection of the gastroduodenal mucosa. *Scand J Gastroenterol* 1986; **21**: suppl 125: 71–7.
- 50 Ross IN, Bahari HMM, Turnberg LA. The pH-gradient across mucus adherent to rat fundus mucosal in vivo and the effects of potential damaging agents. *Gastroenterology* 1981; **81**: 713–8.
- 51 Ross IN, Turnberg LA. Studies of the ‘mucus-bicarbonate’ barrier on rat fundic mucosa: the effects of luminal pH and a stable prostaglandin analogue. *Gut* 1983; **24**: 1030–3.
- 52 Takeuchi K, Magee D, Critchlow J, Matthews JB, Silen W. Studies of the pH gradient and thickness of frog gastric mucus gel. *Gastroenterology* 1983; **84**: 331–40.
- 53 Allen A, Hutton DA, McQueen S, Garner A. Dimensions of gastroduodenal surface pH gradient exceed those of adherent mucus gel layers. *Gastroenterology* 1983; **85**: 463–6.
- 54 McQueen S, Hutton D, Allen A, Garner A. Gastric and duodenal surface mucus gel thickness in rat: effects of prostaglandins and damaging agents. *Am J Physiol* 1983; **245**: G388–93.
- 55 McQueen A, Allen A, Garner A. Measurement of gastric and duodenal mucus gel thickness. In: Allen A, Flemstrom G, Garner A, *et al*, eds. *Mechanisms of mucosal protection in the upper gastrointestinal tract*. New York: Raven Press, 1984: 215–21.
- 56 Kivilaakso E, Voipio J. Control of intracellular pH (pHi) in isolated necturus antral mucosa exposed to luminal acid. *Gastroenterology* 1987; **92**: 1469.
- 57 Sanders MJ, Avalon A, Roll M, Soll AH. The apical surface of canine chief cells resists  $\text{H}^+$ -back diffusion. *Nature* 1985; **313**: 52–4.
- 58 Kivilaakso E. Contribution of ambient  $\text{HCO}_3^-$  to mucosal protection and intracellular pH in isolated amphibian gastric mucosa. *Gastroenterology* 1983; **85**: 1284–9.
- 59 Konturek StJ, Bilski J, Tasler J, Laskiewicz J. Gastroduodenal alkaline response to acid and taurocholate in conscious dogs. *Am J Physiol* 1984; **247**: G149–54.
- 60 Konturek StJ, Bilski J, Tasler J, Laskiewicz J. Gut hormones in stimulation of gastroduodenal alkaline secretion in conscious dogs. *Am J Physiol* 1985; **248**: G677–91.
- 61 Konturek StJ, Thor P. Relation between duodenal alkaline secretion and motility in fasted and sham-fed dogs. *Am J Physiol* 1986; **251**: G591–6.
- 62 Kauffman GL, Reeve JJ, Grossman MI. Gastric bicarbonate secretion: effect of topical and intravenous 16-16-dimethyl prostaglandin  $\text{E}_2$ . *Am J Physiol* 1980; **239**: G44–8.
- 63 Heylings JR, Garner A, Flemström G. Regulation of gastroduodenal  $\text{HCO}_3^-$  transport by luminal acid in the frog in vitro. *Am J Physiol* 1984; **246**: G235–42.
- 64 Isenberg JI, Hogan DL, Koss AM, Selling JA. Human duodenal mucosal bicarbonate

- secretion and stimulation by hydrochloric acid and a synthetic prostaglandin E<sub>1</sub> analog. *Gastroenterology* 1986; **91**: 370–8.
- 65 Isenberg JI, Smedfors B, Johansson C. Effect of graded doses of intraluminal H<sup>+</sup>, prostaglandin E<sub>2</sub> and inhibition of endogenous prostaglandin synthesis on proximal duodenal bicarbonate secretion in unanesthetized rats. *Gastroenterology* 1985; **88**: 303–7.
- 66 Smeaton LA, Hirst BH. Gastroduodenal ion-outputs: Prostaglandins and hyperosmolar solutions stimulate via different mechanisms. In: Allen A, Flemström G, Garner A, *et al*, eds. *Mechanisms of mucosal protection in the upper gastrointestinal tract*. New York: Raven Press, 1984: 107–12.
- 67 Svanes K, Ito S, Takeuchi K, Silen W. Restitution of the surface epithelium of the in vitro frog gastric mucosa after damage with hyperosmolar sodium chloride. *Gastroenterology* 1982; **82**: 1409–26.
- 68 Dayton MT, Kauffman GK, Schlegel JF, Code ChF, Steinbach JH. Gastric bicarbonate appearance with ethanol ingestion. Mechanism and significance. *Dig Dis Sci* 1983; **28**: 449–5.
- 69 Morris GP, Harding PL, Wallace JP. A functional model for extracellular gastric mucus in the rat. *Virch Arch [Cell Pathol]* 1984; **46**: 239–51.
- 70 Wallace JL, Whittle BJR. Role of mucus in the repair of gastric epithelial damage in the rat. Inhibition by epithelial recovery by mucolytic agents. *Gastroenterology* 1986; **91**: 603–11.