

Appendix 1: Baseline characteristics of patients in the ASSENT-3 PLUS and ASSENT-3 trials, by treatment group

Characteristic	ASSENT-3 PLUS						ASSENT-3					
	Enoxaparin, n = 818			UF heparin, n = 821			Enoxaparin, n = 2040			UF heparin, n = 2038		
	No.	%*	n	No.	%*	n	No.	%*	n	No.	%*	n
Age, yr, mean (SD)	62 (13)			62 (13)			61 (12)			61 (13)		
Patients aged > 75 yr	149	18	817	132	16	820	263	13	2040	271	13	2038
Women	194	24	818	184	22	821	463	23	2040	478	23	2038
Women aged > 75 yr	73	38	193	56	30	184	110	24	463	129	27	478
Hypertension	289	35	815	298	36	820	826	41	2037	824	41	2031
Diabetes	115	14	816	128	16	820	381	19	2039	363	18	2031
Previous myocardial infarction	120	15	816	117	14	820	290	14	2040	293	14	2035
Prior CABG	21	2.6	814	13	1.6	819	73	3.6	2038	53	2.6	2029
Prior PCI	60	7.4	806	52	6.4	816	126	6.2	2019	128	6.4	2012
Current smoker	336	42	801	365	45	809	884	44	1992	930	47	1988
Infarct location												
Anterior	341	42	813	350	43	820	802	39	2040	772	38	2037
Inferior	439	54	813	444	54	820	1144	56	2040	1164	57	2037
Other	33	4.0	813	26	3.2	820	94	4.6	2040	101	5.0	2037
Heart rate, beats/min, mean (SD)	74 (19)			74 (19)			75 (17)			74 (17)		
Systolic BP, mm Hg, mean (SD)	134 (25)			132 (25)			134 (22)			133 (23)		
Killip class I†	593	92	642	603	92	654	1823	89	2039	1786	88	2033

Note: ASSENT = Assessment of the Safety and Efficacy of a New Thrombolytic Agent, UF = unfractionated, SD = standard deviation, CABG = coronary artery bypass graft, PCI = percutaneous coronary intervention, BP = blood pressure.

*Unless indicated otherwise.

†Killip class was unrecorded in the records of 21% of ASSENT-3 PLUS patients.