

Reduced Variation on the Chicken Z Chromosome

Hannah Sundström, Matthew T. Webster and Hans Ellegren¹

Department of Evolutionary Biology, Evolutionary Biology Centre, Uppsala University, SE-752 36 Uppsala, Sweden

Manuscript received December 29, 2003

Accepted for publication January 27, 2004

ABSTRACT

Understanding the population genetic factors that shape genome variability is pivotal to the design and interpretation of studies using large-scale polymorphism data. We analyzed patterns of polymorphism and divergence at Z-linked and autosomal loci in the domestic chicken (*Gallus gallus*) to study the influence of mutation, effective population size, selection, and demography on levels of genetic diversity. A total of 14 autosomal introns (8316 bp) and 13 Z-linked introns (6856 bp) were sequenced in 50 chicken chromosomes from 10 highly divergent breeds. Genetic variation was significantly lower at Z-linked than at autosomal loci, with one segregating site every 39 bp at autosomal loci ($\theta_w = 5.8 \pm 0.8 \times 10^{-3}$) and one every 156 bp on the Z chromosome ($\theta_w = 1.4 \pm 0.4 \times 10^{-3}$). This difference may in part be due to a low male effective population size arising from skewed reproductive success among males, evident both in the wild ancestor—the red jungle fowl—and in poultry breeding. However, this effect cannot entirely explain the observed three- to fourfold reduction in Z chromosome diversity. Selection, in particular selective sweeps, may therefore have had an impact on reducing variation on the Z chromosome, a hypothesis supported by the observation of heterogeneity in diversity levels among loci on the Z chromosome and the lower recombination rate on Z than on autosomes. Selection on sex-linked genes may be particularly important in organisms with female heterogamety since the heritability of sex-linked sexually antagonistic alleles advantageous to males is improved when fathers pass a Z chromosome to their sons.

SINGLE-NUCLEOTIDE polymorphisms (SNPs) constitute a primary source of the variability that underlies differences in the genetic make-up of individuals within species. Currently, the application of SNPs to genomics and population genetics is rapidly expanding and the increasing numbers of species in which an extensive number of SNPs have been identified include humans (INTERNATIONAL SNP MAP WORKING GROUP 2001), mouse (WADE *et al.* 2002), *Arabidopsis thaliana* (CHO *et al.* 1999), *Caenorhabditis elegans* (KOCH *et al.* 2000), maize (TENAILLON *et al.* 2001), and soybean (ZHU *et al.* 2003). Population genetics theory stipulates that the degree and character of genetic variability are governed by factors such as mutation, selection, and effective population size. Understanding the role of these factors in shaping patterns of polymorphism within genomes is key to the interpretation of SNP data in, for example, association studies and for the quantification of linkage disequilibrium (LD) through analysis of haplotype diversity. It is also required for tracing the evolutionary history of genomes and species, on the basis of polymorphism data.

Within-genome variation in SNP frequency is likely to occur at several different scales. To name but one reason, hot-spot sites for mutation as well as mutation rate variation at larger scales imply heterogeneity in the distribution of SNPs (EBERSBERGER *et al.* 2002; INTERNATIONAL HUMAN GENOME SEQUENCING CONSORTIUM 2002; SMITH *et al.* 2002; HARDISON *et al.* 2003). Empirical data from the human (MULLIKIN *et al.* 2000; INTERNATIONAL SNP MAP WORKING GROUP 2001; REICH *et al.* 2002) and mouse (WADE *et al.* 2002) genomes provide evidence of significant variation in the density of SNPs along chromosomes. There is also the potential of variation in SNP frequency *between* chromosomes and this applies in particular to differences between autosomes and sex chromosomes (BEGUN and WHITLEY 2000; INTERNATIONAL SNP MAP WORKING GROUP 2001). These chromosome classes differ with respect to several of the factors thought to affect nucleotide diversity and contrasting the levels of diversity in the different classes may thus be informative for addressing the role of population genetic factors in shaping genomic variability.

Preliminary surveys of the incidence of SNPs in the genome of the domestic chicken have indicated genetic variability to be at relatively high levels. No detailed estimates of nucleotide diversity are available but there are reports of one SNP every 60–500 bp (SCHMID *et al.* 2000, 2001; PARSANEJAD *et al.* 2002; SMITH *et al.* 2002; LIU *et al.* 2003), the broad range of estimated frequencies probably reflecting that only a limited number of genes have been analyzed, that coding as well as untrans-

Sequence data from this article have been deposited with the EMBL/GenBank Data Libraries under accession nos. AF165971, AF526014, AY139845–7, AY139851, AY139861, AY142943–4, AY144673–6, AY144678–9, AY144681–2, AY189760, AY189776, AY194143, AY194147, AY298982, AY298987, AY298992, AY380786, and AY380788–9.

¹Corresponding author: Department of Evolutionary Biology, Evolutionary Biology Centre, Uppsala University, Norbyvägen 18D, SE-752 36 Uppsala, Sweden. E-mail: hans.ellegren@ebc.uu.se

lated and intron regions have been surveyed, and that a varying number of breeds/lines have been sampled. It thus appears that despite strong artificial selection during domestication, average genome variability is higher in chicken than in, for instance, humans (INTERNATIONAL SNP MAP WORKING GROUP 2001). In contrast to mammals, birds have female heterogamety (males ZZ; females ZW) and the avian sex chromosomes have evolved independently from mammalian X and Y (FRIDOLFSSON *et al.* 1998; NANDA *et al.* 2002); *i.e.*, Z and X are not syntenic. Moreover, due to female heterogamety the population genetics of avian and mammalian sex chromosomes differ in several respects. In this study we survey genetic variability in noncoding regions of the chicken genome, including autosomes as well as the Z chromosome, using a panel of highly divergent breeds. We find nucleotide diversity to be strikingly higher in autosomes than in the Z chromosome, an observation that suggests low male effective population size and a role of selection in reducing levels of genetic variability on the Z chromosome.

MATERIALS AND METHODS

DNA samples: We used genomic DNA from 25 male chickens (*Gallus gallus*), giving 50 chromosomes of autosomal as well as Z sequences. The sample included two to three individuals from each of 10 diverse breeds, which was a subset of the 52 breeds used by HILLEL *et al.* (2003). This subset of breeds has been identified as constituting a broad representation of the total gene pool of domestic chicken (ROSENBERG *et al.* 2001). Two individuals were included from broiler sire line B, brown-egg-layer line D, broiler dam line D, Icelandic landrace, and captive red jungle fowl *G. g. gallus*. Three individuals were analyzed from Fayoumi, Marans, Transylvanian naked neck, Green-legged partridge, and White-egg-layer line A. For all introns analyzed we also determined the orthologous sequence in turkey, to be able to contrast levels of intraspecific diversity and interspecific divergence. Chicken and turkey have an estimated divergence time of 28 million years ago (DIMCHEFF *et al.* 2002).

PCR: We amplified 27 introns from 13 autosomal and 9 Z-linked genes (Table 1). While most introns were >400 bp, they were analyzed in overlapping ~250-bp amplicons. PCR was performed in 25- μ l reactions containing 20 ng DNA, 1 \times PCR Gold buffer (Applied Biosystems, Foster City, CA), 0.2 μ M of each primer, 2.0–2.5 mM MgCl₂, 2.0 μ M dNTPs, and 1 unit AmpliTaq Gold (Applied Biosystems). Amplification was performed using an initial denaturation of 95° for 5 min, followed by 33–40 cycles at 94° for 30 sec, a touchdown annealing temperature profile for 30 sec, and extension at 72° for 45 sec. Primer sequences, annealing temperatures, and MgCl₂ concentrations for each marker are available as supplementary information at <http://www.genetics.org/supplemental/>.

Sequencing and SNP detection: The fragments were purified using ExoSAP-IT (Amersham Biosciences, Arlington Heights, IL) and direct sequenced using original PCR primers and the DYEnamic cycle sequencing kit (Amersham Biosciences) and analyzed on a MegaBACE 1000 (Amersham Biosciences) instrument. Sequences from both directions were aligned and edited in the program AutoAssembler (Applied Biosystems), which also was used for defining heterozygous

positions. Singletons were confirmed by independent sequencing reactions using new amplification products. In cases where a fragment was found to be heterozygous for an insertion or deletion polymorphism, the fragment was resequenced from both directions using a new amplification product.

Complete diploid intron sequences were assembled from the overlapping fragments using Sequencher (Gene Codes, Ann Arbor, MI). Since phase was unknown, each sequence was then separated into two “pseudo-haplotypes” on the basis of the ambiguity codes produced by Sequencher using a Perl script. All chicken and turkey sequences at each locus were then aligned using a ClustalW algorithm (THOMPSON *et al.* 1994) in Sequence Navigator (Applied Biosystems) and improved by manual adjustment. A number of additional Perl programs were used during the analysis to extract polymorphism information from the sequences and to format them for use in data analysis programs. There were two segregating sites with three alleles, which indicates a frequency of multiallelic sites similar to that found in the human genome (PATIL *et al.* 2001).

Statistical analysis: Haplotypes were inferred at each locus, using the expectation-maximization (EM) algorithm implemented in Arlequin (SCHNEIDER *et al.* 2000); the inferred haplotypes are provided as supplementary information at <http://www.genetics.org/supplemental/>. We then used DnaSP (ROZAS and ROZAS 1999) to analyze patterns of polymorphism and divergence and perform tests of neutrality on the basis of the allele frequency spectrum. These included Tajima's *D* (TAJIMA 1989a) and Fu and Li's *D* and *F* (Fu and Li 1993) test statistics. These tests measure whether the observed frequencies of segregating mutations are compatible with the frequencies expected under a standard neutral model. Positive selection or the presence of weakly deleterious mutations (as well as population growth) tends to give an excess of low-frequency variants, resulting in negative test values. Balancing selection (or population contraction) may cause an excess of intermediate-frequency variants and positive test values. Multilocus Hudson-Kreitman-Aguadé (HKA) tests (HUDSON *et al.* 1987) were performed using the HKA program written by Jody Hey and available at <http://lifesci.rutgers.edu/~hey/lab/index.html>. This program allows tests assuming sex ratios different from 1:1. The basic principle of the HKA test is to test whether the levels of intraspecific diversity and interspecific divergence are positively correlated in two or more genomic regions, as predicted by the neutral mutation hypothesis. If they are not, selection may be interpreted as having affected polymorphism levels in at least one of the regions analyzed. AMOVA analysis of population subdivision was performed in Arlequin. Confidence intervals for the ratio of average levels of variation on autosomes and Z were determined by nonparametric bootstrapping: To account for potential mutation rate variation between loci, segregating sites from each chromosomal class were resampled with replacement first by site and then by locus, using 10,000 replicates.

RESULTS

We amplified and sequenced 27 different introns, from 22 different genes spread over the genome, in 25 male chickens from 10 diverse breeds to get a picture of the patterns of genetic variability in the chicken genome (Table 1). With this experimental design we sought to account for regional variation in the mutation rate and for local effects of selection. In mammals, regions of local similarity in mutation rate have been observed at

TABLE 1
Summary statistics of polymorphism screening in chicken introns

Locus	Intron	GenBank acc. no.		<i>L</i>	Average divergence (bp)	Polymorphism					Neutrality tests		
		<i>G. gallus</i>	<i>M. gallopavo</i>			<i>S</i>	π	θ_w	<i>H</i>	R_M	D_T	D_{FL}	F_{FL}
Autosomes													
<i>ACGR</i>	F	K02904	AY144678	702	0.0641	13	0.0034	0.0041	20	2	-0.51	-0.60	-0.68
<i>CEPU</i>	1	AJ225897	AY142943	611	0.0648	17	0.0081	0.0062	30	6	0.95	1.17	1.15
<i>CLTC</i>	5	AJ429073	AY142944	670	0.0834	8	0.0022	0.0027	7	2	-0.45	0.59	0.29
<i>DEF1</i>	3	D76434	AY144673	641	0.0911	17	0.0080	0.0059	23	6	1.10	1.28	1.46
<i>GHI</i>	2	AJ289775	AY144674	442	0.0881	11	0.0055	0.0056	18	4	-0.27	-0.77	-0.71
<i>LHCGR</i>	7	D10484	AY144675	500	0.0935	15	0.0085	0.0067	26	5	0.61	1.17	1.09
<i>L37a</i>	3	D14167	AY298987	844	0.0880	28	0.0086	0.0074	28	11	0.55	0.94	1.03
<i>MLC</i>	6	X00460	AY144676	469	0.1152	9	0.0051	0.0043	16	3	0.53	1.40	1.32
<i>RHO</i>	1	D00702	AY144679	797	0.1337	10	0.0033	0.0028	19	4	0.49	1.46	1.35
<i>RPL5</i>	4	D10737	AY380788	598	0.0768	9	0.0048	0.0034	17	4	1.19	0.70	1.03
<i>RPL7a</i>	3	D14522	AY380789	351	0.1233	13	0.0100	0.0083	18	6	0.62	1.59*	1.50
<i>RPL7a</i>	4	D14522	AY298982	337	0.0618	7	0.0041	0.0046	8	2	-0.31	1.28	0.90
<i>RPL30</i>	3	D14521	AY144681	748	0.1025	37	0.0130	0.0110	36	14	0.57	0.33	0.59
<i>VIM</i>	E	M15852	AY144682	606	0.1022	20	0.0061	0.0074	21	5	-0.54	0.21	-0.06
Z chromosome													
<i>AD012Z</i>	1	AY189761	AY189760	1230	0.0704	3	0.0008	0.0005	5	1	1.00	0.89	1.08
<i>AD012Z</i>	2	AY189777	AY189776	406	0.0837	0	0.0000	0.0000	1	—	NA	NA	NA
<i>ALDOB</i>	1	AY380785	AY380786	280	0.0875	2	0.0030	0.0016	3	0	1.56	0.74	1.15
<i>ALDOB</i>	5	AY139839	AY139845	473	0.0931	7	0.0029	0.0033	7	1	-0.34	0.46	0.23
<i>ALDOB</i>	6	AY139840	AY139846	143	0.0965	2	0.0059	0.0031	3	0	1.56	0.74	1.15
<i>ALDOB</i>	7	AY139841	AY139847	415	0.0817	4	0.0047	0.0022	8	3	2.66***	1.01	1.79*
<i>ATP5A1Z</i>	3	AF165969	AF165971	833	0.1121	12	0.0041	0.0032	21	5	0.823	0.97	1.10
<i>BRM</i>	15	AY298991	AY298992	280	0.0930	3	0.0042	0.0024	7	2	1.56	0.89	1.28
<i>CHD1Z</i>	15	AF525980	AF526014	424	0.1260	5	0.0035	0.0026	8	2	0.83	1.01	1.10
<i>PKCIZ</i>	2	AB026676	AY194147	740	0.1049	1	0.0007	0.0003	2	0	1.60	NA	NA
<i>PTCH</i>	6	AY139859	AY139861	530	0.0811	1	0.0001	0.0004	2	0	-1.10	NA	NA
<i>SPINZ</i>	2	AY194142	AY194143	666	0.1005	0	0.0000	0.0000	1	—	NA	NA	NA
<i>VLDL</i>	7	AY139848	AY139851	436	0.1075	4	0.0017	0.0021	5	0	-0.37	-1.26	-1.16
Total autosomes						214	0.0065	0.0058			0.44	1.11	1.01
Total Z chromosomes						44	0.0020	0.0014			1.23	1.21	1.48

Average divergence is the mean of all comparisons between individual chicken haplotypes and the turkey outgroup sequence. *L*, sequence length in base pairs; *S*, number of segregating sites; π , average difference per base pair between two randomly chosen sequences; θ_w , Watterson's estimate of θ per site; *H*, number of haplotypes; R_M , minimum number of recombination events; D_T , Tajima's *D*; D_{FL} , Fu and Li's *D*; F_{FL} , Fu and Li's *F*; * $P < 0.05$; *** $P < 0.001$.

scales >1 Mb (LERCHER *et al.* 2001). This implies that to obtain a representative genomic sample, choosing many segments of relatively short length should be preferable to the use of a limited number of longer segments (ELLEGREN *et al.* 2003). As the aim of the study was to get a broad idea of levels of polymorphism in the chicken genome and not to characterize the levels in particular breeds, we chose to analyze a few individuals from many different breeds rather than many individuals from one or only a few breeds. The sample also included two males of the red jungle fowl, the wild ancestor to domestic chicken. The total length of the sequence surveyed was ~15.2 kb, divided into 8316 bp from autosomes (14 introns) and 6856 bp from the Z chromosome (13 introns).

The total numbers of segregating sites were 214 on the autosomes (one every 39 bp) and 44 on the Z chromosome (one every 156 bp). For autosomal introns, average nucleotide diversity (π) calculated from the average number of pairwise differences was $6.5 \pm 0.3 \times 10^{-3}$ and Watterson's estimate of θ per site (θ_w ; basically, the proportion of segregating sites in a sample) was $5.8 \pm 0.8 \times 10^{-3}$. For Z chromosome introns, π was $2.0 \pm 0.1 \times 10^{-3}$ and θ_w was $1.4 \pm 0.4 \times 10^{-3}$. These observations suggest distinct differences in levels of genetic variability on autosomes and the Z chromosome of chicken; the A/Z ratio for π is 3.2 and for θ_w it is 4.1. Sixteen insertion or deletion polymorphisms (indels) were identified (excluding regions with length polymorphism in tandem repetitive DNA sequences);

13 indels were in autosomal and 3 were in Z-linked introns. For this type of polymorphism too, autosomal diversity thus seemed to exceed that of the Z chromosome. Details on polymorphism data are presented in Table 1.

We also estimated π for individual breeds. Although only a limited number of chromosomes per breed were sampled, it was evident that the trend of higher autosomal variability was present within breeds as well. The within-breed A/Z ratio for π was in the range of 2.4–11.1, with a mean of 5.4 ± 2.7 . A high A/Z ratio was also seen in the few red jungle fowl analyzed (3.11).

It should be noted that variation in the underlying rate of mutation is highly unlikely to account for reduced Z chromosome variation. Differences in the germline mutation rate between autosomes and sex chromosomes generally follow from the male mutation bias (α_m), leading to higher mutation rates the more time a chromosome class spends in the male germline (MIYATA *et al.* 1987; HURST and ELLEGREN 1998; LI *et al.* 2002). In birds, where α_m is estimated at 2–4 (ELLEGREN and FRIDOLFSSON 1997; KAHN and QUINN 1999; CARMICHAEL *et al.* 2000; BARTOSCH-HÄRLID *et al.* 2003), Z is two-thirds of the time in the male germline and should therefore be expected to mutate at a higher rate than autosomes, an assumption supported by sequence data (AXELSSON *et al.* 2004). If anything, the effect of this difference would be to give high levels of Z chromosome diversity, which is opposite to what we find in this study.

An HKA test was performed to test for evolutionary heterogeneity between intraspecific variation in chicken and interspecific divergence with the turkey outgroup, using all autosomal loci combined and all Z-linked loci combined (Table 2). This test took into account the mode of inheritance of the Z chromosome, which, assuming random mating, reduces its effective population size to three-quarters that of autosomal loci. Statistically significant deviations from the neutral model were observed ($P = 0.006$), suggesting that the observed ratio of average variation at autosomal and Z-linked loci is incompatible with a neutral model.

Tajima's D (TAJIMA 1989a) and Fu and Li's D and F (FU and LI 1993) statistics based on allele frequency spectra at all loci (Table 1) broadly indicate that a neutral model cannot be rejected. Only two loci (*RPL7a* intron 3 and *ALDOB* intron 7) have significant values for one or more of the test statistics, but this is not unexpected when multiple tests are performed, particularly when short sequences are used as is the case here. However, the average values of all test statistics, including Tajima's D , for both Z-linked and autosomal loci are positive, although not significant, indicating a weak excess of common variants.

Two further HKA tests were performed to examine heterogeneity (a) between all autosomal loci and (b) between all Z-linked loci (Table 2). There is no evidence for heterogeneity between autosomal loci ($P = 0.527$)

but Z chromosome loci exhibited significant deviations from the neutral model ($P = 0.024$). This suggests that, in addition to a reduced average variation on the Z chromosome compared to autosomal loci, there is evidence for heterogeneity within the Z chromosome.

DISCUSSION

There are two immediate conclusions of this study: (a) that Z-linked loci of domestic chicken have significantly reduced levels of variation in noncoding regions compared with autosomal loci and (b) that there is significant heterogeneity in levels of variation between loci on the Z chromosome. In the following we consider these observations in relation to effective population size, selection, and demography. However, it should be noted directly that the evolutionary history of chickens is complex, involving natural processes in populations of the wild ancestor as well as artificial selection and changes in population size and structure during domestication and breeding. Because of this, patterns of genetic variability in the chicken genome are likely to have a complex background, probably affected by processes prior to, during, and subsequent to domestication. Disentangling the relative importance of these processes may be difficult.

A low male effective population size is compatible with reduced Z chromosome variability: Under random mating, the effective population size of Z is three-quarters that of autosomes and polymorphism levels should thus be expected to scale accordingly. However, an HKA test demonstrates that our observation of three to four times higher average nucleotide variation at autosomal than at Z-linked loci is significantly greater than expected under neutrality ($P = 0.006$). A potential cause of this departure from the neutral model is that the assumptions of random mating are violated in chickens as domestication is likely to have been associated with skewed reproductive success among cocks, as is currently practiced in poultry breeding (MUIR and AGGREY 2003). Moreover, in the wild ancestor of the domestic chicken, the red jungle fowl, there is a high variance in male reproductive success with a suggested twofold difference in the effective population size of males and females (ZUK *et al.* 1990a,b; COLLIAS and COLLIAS 1996).

A low male effective population size will lead to a lowered effective population size of Z. If we conservatively assume that mutation rates at Z-linked and autosomal loci are the same, then the ratio of the number of autosomes to the number of Z chromosomes approaches 2 as the bias in the sex ratio of successfully reproducing birds becomes severe. This means that a twofold excess of polymorphism on autosomes compared to Z is the maximum possible difference one should expect from nonrandom mating. When twice as many females as males are contributing to the gene pool, as suggested by red jungle fowl data, the effective

TABLE 2
Results from HKA tests

Locus	Intron	Segregating sites		Average divergence	
		Observed	Expected	Observed	Expected
HKA test between autosomes and Z chromosome ($P = 0.006$)					
Total autosomes		204	157.97	755.98	802.01
Total Z chromosomes		42	88.03	591.36	545.338
HKA test within autosomes ($P = 0.527$)					
ACGR	F	13	12.18	44.38	45.20
CEPU	1	16	11.67	39.20	43.53
CLTC	5	8	13.36	55.20	49.84
DEF1	3	17	15.83	57.40	58.57
GHI	2	11	10.48	38.34	38.86
LHCGR	7	14	12.75	46.58	47.83
L37a	3	25	20.86	72.10	76.24
MLC	6	9	13.21	53.32	49.11
RHO	1	10	24.43	105.74	91.31
RPL5	4	9	11.52	45.52	43.00
RPL7a	3	13	11.82	42.18	43.36
RPL7a	4	7	5.84	20.84	22.00
RPL30	3	32	22.84	74.92	84.08
VIM	E	20	17.21	60.22	63.01
HKA test within Z chromosome ($P = 0.024$)					
AD012Z	1	3	5.54	83.6	81.06
AD012Z	2	0	2.10	34.00	31.90
ALDOB	1	2	1.64	24.32	24.68
ALDOB	5	7	3.16	43.02	46.86
ALDOB	6	2	0.98	13.7	14.72
ALDOB	7	4	2.35	33.66	35.31
ATP5A1Z	3	12	6.54	88.42	93.88
BRM	15	3	1.80	25.76	26.96
CHD1Z	15	5	3.62	52.92	54.30
PKCIZ	2	0	4.77	70.5	65.73
PTCH	6	0	2.16	9.00	6.84
SPINZ	2	0	4.20	66.00	61.80
VLDL	7	4	3.15	46.46	47.31

Twelve out of 258 SNPs from Table 1 were excluded from this analysis since they were within intron segments not found in turkey (*i.e.*, representing insertions in the chicken lineage or deletions in the turkey lineage).

population size of autosomes is 1.5 times that of Z; when there is 1 male for every 10 females it is 1.8 (22/12). However, even when an extreme operational sex ratio of 1:10 is incorporated in the HKA test, Z chromosome variability is still less than expected under neutrality ($P = 0.04$). We conclude that a reduced male effective population size is likely to have led to reduced levels of Z chromosome diversity in chicken, although it fails to account for all of the difference in polymorphism levels between autosomes and the Z.

Selection: We next consider the possibility of selection contributing to the observed discrepancy in polymorphism levels between autosomes and Z. As it would seem improbable that selection has independently *increased* variation at several unlinked autosomal loci (for example, due to balancing selection), it can be hypothesized that selection has *decreased* variation on the Z chromo-

some. The influence of selection on the Z chromosome is supported by an HKA test showing significant heterogeneity in polymorphism levels between loci on the Z chromosome ($P = 0.024$), which cannot be easily explained by differences in effective population size between chromosomal classes. Two possible factors could cause selection to have a greater effect on variation on the Z chromosome: (i) lower recombination rate on the Z chromosome resulting in a greater effect of selection on linked neutral variants and (ii) greater incidence of selection at Z-linked loci. We now examine the evidence for these two possibilities.

Positive selection and background selection both cause a reduction of genetic variation at linked neutral sites. During positive selection, the sweep through the population of an adaptive mutation will drag with it alleles at linked neutral sites. Fixation of the adaptive

mutation may thereby lead to the associated fixation of linked variants, often referred to as “genetic hitchhiking” (MAYNARD-SMITH and HAIGH 1974; KAPLAN *et al.* 1989). For background (negative, purifying) selection, the removal from the population of chromosomes containing deleterious alleles will also lead to removal of linked variants (CHARLESWORTH *et al.* 1993, 1995). A common prediction from both models is that the strength of the effect of selection on polymorphism levels at linked sites will be inversely proportional to the recombination rate. In other words, when there is more recombination, the association between a locus under selection and linked alleles will be uncoupled more quickly.

Data from chicken genome mapping provide evidence for significant heterogeneity in the sex-averaged recombination rate between autosomes and the Z chromosome (LEVIN *et al.* 1993; SMITH and BURT 1998; GROENEN *et al.* 2000; SCHMID *et al.* 2000): microchromosomes, ~ 5 cM/Mb; macrochromosomes, ~ 2.5 cM/Mb; Z chromosome, ~ 1.3 cM/Mb. The recombination rate on Z is thus ~ 2.5 times less than the average autosomal rate. This suggests that the effects of selection on linked neutral sites would stretch much farther on average from a locus under selection on the Z chromosome than from an autosomal locus. As a consequence, lower levels of variability on the Z chromosome than on autosomes are predicted.

If selection is an important factor to explain the contrasting levels of nucleotide diversity on chicken autosomes and the Z chromosome, is it through the action of selective sweeps or background selection? Our data are compatible with a scenario of selective sweeps as positive selection can be expected to be more effective in reducing neutral polymorphism on Z than on autosomes (BEGUN and WHITLEY 2000). In contrast, as background selection should be expected to quickly eliminate recessive deleterious mutations when exposed on the hemizygote Z chromosome, Z should have a larger proportion of mutation-free chromosomes than autosomes and, as a consequence, greater neutral polymorphism (CROW and KIMURA 1970; AQUADRO *et al.* 1994). However, the predictions made by the models assume equal rates of recombination in the two chromosome classes, which we know is not the case. Moreover, while the observed heterogeneity in polymorphism levels on Z would seem to suggest the action of selective sweeps, background selection can also cause heterogeneity in polymorphism levels when there is variation in the local rate of recombination (CHARLESWORTH *et al.* 1993, 1995).

It is also possible that the strength and frequency of selection differ between autosomes and the Z chromosomes. Important in this context are the observations of disproportionate sex linkage of traits involved in sexual selection (REINHOLD 1998; PIZZARI and BIRKHEAD 2002), speciation (ORR and COYNE 1989; COYNE 1992;

PRESGRAVES *et al.* 2003), and sexually antagonistic fitness variation (GIBSON *et al.* 2002), traits where adaptive evolution is often invoked. Sex chromosomes also show an unusual complement of sex-specific genes. Human X is enriched for genes with male-specific expression (SAIFI and CHANDRA 1999; WANG *et al.* 2001; LERCHER *et al.* 2003) whereas *Drosophila* (SWANSON *et al.* 2001; BETRAN *et al.* 2002; PARISI *et al.* 2003; RANZ *et al.* 2003) and *C. elegans* (REINKE *et al.* 2000) X's show a significant deficit of genes with male-biased expression and some evidence for an excess of genes with female-biased expression. Reproductive genes usually evolve rapidly under positive selection (CIVETTA and SINGH 1995; SWANSON and VACQUIER 2002; TORGERSON *et al.* 2002; SWANSON *et al.* 2003) and there is evidence for rapid evolution of male-biased gene expression in *Drosophila* (MEIKLEJOHN *et al.* 2003). It remains to be elucidated whether genes with sex-specific functions are overrepresented also on the avian Z chromosome.

Demography: Although selection can obviously shape levels and patterns of genomic variability, demography can too. Moreover, rejection of a neutral model in neutrality tests like Tajima's *D* (TAJIMA 1989a) does not necessarily imply that selection has to be invoked; it can as well result from violations of the assumptions of random mating in a population of constant size. For instance, a skewed spectrum of variants toward rare alleles, as indicated by negative values of Tajima's *D*, is expected from directional selection, from weak purifying selection, and from population expansion (TAJIMA 1989b; BRAVERMAN *et al.* 1995; PRZEWORSKI 2002). An excess of intermediate-frequency variants (positive Tajima's *D* values) is expected both from balancing selection and from population bottlenecks (TAJIMA 1989b; SIMONSEN *et al.* 1995; FAY and WU 1999).

Domestication has recently affected the size and structure of chicken populations, likely involving complex episodes of bottlenecks, population expansion, and subdivision. With such complex demographic history it is unclear what should be expected when it comes to the signature left on patterns of variation. We found average Tajima's *D* values to be positive for Z-linked and autosomal sequences but not significantly different from a neutral model. One possible interpretation is that there are two countering effects from population history. A recent contraction in population size associated with chicken domestication, indicated by the presence of only a limited number of mitochondrial DNA lineages (FUMIHITO *et al.* 1994), may have led to an excess of intermediate-frequency variants (rare alleles are more readily lost during a reduction in population size). Assuming an island model of population structure with weak migration between demes, population subdivision following from breed formation should also support positive values of Tajima's *D* (HAMMER *et al.* 2003). However, the breeding population of chicken has expanded significantly after the early days of domestication so

there may be many new mutations segregating at rare frequencies. Moreover, our sampling design with few individuals from many different demes should recover more rare alleles than a single large population sample (PTAK and PRZEWORSKI 2002; HAMMER *et al.* 2003).

The sign of Tajima's D subsequent to a bottleneck will depend on its severity and the length of time since the bottleneck (TAJIMA 1989b; FAY and WU 1999). The fact that, in this study, overall Tajima's D was found to still be positive may reflect that the bottleneck associated with domestication was strong and that relatively limited time has elapsed since population contraction. In any case, we cannot exclude the possibility that the effects of demographic history on allele frequency spectra have masked potential effects from selection.

Conclusions: We favor the idea that a low male effective population size has reduced the levels of polymorphism on the chicken Z chromosome. This could initially have arisen from skewed reproductive success among males in the wild ancestor, but is likely to have been accentuated during chicken breeding. However, as differences in effective population size do not seem to be able to completely explain the observed heterogeneity in polymorphism levels, selection—before, during, or subsequent to domestication—may have been important. Several lines of argument suggest that selective sweeps could be a potent force in shaping Z chromosome variability. In this context it is of interest to note that selection may have a stronger effect on polymorphism levels on Z in species with female heterogamety than on X in systems with male heterogamety, relative to that of the respective autosomes. As suggested by GIBSON *et al.* (2002) and formally demonstrated by REEVE and PFENNIG (2003), the process of sexual selection affecting sex-linked loci can be expected to be facilitated in birds as fathers pass a Z chromosome to their sons, improving the heritability and increasing the strength of selection of sexually antagonistic alleles advantageous to males (in systems with male heterogamety, fathers do not pass the X chromosome to sons). There is evidence for frequent Z-linkage of sexually selected characters in birds (OHNO 1967; SAETRE *et al.* 2003) as well as in butterflies (also ZW; SPERLING 1994; IYENGAR *et al.* 2002). Moreover, in chicken, several genes likely to have been under directional selection during domestication map to the Z chromosome, *e.g.*, the feathering (BITGOOD 1999) and dwarf (BURNSIDE *et al.* 1992) loci. Although speculative, the fact that the excess of polymorphism on autosomes compared to X in humans is less pronounced (~ 1.6 times more variation on autosomes) than that between autosomes and Z in birds could potentially be related to such a difference in the efficiency of selection.

The samples used in this study were kindly provided by Michèle Tixier-Boichard and the AVIANDIV project. Financial support was obtained from the Swedish Research Council. H.E. is a Royal Swedish Academy Research Fellow supported by a grant from the Knut and Alice Wallenberg foundation.

LITERATURE CITED

- AQUADRO, C. F., D. J. BEGUN and E. C. KINDAHL, 1994 Selection, recombination and DNA polymorphism in *Drosophila*, pp. 46–55 in *Nonneutral Evolution: Theories and Molecular Data*, edited by B. GOLDING. Chapman & Hall, London.
- AXELSSON, E., N. G. C. SMITH, H. SUNDSTRÖM, S. BERLIN and H. ELLEGREN, 2004 Male-biased mutation rate and divergence in autosomal, Z-linked and W-linked introns of chicken and turkey. *Mol. Biol. Evol.* (in press).
- BARTOSCH-HÄRLID, A., S. BERLIN, N. G. C. SMITH, A. P. MOLLER and H. ELLEGREN, 2003 Life history and the male mutation bias. *Evolution* **57**: 2398–2406.
- BEGUN, D. J., and P. WHITLEY, 2000 Reduced X-linked nucleotide polymorphism in *Drosophila simulans*. *Proc. Natl. Acad. Sci. USA* **97**: 5960–5965.
- BETRAN, E., K. THORNTON and M. LONG, 2002 Retroposed new genes out of the X in *Drosophila*. *Genome Res.* **12**: 1854–1859.
- BITGOOD, J. J., 1999 Linkage relationships of the Z-linked silver, slow feathering, and pop-eye loci. *Poult. Sci.* **78**: 1100–1101.
- BRAVERMAN, J. M., R. R. HUDSON, N. L. KAPLAN, C. H. LANGLEY and W. STEPHAN, 1995 The hitchhiking effect on the site frequency spectrum of DNA polymorphisms. *Genetics* **140**: 783–796.
- BURNSIDE, J., S. S. LIU, C. ZHONG and L. A. COGBURN, 1992 Abnormal growth hormone receptor gene expression in the sex-linked dwarf chicken. *Gen. Comp. Endocrinol.* **88**: 20–28.
- CARMICHAEL, A. N., A. K. FRIDOLFSSON, J. HALVERSON and H. ELLEGREN, 2000 Male-biased mutation rates revealed from Z and W chromosome-linked ATP synthase alpha-subunit (*ATP5A1*) sequences in birds. *J. Mol. Evol.* **50**: 443–447.
- CHARLESWORTH, B., M. T. MORGAN and D. CHARLESWORTH, 1993 The effect of deleterious mutations on neutral molecular variation. *Genetics* **134**: 1289–1303.
- CHARLESWORTH, D., B. CHARLESWORTH and T. M. MORGAN, 1995 The pattern of neutral molecular variation under the background selection model. *Genetics* **141**: 1619–1632.
- CHO, R. J., M. MINDRINOS, D. R. RICHARDS, R. J. SAPOLSKY, M. ANDERSON *et al.*, 1999 Genome-wide mapping with biallelic markers in *Arabidopsis thaliana*. *Nat. Genet.* **23**: 203–207.
- CIVETTA, A., and R. S. SINGH, 1995 High divergence of reproductive tract proteins and their association with postzygotic reproductive isolation in *Drosophila melanogaster* and *Drosophila virilis* group species. *J. Mol. Evol.* **41**: 1085–1095.
- COLLIAS, N. E., and E. COLLIAS, 1996 Social organization of a red junglefowl, *Gallus gallus*, population related to evolution theory. *Anim. Behav.* **51**: 1337–1354.
- COYNE, J. A., 1992 Genetics and speciation. *Nature* **355**: 511–515.
- CROW, J. F., and M. KIMURA, 1970 *An Introduction to Population Genetics Theory*. Harper & Row, New York.
- DIMICHEFF, D. E., S. V. DROVETSKI and D. P. MINDELL, 2002 Phylogeny of *Tetraoninae* and other galliform birds using mitochondrial 12S and ND2 genes. *Mol. Phylogenet. Evol.* **24**: 203–215.
- EBERSBERGER, I., D. METZLER, C. SCHWARZ and S. PAABO, 2002 Genomewide comparison of DNA sequences between humans and chimpanzees. *Am. J. Hum. Genet.* **70**: 1490–1497.
- ELLEGREN, H., and A. K. FRIDOLFSSON, 1997 Male-driven evolution of DNA sequences in birds. *Nat. Genet.* **17**: 182–184.
- ELLEGREN, H., N. G. C. SMITH and M. T. WEBSTER, 2003 Mutation rate variation in the mammalian genome. *Curr. Opin. Genet. Dev.* **13**: 562–568.
- FAY, J. C., and C.-I. WU, 1999 A human population bottleneck can account for the discordance between patterns of mitochondrial versus nuclear DNA variation. *Mol. Biol. Evol.* **16**: 1003–1005.
- FRIDOLFSSON, A. K., H. CHENG, N. G. COPELAND, N. A. JENKINS, H. C. LIU *et al.*, 1998 Evolution of the avian sex chromosomes from an ancestral pair of autosomes. *Proc. Natl. Acad. Sci. USA* **95**: 8147–8152.
- FU, Y.-X., and W.-H. LI, 1993 Statistical tests of neutrality of mutations. *Genetics* **133**: 693–709.
- FUMIHIITO, A., T. MIYAKE, S. SUMI, M. TAKADA and S. OHNO, 1994 One subspecies of the red junglefowl (*Gallus gallus gallus*) suffices as the matriarchic ancestor of all domestic breeds. *Proc. Natl. Acad. Sci. USA* **91**: 12505–12509.
- GIBSON, J. R., A. K. CHIPPINDALE and W. R. RICE, 2002 The X chromosome is a hot spot for sexually antagonistic fitness variation. *Proc. R. Soc. Lond. Ser. B* **269**: 499–505.

- GROENEN, M. A., H. H. CHENG, N. BUMSTEAD, B. F. BENKEL, W. E. BRILES *et al.*, 2000 A consensus linkage map of the chicken genome. *Genome Res.* **10**: 137–147.
- HAMMER, M. F., F. BLACKMER, D. GARRIGAN, M. W. NACHMAN and J. A. WILDER, 2003 Human population structure and its effects on sampling Y chromosome sequence variation. *Genetics* **164**: 1495–1509.
- HARDISON, R. C., K. M. ROSKIN, S. YANG, M. DIEKHANS, W. J. KENT *et al.*, 2003 Covariation in frequencies of substitution, deletion, transposition, and recombination during eutherian evolution. *Genome Res.* **13**: 13–26.
- HILLEL, J., M. A. GROENEN, M. TIXIER-BOICHARD, A. B. KOROL, L. DAVID *et al.*, 2003 Biodiversity of 52 chicken populations assessed by microsatellite typing of DNA pools. *Genet. Sel. Evol.* **35**: 533–557.
- HUDSON, R. R., M. KREITMAN and M. AGUADÉ, 1987 A test of neutral molecular evolution based on nucleotide data. *Genetics* **116**: 153–159.
- HURST, L. D., and H. ELLEGREN, 1998 Sex biases in the mutation rate. *Trends Genet.* **14**: 446–452.
- INTERNATIONAL HUMAN GENOME SEQUENCING CONSORTIUM, 2002 Initial sequencing and comparative analysis of the mouse genome. *Nature* **420**: 520–562.
- INTERNATIONAL SNP MAP WORKING GROUP, 2001 A map of human genome sequence variation containing 1.42 million single nucleotide polymorphisms. *Nature* **409**: 928–933.
- IYENGAR, V. K., H. K. REEVE and T. EISNER, 2002 Paternal inheritance of a female moth's mating preference. *Nature* **419**: 830–832.
- KAHN, N. W., and T. W. QUINN, 1999 Male-driven evolution among Eoaves? A test of the replicative division hypothesis in a heterogametic female (ZW) system. *J. Mol. Evol.* **49**: 750–759.
- KAPLAN, N. L., R. R. HUDSON and C. H. LANGLEY, 1989 The "hitchhiking effect" revisited. *Genetics* **123**: 887–899.
- KOCH, R., H. G. VAN LUENEN, M. VAN DER HORST, K. L. THIJSEN and R. H. PLASTERK, 2000 Single nucleotide polymorphisms in wild isolates of *Caenorhabditis elegans*. *Genome Res.* **10**: 1690–1696.
- LERCHER, M. J., E. J. WILLIAMS and L. D. HURST, 2001 Local similarity in evolutionary rates extends over whole chromosomes in human-rodent and mouse-rat comparisons: implications for understanding the mechanistic basis of the male mutation bias. *Mol. Biol. Evol.* **18**: 2032–2039.
- LERCHER, M. J., A. O. URRUTIA and L. D. HURST, 2003 Evidence that the human X chromosome is enriched for male-specific but not female-specific genes. *Mol. Biol. Evol.* **20**: 1113–1116.
- LEVIN, I., L. B. CRITTENDEN and J. B. DODGSON, 1993 Genetic map of the chicken Z chromosome using random amplified polymorphic DNA (RAPD) markers. *Genomics* **16**: 224–230.
- LI, W. H., S. YI and K. MAKOVA, 2002 Male-driven evolution. *Curr. Opin. Genet. Dev.* **12**: 650–656.
- LIU, W., M. G. KAISER and S. J. LAMONT, 2003 Natural resistance-associated macrophage protein 1 gene polymorphisms and response to vaccine against or challenge with *Salmonella enteritidis* in young chicks. *Poult. Sci.* **82**: 259–266.
- MAYNARD-SMITH, J., and J. HAIGH, 1974 The hitch-hiking effect of a favourable gene. *Genet. Res.* **23**: 23–35.
- MEIKLEJOHN, C. D., J. PARSCH, J. M. RANZ and D. L. HARTL, 2003 Rapid evolution of male-biased gene expression in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **100**: 9894–9899.
- MIYATA, T., H. HAYASHIDA, K. KUMA, K. MITSUYASU, T. YASUNAGA *et al.*, 1987 Male-driven molecular evolution: a model and nucleotide sequence analysis. *Cold Spring Harbor Symp. Quant. Biol.* **52**: 863–867.
- MUIR, W. M., and S. E. AGGREY, 2003 *Poultry Genetics, Breeding and Biotechnology*. CABI, New York.
- MULLIKIN, J. C., S. E. HUNT, C. G. COLE, B. J. MORTIMORE, C. M. RICE *et al.*, 2000 An SNP map of human chromosome 22. *Nature* **407**: 516–520.
- NANDA, I., T. HAAF, M. SCHARTL, M. SCHMID and D. W. BURT, 2002 Comparative mapping of Z-orthologous genes in vertebrates: implications for the evolution of avian sex chromosomes. *Cytogenet. Genome Res.* **99**: 178–184.
- OHNO, S., 1967 *Sex Chromosomes and Sex-Linked Genes*. Springer, Berlin/New York.
- ORR, H. A., and J. A. COYNE, 1989 The genetics of postzygotic isolation in the *Drosophila virilis* group. *Genetics* **121**: 527–537.
- PARISI, M., R. NUTTALL, D. NAIMAN, G. BOUFFARD, J. MALLEY *et al.*, 2003 Paucity of genes on the *Drosophila* X chromosome showing male-biased expression. *Science* **299**: 697–700.
- PARSANEJAD, R., D. ZADWORNÝ and U. KUHNLEIN, 2002 Genetic variability of the cytosolic phosphoenolpyruvate carboxykinase gene in white leghorn chickens. *Poult. Sci.* **81**: 1668–1670.
- PATIL, N., A. J. BERNO, D. A. HINDS, W. A. BARRETT, J. M. DOSHI *et al.*, 2001 Blocks of limited haplotype diversity revealed by high-resolution scanning of human chromosome 21. *Science* **294**: 1719–1723.
- PIZZARI, T., and T. R. BIRKHEAD, 2002 The sexually-selected sperm hypothesis: sex-biased inheritance and sexual antagonism. *Biol. Rev.* **77**: 183–209.
- PRESGRAVES, D. C., L. BALAGOPALAN, S. M. ABMAYR and H. A. ORR, 2003 Adaptive evolution drives divergence of a hybrid inviability gene between two species of *Drosophila*. *Nature* **423**: 715–719.
- PRZEWORSKI, M., 2002 The signature of positive selection at randomly chosen loci. *Genetics* **160**: 1179–1189.
- PTAK, S. E., and M. PRZEWORSKI, 2002 Evidence for population growth in humans is confounded by fine-scale population structure. *Trends Genet.* **18**: 559–563.
- RANZ, J. M., C. I. CASTILLO-DAVIS, C. D. MEIKLEJOHN and D. L. HARTL, 2003 Sex-dependent gene expression and evolution of the *Drosophila* transcriptome. *Science* **300**: 1742–1745.
- REEVE, H. K., and D. W. PFENNIG, 2003 Genetic biases for showy males: Are some genetic systems especially conducive to sexual selection? *Proc. Natl. Acad. Sci. USA* **100**: 1089–1094.
- REICH, D. E., S. F. SCHAFFNER, M. J. DALY, G. MCVEAN, J. C. MULLIKIN *et al.*, 2002 Human genome sequence variation and the influence of gene history, mutation and recombination. *Nat. Genet.* **32**: 135–142.
- REINHOLD, K., 1998 Sex linkage among genes controlling sexually selected traits. *Behav. Ecol. Sociobiol.* **44**: 1–7.
- REINKE, V., H. E. SMITH, J. NANCE, J. WANG, C. VAN DOREN *et al.*, 2000 A global profile of germline gene expression in *C. elegans*. *Mol. Cell* **6**: 605–616.
- ROSENBERG, N. A., T. BURKE, K. ELO, M. W. FELDMAN, P. J. FREIDLIN *et al.*, 2001 Empirical evaluation of genetic clustering methods using multilocus genotypes from 20 chicken breeds. *Genetics* **159**: 699–713.
- ROZAS, J., and R. ROZAS, 1999 DnaSP version 3: an integrated program for molecular population genetics and molecular evolution analysis. *Bioinformatics* **15**: 174–175.
- SAETRE, G. P., T. BORGE, K. LINDROOS, J. HAAVIE, B. C. SHELDON *et al.*, 2003 Sex chromosome evolution and speciation in *Ficedula* flycatchers. *Proc. R. Soc. Lond. Ser. B* **270**: 53–59.
- SAIFI, G. M., and H. S. CHANDRA, 1999 An apparent excess of sex- and reproduction-related genes on the human X chromosome. *Proc. R. Soc. Lond. Ser. B* **266**: 203–209.
- SCHMID, M., I. NANDA, M. GUTTENBACH, C. STEINLEIN, M. HOEHN *et al.*, 2000 First report on chicken genes and chromosomes 2000. *Cytogenet. Cell Genet.* **90**: 169–218.
- SCHNEIDER, S., D. ROESSLI and L. EXCOFIER, 2000 *Arlequin: A Software for Population Genetics Data Analysis*. Genetics and Biometry Lab, Department of Anthropology, University of Geneva, Geneva.
- SIMONSEN, K. L., G. A. CHURCHILL and C. F. AQUADRO, 1995 Properties of statistical tests of neutrality for DNA polymorphism data. *Genetics* **141**: 413–429.
- SMITH, E. J., L. SHI, P. DRUMMOND, L. RODRIGUEZ, R. HAMILTON *et al.*, 2001 Expressed sequence tags for the chicken genome from a normalized 10-day-old White Leghorn whole embryo cDNA library. 1. DNA sequence characterization and linkage analysis. *J. Hered.* **92**: 1–8.
- SMITH, E. J., L. SHI and G. SMITH, 2002 Expressed sequence tags for the chicken genome from a normalized 10-day-old white leghorn whole-embryo cDNA library. 3. DNA sequence analysis of genetic variation in commercial chicken populations. *Genome* **45**: 261–267.
- SMITH, J., and D. W. BURT, 1998 Parameters of the chicken genome (*Gallus gallus*). *Anim. Genet.* **29**: 290–294.
- SMITH, N. G., M. T. WEBSTER and H. ELLEGREN, 2002 Deterministic mutation rate variation in the human genome. *Genome Res.* **12**: 1350–1356.
- SPELTING, F. A. H., 1994 Sex-linked genes and species-differences in *Lepidoptera*. *Can. Entomol.* **126**: 807–818.
- SWANSON, W. J., and V. D. VACQUIER, 2002 The rapid evolution of reproductive proteins. *Nat. Rev. Genet.* **3**: 137–144.

- SWANSON, W. J., A. G. CLARK, H. M. WALDRIP-DAIL, M. F. WOLFNER and C. F. AQUADRO, 2001 Evolutionary EST analysis identifies rapidly evolving male reproductive proteins in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **98**: 7375–7379.
- SWANSON, W. J., R. NIELSEN and Q. YANG, 2003 Pervasive adaptive evolution in mammalian fertilization proteins. *Mol. Biol. Evol.* **20**: 18–20.
- TAJIMA, F., 1989a Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* **123**: 585–595.
- TAJIMA, F., 1989b The effect of change in population size on DNA polymorphism. *Genetics* **123**: 597–601.
- TENAILLON, M. L., M. C. SAWKINS, A. D. LONG, R. L. GAUT, J. F. DOEBLEY *et al.*, 2001 Patterns of DNA sequence polymorphism along chromosome 1 of maize (*Zea mays* ssp. *mays* L.). *Proc. Natl. Acad. Sci. USA* **98**: 9161–9166.
- THOMPSON, J. D., D. G. HIGGINS and T. J. GIBSON, 1994 CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**: 4673–4680.
- TORGERSON, D. G., R. J. KULATHINAL and R. S. SINGH, 2002 Mammalian sperm proteins are rapidly evolving: evidence of positive selection in functionally diverse genes. *Mol. Biol. Evol.* **19**: 1973–1980.
- WADE, C. M., E. J. KULBOKAS, A. W. KIRBY, M. C. ZODY, J. C. MULLIKIN *et al.*, 2002 The mosaic structure of variation in the laboratory mouse genome. *Nature* **420**: 574–578.
- WANG, P. J., J. R. MCCARREY, F. YANG and D. C. PAGE, 2001 An abundance of X-linked genes expressed in spermatogonia. *Nat. Genet.* **27**: 422–426.
- ZHU, Y. L., Q. J. SONG, D. L. HYTEN, C. P. VAN TASSELL, L. K. MATUKUMALLI *et al.*, 2003 Single-nucleotide polymorphisms in soybean. *Genetics* **163**: 1123–1134.
- ZUK, M., K. JOHNSON, R. THORNHILL and J. D. LIGON, 1990a Mechanisms of female choice in red jungle fowl. *Evolution* **44**: 477–485.
- ZUK, M., R. THORNHILL, J. D. LIGON, K. JOHNSON, S. AUSTAD *et al.*, 1990b The role of male ornaments and courtship behavior in female mate choice of red jungle fowl. *Am. Nat.* **136**: 459–473.

Communicating editor: N. A. JENKINS

