

Supporting information for Baechler *et al.* (2003) *Proc. Natl. Acad. Sci. USA*,  
10.1073/pnas.0337679100

**Table 7. Comparison of clinical features between IFN-high and IFN-low patients**

	Summation	Euclidean distance	Subtract mean/median	Ratio vs. mean	Ratio vs. median
Age at diagnosis (yrs)	IFN-high 25.2±12.2	IFN-high 27.2±14.6	IFN-high 25.2±12.2	IFN-high 27.5±14.5	IFN-high 29.8±15.1
	IFN-low 30.0±13.3	IFN-low 28.0±11.1	IFN-low 30.0±13.3	IFN-low 27.8±11.3	IFN-low 25.5±10.0
	<i>t</i> test 0.192	<i>t</i> test 0.817	<i>t</i> test 0.192	<i>t</i> test 0.938	<i>t</i> test 0.252
# Criteria	IFN-high 6.8±1.2	IFN-high 6.8±1.3	IFN-high 6.8±1.2	IFN-high 6.6±1.4	IFN-high 6.6±1.4
	IFN-low 5.7±1.1	IFN-low 5.7±1.1	IFN-low 5.7±1.1	IFN-low 5.8±1.1	IFN-low 5.8±1.0
	<i>t</i> test 0.002	<i>t</i> test 0.004	<i>t</i> test 0.002	<i>t</i> test 0.032	<i>t</i> test 0.032
% Renal/ CNS	IFN-high 62.5%	IFN-high 58.3%	IFN-high 62.5%	IFN-high 54.2%	IFN-high 50.0%
	IFN-low 20.8%	IFN-low 25.0%	IFN-low 20.8%	IFN-low 29.2%	IFN-low 33.3%
	Chi-square 7.7E-06	Chi-square 0.001	Chi-square 7.7E-06	Chi-square 0.023	Chi-square 0.160
% Hematologic	IFN-high 75.0%	IFN-high 70.8%	IFN-high 75.0%	IFN-high 70.8%	IFN-high 70.8%
	IFN-low 20.8%	IFN-low 25.0%	IFN-low 20.8%	IFN-low 25.0%	IFN-low 25.0%
	Chi-square 6.1E-09	Chi-square 7.1E-06	Chi-square 6.1E-09	Chi-square 7.1E-06	Chi-square 7.1E-06
Serositis	IFN-high 66.7%	IFN-high 66.7%	IFN-high 66.7%	IFN-high 66.7%	IFN-high 62.5%
	IFN-low 75.0%	IFN-low 75.0%	IFN-low 75.0%	IFN-low 75.0%	IFN-low 79.2%
	Chi-square 0.637	Chi-square 0.637	Chi-square 0.637	Chi-square 0.637	Chi-square 0.359

