

Table 9. Correlation coefficients of clinical features with IFN scores

	Summation	Euc. D. vs. mean	Euc. D. vs. median	Subtract mean	Subtract median	Ratio vs. mean	Ratio vs. median
# SLE criteria	0.51 (0.00024) ^a	0.52 (0.00017)	0.52 (0.00018)	0.50 (0.00031)	0.50 (0.00030)	0.35 (0.016)	0.39 (0.0062)
Renal	0.26 (0.070)	0.24 (0.099)	0.24 (0.094)	0.23 (0.116)	0.24 (0.102)	0.14 (0.340)	0.21 (0.162)
CNS	0.13 (0.363)	0.12 (0.427)	0.12 (0.417)	0.12 (0.416)	0.12 (0.399)	0.07 (0.651)	0.12 (0.417)
Renal and/or CNS	0.37 (0.009)	0.34 (0.016)	0.35 (0.015)	0.34 (0.018)	0.35 (0.015)	0.21 (0.144)	0.32 (0.026)
Hematologic	0.53 (0.00012)	0.51 (0.00021)	0.51 (0.00019)	0.51 (0.00023)	0.51 (0.00019)	0.34 (0.019)	0.45 (0.00148)
Serositis	-0.12 (0.404)	-0.13 (0.390)	-0.13 (0.390)	-0.13 (0.379)	-0.13 (0.382)	-0.06 (0.698)	-0.11 (0.436)
anti-dsDNA	0.35 (0.014)	0.35 (0.014)	0.35 (0.015)	0.35 (0.016)	0.35 (0.016)	0.10 (0.489)	0.19 (0.194)
Age at diagnosis	-0.22 (0.138)	-0.19 (0.205)	-0.18 (0.209)	-0.19 (0.189)	-0.20 (0.184)	0.07 (0.641)	0.02 (0.916)
0 major criteria	-0.44 (0.002)	-0.41 (0.004)	-0.42 (0.003)	-0.41 (0.004)	-0.42 (0.003)	-0.19 (0.185)	-0.35 (0.015)
1 major criteria	0.01 (0.934)	-0.01 (0.927)	-0.01 (0.952)	-0.01 (0.953)	0.00 (0.987)	-0.18 (0.230)	-0.04 (0.789)
2/3 major criteria	0.45 (0.0012)	0.44 (0.0017)	0.44 (0.0016)	0.44 (0.0020)	0.44 (0.0017)	0.37 (0.0106)	0.40 (0.0047)

^aData are expressed as r values followed by P values from linear regression.