

Update on Abiotic Stress

Betaines and Related Osmoprotectants. Targets for Metabolic Engineering of Stress Resistance¹

Scott D. McNeil, Michael L. Nuccio, and Andrew D. Hanson*

Horticultural Sciences Department, University of Florida, Gainesville, Florida 32611

Osmoprotectants (also termed compatible solutes) occur in all organisms from archaeobacteria to higher plants and animals. They are highly soluble compounds that carry no net charge at physiological pH and are nontoxic at high concentrations. Osmoprotectants serve to raise osmotic pressure in the cytoplasm and can also stabilize proteins and membranes when salt levels or temperatures are unfavorable. Osmoprotectants therefore play important roles in the adaptation of cells to various adverse environmental conditions (Yancey, 1994). Chemically, there are three types: betaines and allied compounds, polyols and sugars (e.g. mannitol and trehalose), and amino acids such as Pro. This *Update* will focus on betaines and their biosynthetic pathways. Advances in polyol and Pro research are covered elsewhere (Delauney and Verma, 1993; Stoop et al., 1996).

Betaines are amino acid derivatives in which the nitrogen atom is fully methylated, i.e. they are quaternary ammonium compounds. Figure 1 shows the structures of the three best-known betaines from plants, Gly betaine, Pro betaine (stachydrine), and β -Ala betaine, as well as choline-*O*-sulfate and DMSP (Rhodes and Hanson, 1993). The last two compounds are strictly speaking not betaines because DMSP has a tertiary sulfonium in place of a quaternary ammonium group, and choline-*O*-sulfate has a sulfate ester instead of a carboxyl group—but they are close structural analogs of betaines and have similar chemical and physiological properties.

The compounds in Figure 1 differ in their taxonomic distribution (Blunden and Gordon, 1986; Rhodes and Hanson, 1993). For instance, Gly betaine is widespread among both flowering plants and algae, whereas DMSP is rare in higher plants but common in algae. Certain crop plants such as rice, soybeans, and potatoes lack significant amounts of betaines or any other osmoprotectant. This deficiency is the rationale for recent interest in using metabolic engineering technology to install the synthesis of

osmoprotectants in such crops in order to improve their tolerance to drought, salinity, and other stresses.

The levels of betaines and other osmoprotectants typically rise during exposure to stresses such as salinity, water deficit, and low temperature because the biosynthetic enzymes are stress induced. Osmoprotectants are largely confined to the cytoplasm (including organelles) and are almost absent from the vacuole, which generally occupies about 90% of the cell volume. For example, the halophyte *Atriplex gmelini* was found to have 320 mM Gly betaine in the cytoplasm, but only 0.24 mM in the vacuole (Matoh et al., 1987). Isolated chloroplasts of various species have also been shown to contain high concentrations of Gly betaine or DMSP, particularly when isolated from salt-stressed plants; Figure 1B illustrates this point with data from three species.

MAGNITUDE AND MECHANISM OF PROTECTIVE EFFECTS

The protective properties of betaines were first recognized in experiments in which they were supplied to bacteria whose growth was inhibited by high salt concentrations (Le Rudulier et al., 1984). Typical data for Gly betaine and DMSP are shown in Figure 2: In media containing 0.6 M NaCl, the bacteria grow very slowly unless supplied with one of these compounds, which they take up from the medium and accumulate to intracellular levels of >1 M. The physicochemical basis for this striking osmoprotective effect is not fully understood, but there is good evidence that it lies partly in the exclusion of osmoprotectant molecules from the water layer in contact with protein surfaces (Timasheff, 1992). This creates a situation in which native (i.e. folded) protein structures are thermodynamically favored because they present the least possible surface area to the water. Most other solutes such as NaCl or MgSO₄ interact directly with protein surfaces and favor unfolding, which leads to denaturation. Osmoprotectants also have cryoprotectant and heat-protectant properties, and exclusion from protein surfaces is probably part of the protective mechanism in these cases as well (Carpenter and Crowe, 1988; Winzor et al., 1992).

Abbreviations: BADH, betaine aldehyde dehydrogenase; CMO, choline monooxygenase; DMSP, 3-dimethylsulfoniopropionate; DMSP-aldehyde, 3-dimethylsulfoniopropionaldehyde; DMSP-amine, 3-dimethylsulfoniopropylamine; SMM, S-methylmethionine.

¹ This research was supported in part by grants to A.D.H. from the U.S. Department of Agriculture National Research Initiative Competitive Grants Program (no. 98-35100-6149) and the National Science Foundation (nos. IBN-9816075 and IBN-9813999), by an endowment from the C.V. Griffin, Sr., Foundation, and by the Florida Agricultural Experiment Station. This article is journal series no. R-06838.

* Corresponding author; e-mail adha@gnv.ifas.ufl.edu; fax 352-392-6479.




Figure 1. A, Chemical structures of betaines and related compounds that occur in flowering plants and marine algae. B, Concentrations of Glycine betaine or DMSP measured in chloroplasts isolated from leaves of salt-tolerant plants grown in nonsaline (black bars) or saline (white bars) conditions. Data are from Robinson and Jones (1986), Génard et al. (1991), and Trossat et al. (1998). Conc., Concentration.

BIOSYNTHETIC PATHWAYS, ENZYMES, AND GENES

Detailed knowledge of biochemical pathways is a prerequisite for metabolic engineering. Biosynthetic routes have now been established for all of the compounds shown in Figure 1. Most of the enzymes participating in these pathways have been identified, and genes for some of them have been cloned. The following is a summary of the current status of knowledge for each compound.

Glycine Betaine

Glycine betaine occurs in diverse marine algae and at least 10 flowering plant families, including Chenopodiaceae, Amaranthaceae, Gramineae, Compositae, and Malvaceae (Blunden and Gordon, 1986; Rhodes and Hanson, 1993). Its synthesis has been studied mainly in species of Chenopodiaceae, and to some extent in Amaranthaceae and Gramineae. In these cases, Glycine betaine is synthesized via a two-step oxidation of choline (Fig. 3). In Chenopodiaceae and Amaranthaceae, the first step (choline → betaine aldehyde) is mediated by CMO, an unusual Fd-dependent monooxygenase with a Rieske-type iron-sulfur cluster, which has been characterized and cloned. The active en-

zyme is a dimer or trimer of identical 43-kD subunits and is localized in the chloroplast stroma (Burnet et al., 1995; Rathinasabapathi et al., 1997). Because reduced Fd is generated by photosynthetic electron transport, CMO activity in vivo is strongly light dependent. Both drought and salinity induce CMO expression (Rathinasabapathi et al., 1997; Russell et al., 1998). As CMO has only been found in Chenopodiaceae and Amaranthaceae (Russell et al., 1998), other families may have different choline-oxidizing enzymes that may not be located in plastids. Could these families have choline dehydrogenases or choline oxidases similar to the enzymes known to catalyze choline oxidation in bacteria (Le Rudulier et al., 1984; Hayashi et al., 1997)?

The second step of Glycine betaine synthesis is catalyzed by BADH (Fig. 3), an NAD-dependent dehydrogenase that has been characterized and cloned from Chenopodiaceae, Amaranthaceae, and Gramineae. BADH is not unique to Glycine betaine synthesis; it also attacks DMSP-aldehyde (see below) and seems to be identical to ω-aminoaldehyde dehydrogenase, a ubiquitous enzyme of polyamine metabolism (Trossat et al., 1997). This probably explains why plants with no Glycine betaine have some BADH activity (Holmstrom et al., 1994; Rathinasabapathi et al., 1994). BADH is a dimer of identical 54-kD subunits. It is a stromal enzyme in Chenopodiaceae, and, surprisingly, lacks a typical transit peptide. Nonetheless, BADH is efficiently targeted to chloroplasts in transgenic tobacco (Rathinasabapathi et al., 1994). In Gramineae, however, most of the BADH may be peroxisomal (Nakamura et al., 1997), which reinforces the idea that the choline-oxidizing enzyme in this family may not be CMO, as CMO could not function in peroxisomes where there is no reduced Fd. BADH is induced by osmotic stress in both Chenopodiaceae and Gramineae (Rhodes and Hanson, 1993).

Choline-O-Sulfate

Choline-O-sulfate occurs throughout the Plumbaginaceae (Hanson et al., 1994) and in some marine algae (Blunden




Figure 2. An illustration of the osmoprotective effect of Glycine betaine or DMSP on cells of the bacterium *Escherichia coli* cultured in a highly saline medium (0.6 M NaCl). Growth in the presence of 1 mM Glycine betaine or DMSP was far more rapid than in their absence (control). These compounds accumulated to intracellular levels of >1 M. Bacterial growth was measured by the optical density of the cultures at 420 nm. Data are from Paquet et al. (1994).




Figure 3. Biosynthetic pathways of Gly betaine in Chenopodiaceae and of choline-*O*-sulfate in Plumbaginaceae. The product of the CMO reaction is the hydrate form of betaine aldehyde. CST, Choline sulfotransferase; PAP, 3'-phosphoadenosine-5'-phosphate; PAPS, 3'-phosphoadenosine-5'-phosphosulfate.

and Gordon, 1986). In addition to being an osmoprotectant, it also provides a means of detoxifying sulfate, which is often abundant in saline environments (Hanson et al., 1994). The conversion of choline to choline-*O*-sulfate is catalyzed by a 3'-phosphoadenosine-5'-phosphosulfate-dependent choline sulfotransferase (Fig. 3) (Rivoal and Hanson, 1994). This sulfotransferase is a soluble enzyme with an extremely high affinity for choline ($K_m = 5.5 \mu\text{M}$), which probably reflects the low cytoplasmic choline levels that prevail in plants (Nuccio et al., 1998). Saline conditions induce sulfotransferase enzyme activity in both leaves and roots (Rivoal and Hanson, 1994).

Pro Betaine and β -Ala Betaine

β -Ala betaine and Pro betaine are found in some species of Plumbaginaceae; Pro betaine also occurs in Rutaceae, Leguminosae, Compositae, and other families (Rhodes and Hanson, 1993; Hanson et al., 1994; Nolte et al., 1997). Both compounds are also found in marine algae (Blunden and Gordon, 1986) and are known to be formed by methylation of the corresponding amino acid; however, the enzyme(s) involved have yet to be identified (Essery et al., 1962; Hanson et al., 1994). For Pro betaine, *in vivo* radiolabeling and phytochemical data suggest that the two methylations are catalyzed by different enzymes (Essery et al., 1962; Naidu et al., 1987).

DMSP

DMSP biosynthesis is important not only in relation to osmoprotection but also because the DMSP produced by marine algae is the precursor of atmospheric dimethylsulfide gas. This gas has a key role in the global sulfur cycle and influences climate (Malin and Kirst, 1997). DMSP is present in diverse marine algae (Blunden and Gordon, 1986; Malin and Kirst, 1997), but has so far been found in only a few flowering plants: *Spartina* spp. and sugarcane from the Gramineae and *Wollastonia biflora* from the Compositae. In all cases DMSP is synthesized from Met. However, conversion of Met to DMSP seems to have evolved independently three times, once in algae and twice in

flowering plants, because there are three different pathways (Fig. 4).

DMSP Synthesis in Algae

The steps involved in DMSP synthesis have been demonstrated in the green macroalga *Enteromorpha intestinalis*, and appear to be the same in diverse microalgae (Gage et al., 1997). They are: transamination of Met to give the corresponding 2-oxo acid, followed by reduction to a 2-hydroxy acid, then *S*-methylation and oxidative decarboxylation (Fig. 4). The enzymes catalyzing the first three steps have been demonstrated *in vitro* (Summers et al., 1998); *in vivo* $^{18}\text{O}_2$ -labeling data indicate that the final step is mediated by an oxygenase (Gage et al., 1997).

DMSP Synthesis in Flowering Plants

DMSP synthesis in flowering plants has been investigated in *W. biflora* and *S. alterniflora*, which have somewhat different pathways. In both species, the first step is conversion of Met to SMM, catalyzed by Met *S*-methyltransferase, and the last is oxidation of DMSP-aldehyde, catalyzed by




Figure 4. Biosynthetic pathways of DMSP in marine algae and flowering plants. The algal pathway has been found in the green macroalga *E. intestinalis* and various plankton species. Two related pathways have been found in flowering plants, one in the saltmarsh grass *S. alterniflora* and the other in the Pacific coastal plant *W. biflora*. The breakdown of DMSP to give dimethylsulfide (DMS) and acrylate is also shown; this reaction has so far only been demonstrated in marine algae and bacteria. DMSPald, DMSP-aldehyde; D-DMSHB, D-4-dimethylsulfonio-2-hydroxybutyrate; D-MTHB, D-4-methylthio-2-hydroxybutyrate; MAT, Met aminotransferase; MMT, Met *S*-methyltransferase; MTHBMT, 4-methylthio-2-hydroxybutyrate *S*-methyltransferase; MTOB, 4-methylthio-2-oxobutyrate; MTOBR, MTOB reductase.

BADH (Trossat et al., 1996, 1997; Kocsis et al., 1998). Met S-methyltransferase has been shown to be cytosolic and has been cloned (Trossat et al., 1996; Bourgis et al., 1999). In *W. biflora*, SMM is apparently converted to DMSP-aldehyde in a single step via an unusual coupled transamination-decarboxylation reaction (Fig. 4) (Rhodes et al., 1997). This conversion is known to occur in the chloroplast (Trossat et al., 1996), but the enzyme(s) involved has not yet been identified. On the other hand, *S. alterniflora* converts SMM to DMSP-aldehyde in two steps via the intermediate DMSP-amine (Kocsis et al., 1998) (Fig. 4). The enzymes have not yet been isolated, but in vivo radiolabeling evidence points to an SMM-specific decarboxylase plus a specific amine oxidase, dehydrogenase or transaminase (Kocsis et al., 1998).

METABOLIC ENGINEERING OF Gly BETAINE SYNTHESIS

Glycine betaine accumulation has long been a target for engineering stress resistance (Le Rudulier et al., 1984). The idea that introducing the Glycine betaine pathway into plants that lack it will enhance their stress tolerance is based both on comparative physiology (Yancey, 1994) and on genetic evidence from a mutation in maize that abolishes Glycine betaine synthesis and reduces salt and heat tolerance (Saneoka et al., 1995; Yang et al., 1996). Wide-crossing work on *Lophopyrum elongatum* and wheat has provided further physiological-genetic evidence that Glycine betaine accumulation contributes to salt tolerance (Colmer et al., 1995). Because plants have been found not to catabolize Glycine betaine (Rhodes and Hanson, 1993; Nuccio et al., 1998), it has been reasonably assumed that engineering the synthesis of Glycine betaine will lead to its accumulation.

Several groups have taken the first step toward this goal by expressing choline-oxidizing enzymes from bacteria (Lilius et al., 1996; Hayashi et al., 1997; Alia et al., 1998; Sakamoto et al., 1998) or spinach CMO (Nuccio et al., 1998) in tobacco and other plants that do not contain Glycine betaine ("nonaccumulators"). The transgenic plants produced a little Glycine betaine and, in some cases, showed small but significant increases in tolerance to various stresses (for review, see Nuccio et al., 1999). These results are encouraging inasmuch as they confirm that Glycine betaine synthesis can be engineered. However, the Glycine betaine levels obtained to date in transgenic plants (typically about 0.1–1 $\mu\text{mol g}^{-1}$ fresh weight) are only a small percentage of those in spinach, sugar beet, and other plants that are natural Glycine betaine accumulators. The main constraint on Glycine betaine production in transgenic plants appears to be the endogenous choline supply, because providing choline exogenously leads to a massive increase in Glycine betaine synthesis (Nuccio et al., 1998). It will therefore most likely be necessary to up-regulate the de novo synthesis of choline in order to increase Glycine betaine synthesis in nonaccumulators expressing foreign choline-oxidizing enzymes (Nuccio et al., 1998).

CONCLUSIONS AND PROSPECTS

Characterization and cloning of the enzymes of Glycine betaine synthesis has enabled us to start using transgenic plants to understand the role of Glycine betaine in stress adaptation. This in turn is helping to define the potential of osmoprotectant engineering in crop improvement. Now that pathways to DMSP have been established, a similar approach could be followed for this compound; the same is true for the other osmoprotectants discussed above. There are two reasons to do this. First, some osmoprotectants may be better than Glycine betaine in certain environments, which has important implications for engineering crops. We have already indicated that choline-O-sulfate could be particularly suitable in high-sulfate conditions because its synthesis can detoxify the sulfate anion. DMSP is another example; since it does not require nitrogen to produce, it may be a better choice than Glycine betaine for environments that are poor in nitrogen. The second reason to transgenically express enzymes that produce various osmoprotectants is to explore the in vivo control of metabolism, about which we currently know very little. Introducing novel pathways increases the demand for precursors, and quantitative analysis of how this impacts metabolic fluxes, pool sizes, and gene expression can be highly informative in regard to the control architecture of metabolic networks (Bailey, 1991). For instance, because betaines and related compounds require very large amounts of methyl groups relative to primary metabolism, engineering betaine overproduction has the potential to teach us much about how the supply of one-carbon units is regulated.

Received March 5, 1999; accepted April 19, 1999.

LITERATURE CITED

- Alia, Hayashi H, Sakamoto A, Murata N (1998) Enhancement of the tolerance of *Arabidopsis* to high temperatures by genetic engineering of the synthesis of glycinebetaine. *Plant J* **16**: 155–161
- Bailey JE (1991) Toward a science of metabolic engineering. *Science* **252**: 1668–1675
- Blunden G, Gordon SM (1986) Betaines and their sulphonio analogues in marine algae. *Prog Phycol Res* **4**: 39–80
- Bourgis F, Roje S, Nuccio ML, Fisher DB, Tarczynski MC, Li C, Herschbach C, Rennenberg H, Pimenta MJ, Shen T-L, Gage D, Hanson AD (1999) S-Methylmethionine has a major role in phloem sulfur transport, and is synthesized by a novel type of methyltransferase. *Plant Cell* (in press)
- Burnet M, Lafontaine PJ, Hanson AD (1995) Assay, purification, and partial characterization of choline monooxygenase from spinach. *Plant Physiol* **108**: 581–588
- Carpenter JF, Crowe JH (1988) The mechanism of cryoprotection of proteins by solutes. *Cryobiology* **25**: 244–255
- Colmer TD, Epstein E, Dvorak J (1995) Differential solute regulation in leaf blades of various ages in salt-sensitive wheat and a salt-tolerant wheat \times *Lophopyrum elongatum* (Host) A. Love amphiploid. *Plant Physiol* **108**: 1715–1724
- Delauney AJ, Verma DPS (1993) Proline biosynthesis and osmoregulation in plants. *Plant J* **4**: 215–223
- Essery JM, McCaldin DJ, Marion L (1962) The biogenesis of stachydrine. *Phytochemistry* **1**: 209–213
- Gage DA, Rhodes D, Nolte KD, Hicks WA, Leustek T, Cooper AJL, Hanson AD (1997) A new route for synthesis of dimethylsulphonioacetate in marine algae. *Nature* **387**: 891–894

- Génard H, Le Saos J, Billard J-P, Trémolières A, Boucaud J (1991) Effect of salinity on lipid composition, glycine betaine content and photosynthetic activity in chloroplasts of *Suaeda maritima*. *Plant Physiol Biochem* **29**: 421–427
- Hanson AD, Rathinasabapathi B, Rivoal J, Burnet M, Dillon MO, Gage DA (1994) Osmoprotective compounds in the Plumbaginaceae: a natural experiment in metabolic engineering of stress tolerance. *Proc Natl Acad Sci USA* **91**: 306–310
- Hayashi H, Mustardy L, Deshniun P, Ida M, Murata N (1997) Transformation of *Arabidopsis thaliana* with the *codA* gene for choline oxidase: accumulation of glycinebetaine and enhanced tolerance to salt and cold stress. *Plant J* **12**: 133–142
- Holmstrom KO, Welin B, Mandal A, Kristiansdottir I, Teeri TH, Lamark T, Strom AR, Palva ET (1994) Production of the *Escherichia coli* betaine-aldehyde dehydrogenase, an enzyme required for the synthesis of the osmoprotectant glycine betaine, in transgenic plants. *Plant J* **6**: 749–758
- Kocsis MG, Nolte KD, Rhodes D, Shen T-L, Gage DA, Hanson AD (1998) Dimethylsulfoniopropionate biosynthesis in *Spartina alterniflora*: evidence that S-methylmethionine and dimethylsulfoniopropylamine are intermediates. *Plant Physiol* **117**: 273–281
- Le Rudulier D, Strom AR, Dandekar AM, Smith LT, Valentine RC (1984) Molecular biology of osmoregulation. *Science* **224**: 1064–1068
- Lilius G, Holmberg N, Bülow L (1996) Enhanced NaCl stress tolerance in transgenic tobacco expressing bacterial choline dehydrogenase. *Bio-Technology* **14**: 177–180
- Malin G, Kirst GO (1997) Algal production of dimethyl sulphide and its atmospheric role. *J Phycol* **33**: 889–896
- Matoh T, Watanabe J, Takahashi E (1987) Sodium, potassium, chloride, and betaine concentrations in isolated vacuoles from salt-grown *Atriplex gmelini* leaves. *Plant Physiol* **84**: 173–177
- Naidu BP, Jones GP, Paleg LG, Poljakoff-Mayber A (1987) Proline analogues in *Melaleuca* species: response of *Melaleuca lanceolata* and *M. uncinata* to water stress and salinity. *Aust J Plant Physiol* **14**: 669–677
- Nakamura T, Yokota S, Muramoto Y, Oguri Y, Fukui K, Takabe T (1997) Expression of a betaine aldehyde dehydrogenase gene in rice, a glycinebetaine nonaccumulator, and possible localization of its protein in peroxisomes. *Plant J* **11**: 1115–1120
- Nolte KD, Hanson AD, Gage DA (1997) Proline accumulation and methylation to proline betaine in *Citrus*: implications for genetic engineering of stress resistance. *J Am Soc Hortic Sci* **122**: 8–13
- Nuccio ML, Rhodes D, McNeil SD, Hanson AD (1999) Metabolic engineering of plants for osmotic stress resistance. *Curr Opin Plant Biol* (in press)
- Nuccio ML, Russell BL, Nolte KD, Rathinasabapathi B, Gage DA, Hanson AD (1998) The endogenous choline supply limits glycine betaine synthesis in transgenic tobacco expressing choline monooxygenase. *Plant J* **16**: 487–498
- Paquet L, Rathinasabapathi B, Saini H, Zamir L, Gage DA, Huang Z-H, Hanson AD (1994) Accumulation of the compatible solute 3-dimethylsulfoniopropionate in sugarcane and its relatives, but not other gramineous crops. *Aust J Plant Physiol* **21**: 37–48
- Rathinasabapathi B, Burnet M, Russell BL, Gage DA, Liao P-C, Nye GJ, Scott P, Golbeck JH, Hanson AD (1997) Choline monooxygenase, an unusual iron-sulfur enzyme catalyzing the first step of glycine betaine synthesis in plants: prosthetic group characterization and cDNA cloning. *Proc Natl Acad Sci USA* **94**: 3454–3458
- Rathinasabapathi B, McCue KF, Gage DA, Hanson AD (1994) Metabolic engineering of glycine betaine synthesis: plant betaine aldehyde dehydrogenases lacking typical transit peptides are targeted to tobacco chloroplasts where they confer betaine aldehyde resistance. *Planta* **193**: 155–162
- Rhodes D, Gage DA, Cooper AJL, Hanson AD (1997) S-Methylmethionine conversion to dimethylsulfoniopropionate: evidence for an unusual transamination reaction. *Plant Physiol* **115**: 1541–1548
- Rhodes D, Hanson AD (1993) Quaternary ammonium and tertiary sulfonium compounds in higher plants. *Annu Rev Plant Physiol Plant Mol Biol* **44**: 357–384
- Rivoal J, Hanson AD (1994) Choline-O-sulfate biosynthesis in plants: identification and partial characterization of a salinity-inducible choline sulfotransferase from species of *Limonium* (Plumbaginaceae). *Plant Physiol* **106**: 1187–1193
- Robinson SP, Jones GP (1986) Accumulation of glycinebetaine in chloroplasts provides osmotic adjustment during salt stress. *Aust J Plant Physiol* **13**: 659–668
- Russell BL, Rathinasabapathi B, Hanson AD (1998) Osmotic stress induces expression of choline monooxygenase in sugar beet and amaranth. *Plant Physiol* **116**: 859–865
- Sakamoto A, Alia, Murata N (1998) Metabolic engineering of rice leading to biosynthesis of glycinebetaine and tolerance to salt and cold. *Plant Mol Biol* **38**: 1011–1019
- Saneoka H, Nagasaka C, Hahn DT, Yang W-J, Premachandra GS, Joly RJ, Rhodes D (1995) Salt tolerance of glycinebetaine-deficient and -containing maize lines. *Plant Physiol* **107**: 631–638
- Stoop JMH, Williamson JD, Pharr DM (1996) Mannitol metabolism in plants: a method for coping with stress. *Trends Plant Sci* **1**: 139–144
- Summers PS, Nolte KD, Cooper AJL, Borgeas H, Leustek T, Rhodes D, Hanson AD (1998) Identification and stereospecificity of the first three enzymes of 3-dimethylsulfoniopropionate biosynthesis in a chlorophyte alga. *Plant Physiol* **116**: 369–378
- Timasheff SN (1992) A physicochemical basis for selection of osmolytes by nature. In GN Somero, CB Osmond, CL Bolis, eds, *Water and Life*. Springer-Verlag, Berlin, pp 70–84
- Trossat C, Nolte KD, Hanson AD (1996) Evidence that the pathway of dimethylsulfoniopropionate biosynthesis begins in the cytosol and ends in the chloroplast. *Plant Physiol* **111**: 965–973
- Trossat C, Rathinasabapathi B, Hanson AD (1997) Transgenically expressed betaine aldehyde dehydrogenase efficiently catalyzes oxidation of dimethylsulfoniopropionaldehyde and ω -aminoaldehydes. *Plant Physiol* **113**: 1457–1461
- Trossat C, Rathinasabapathi B, Weretilnyk EA, Shen T-L, Huang Z-H, Gage DA, Hanson AD (1998) Salinity promotes accumulation of 3-dimethylsulfoniopropionate and its precursor S-methylmethionine in chloroplasts. *Plant Physiol* **116**: 165–171
- Winzor CL, Winzor DJ, Paleg LG, Jones GP, Naidu BP (1992) Rationalization of the effects of compatible solutes on protein stability in terms of thermodynamic nonideality. *Arch Biochem Biophys* **296**: 102–107
- Yancey PH (1994) Compatible and counteracting solutes. In K Strange, ed, *Cellular and Molecular Physiology of Cell Volume Regulation*. CRC Press, Boca Raton, FL, pp 81–109
- Yang G, Rhodes D, Joly RJ (1996) Effects of high temperature on membrane stability and chlorophyll fluorescence in glycinebetaine-deficient and glycinebetaine-containing maize lines. *Aust J Plant Physiol* **23**: 437–443