

# Inappropriate secretion of antidiuretic hormone in infants with respiratory infections

RODNEY P A RIVERS, MARY L FORSLING, AND RICHARD P OLVER

*Department of Paediatrics, University College Hospital Medical School, and Department of Physiology, Middlesex Hospital Medical School*

**SUMMARY** Four infants in whom excessive secretion of antidiuretic hormone was associated with pulmonary infections are reported. Severe hyponatraemia was noted in 3 of them; in the fourth, fluid restriction may have prevented this complication.

In several reports associations have been noted between the excessive secretion of antidiuretic hormone (ADH) and disorders encountered in the neonatal period and in later childhood. Those in which pulmonary disorders have been implicated include viral and bacterial pneumonias,<sup>1-5</sup> pneumothorax and atelectasis,<sup>6</sup> acute asthmatic attacks,<sup>7,8</sup> and lung distension during continuous positive airways pressure (CPAP) administration and mechanical ventilation<sup>9</sup> (although some of the neonates in the last-mentioned series also had intracranial disease).

This report draws attention to the consequences of excessive secretion of ADH in association with common pulmonary infections in infancy. The severity and rapidity of development of hyponatraemia that resulted in 3 of the cases is noteworthy. In the fourth case, prospective therapeutic measures may have prevented the development of this complication. Possible mechanisms for increased ADH secretion are discussed and therapeutic approaches considered.

## Patients and methods

Four infants were admitted during a local outbreak of respiratory syncytial virus (RSV)—positive bronchiolitis and pneumonia. All manifested evidence of increased secretion of ADH.

RSV was identified by immunofluorescence and by culture of material obtained by aspiration of pharyngeal washes.<sup>10,11</sup>

ADH was assayed in plasma by radioimmune assay.<sup>12,13</sup> Urine osmolality was measured by freezing point depression, specific gravity by T S refractometer (American Optical), and biochemical

estimations by routine laboratory procedures. Blood gases were analysed in heparinised arterial blood using Radiometer equipment.

## Case reports

**Case 1.** A 6-week-old formula-fed girl, presented with a 4-day history of increasing respiratory difficulty, cough, partial feed refusal, and sleeplessness. She had vomited twice 2 days before admission; there had been no diarrhoea. On the day of admission her parents noted that she felt stiff when handled and earlier that day she had had a grand mal seizure lasting about 30 minutes.

## Examination

Grunting, dyspnoeic, with increased respiratory rate; expiratory rhonchi and fine crepitations were present over both lung fields. She was not apparently dehydrated. A focal left-sided seizure began shortly after admission. The ensuing apnoea necessitated endotracheal intubation and ventilation for 30 minutes.

## Investigations

Normal plasma urea, sugar, calcium, magnesium, cerebrospinal fluid, electrocardiogram, and skull radiograph. Chest radiograph showed translucent hyperinflated lungs. Plasma sodium 121 mmol/l, urine osmolality 275 mmol/kg. Blood gas analysis: PaO<sub>2</sub> 5.1 kPa (38 mmHg), and Paco<sub>2</sub> 6.8 kPa (45 mmHg) breathing air spontaneously before intubation. Immunofluorescence for RSV was positive.

**Course.** Treated with digoxin, anticonvulsants, and oxygen and, initially, with antibiotics.

A rise in plasma sodium concentration to 134 mmol/l was achieved in 12 hours in association with improved oxygenation, fluid restriction (100 ml/kg in 24 hours), and administration of hypertonic (270 mmol/l) sodium chloride followed by 0.9% sodium chloride. Urine sodium excretion occurred concurrently and equalled 11 mmol of the 57 mmol infused in 24 hours.  $Paco_2$  was normal within 48 hours and oxygen supplementation was stopped after 72 hours. The infant subsequently was shown to have a normal electroencephalogram; a synacthen stimulation test induced normal cortisol release.

**Case 2.** A 7-week-old breast-fed boy presented with history of nasal obstruction for 4 days. He had become irritable in association with rapid breathing for 2 hours before admission. There was no diarrhoea.

**Examination**

Tachycardia 156/minute, grade 2/6 apical systolic murmur; respirations 36/minute, no added sounds in chest. Nasal secretions increased. Liver and spleen were not palpable. After 36 hours the respiratory rate had increased to 130 per minute, and the liver and spleen were palpable.

**Investigations**

Chest radiograph showed translucent markedly hyperinflated lungs with depressed diaphragm.

**Treatment.** Digoxin, a single dose of frusemide 1 mg/kg, fluid restriction (100 ml/kg in 24 hours), and oxygen.

**Course.** Forty-eight hours after admission diffuse fine crepitations were audible over both lung fields. Plasma sodium was normal (135 mmol/l) at this time. Twelve hours later plasma sodium had fallen to 124 mmol/l in association with an inappropriately concentrated urine (534 mmol/kg) (Table 1). Pulmonary gas exchange had deteriorated,  $PaO_2$  6.4 kPa (48 mmHg) in  $F_{iO_2}$  0.21,  $Paco_2$  9.1 kPa (68 mmHg). Further treatment consisted of greater

fluid restriction (50 ml/kg in 24 hours), ampicillin, and continued oxygen supplementation. The course of the illness is shown in Fig. 1. During the next 4 days brief periods of increased fluid intake were associated with falls in plasma sodium concentration, and progressive hypertension was noted. Resolution of electrolyte disturbance took place after 5 days, with the fall in urine concentration occurring during continued fluid restriction. Immunofluorescence for RSV on day 3 of the respiratory illness was negative.




Fig. 1 (Case 2.) Flow diagram of clinical course. Hyponatraemia developed on day 4 of respiratory illness at the peak of recorded hypercarbia in association with production of a non-maximally dilute urine, inappropriately high antidiuretic hormone level, and natriuresis. Fluid restriction was associated with a rise in plasma sodium, urea, and blood pressure. The diminution in urine concentration on day 8 occurred during continued fluid restriction and represents excretion of body water derived from body tissue catabolism and fluid intake, previously retained under the influence of excessive antidiuretic hormone secretion.

Table 1 Results of investigations in Cases 1-4

Case	Plasma		Urine		Body weight (kg)		ADH (pmol/l) (0.96-3.60)	
	Na (mmol/l)	Urea (mmol/l)	Na (mmol/l)	Osmolality (mmol/kg)	At diagnosis of hyponatraemia	On recovery from hyponatraemia	During respiratory distress	On recovery
1	121	2.5	34	275	4.075	3.800	—	—
2	124	3.5	53	534	4.900	4.800	6.24	2.40
3	114	4.2	3	243	2.980	2.960	—	—
4	137	2.5	28	474	6.400	6.500	7.20	1.68

Lowest values of plasma sodium concentrations are shown; these occurred within 4 days of onset of pulmonary symptoms. Urine osmolalities are inappropriately high in all cases. There is no evidence of dehydration either from plasma urea values or on the basis of body-weight measurements made at the time of hyponatraemia compared with those recorded on recovery.

**Case 3.** A girl, the second of twins born at 32 weeks' gestation weighing 1.69 kg, had developed transient tachypnoea in the neonatal period. At age 8 weeks she became unwell with feed refusal and nasal obstruction, and within 24 hours developed rapid respiration of 60/minute and a tachycardia of 190/minute; there were fine crepitations over both lung fields. Chest radiograph showed right upper lobe consolidation. Within 48 hours of onset, pulmonary function deteriorated, PaO<sub>2</sub> 6.0 kPa (45 mmHg), F<sub>I</sub>O<sub>2</sub> 0.30, PaCO<sub>2</sub> 6.7 kPa (50 mmHg). She began convulsing.

Further investigations showed normal plasma calcium, sugar, urea, and cerebrospinal fluid, but marked hyponatraemia (114 mmol/l) and less than maximally dilute urine (243 mmol/kg) (Table 1).

**Treatment.** She was ventilated for 24 hours, fluids were restricted (42 ml/kg in 24 hours), and she was given sodium supplements (7 mmol/kg per day) and antibiotics. An immediate improvement was obtained. Plasma sodium concentration rose to 130 mmol/l within 24 hours and urine osmolality fell to 195 mmol/kg during this period of fluid restriction. RSV was cultured from pharyngeal aspirate.

**Case 4.** A 6-month-old formula-fed boy presented with a 24-hour history of cough, blocked nose, and breathlessness.

#### Examination

Tachypnoea (80/minute), fine crepitations and expiratory rhonchi were present over both lung fields. Liver and spleen were palpable.

#### Investigations

Normal arterial blood gases, electrolytes, and urea. Chest radiograph showed translucent hyperinflated lungs. Immunofluorescence for RSV was positive.

**Treatment.** Fluid restriction (100 ml/kg in 24 hours), ampicillin initially.

**Course.** Increasing dyspnoea over first 24 hours. Plasma sodium ranged from 137 to 139 mmol/l during 5 days of respiratory symptoms.

During the period of maximum respiratory difficulty both the plasma ADH level (7.20 pmol/l), and urine osmolality (474 mmol/kg) were increased. Within 24 hours the urine osmolality had fallen (164 mmol/kg) and remained at between 126 and 164 mmol/kg with the infant receiving 100/ml per kg during 4 days of improving respiratory status (Table 1). By this time the ADH level had fallen to

1.68 pmol/l. Body weight did not vary beyond the range 6400–6600 g during the first 7 days after admission.

#### Discussion

The physiology of ADH was reviewed by Hays.<sup>14</sup> ADH is believed to be released into the plasma from its site of storage in the posterior pituitary gland in response to the following stimuli: an increase in extracellular fluid osmolality,<sup>15 16</sup> reduction in intravascular fluid volume<sup>17</sup> as reflected by a reduction in tension in the wall of the left atrium,<sup>18 19</sup> and stimulation of pressor receptors in the carotid sinus.<sup>20</sup> Beta sympathomimetic drugs and stress itself also stimulate ADH release.<sup>21 22</sup> Acting via adenylate cyclase and cyclic-AMP, ADH renders the distal renal tubules and collecting ducts permeable to water, urea, and other solutes.<sup>23 24</sup> Reabsorption at these sites depends on the existing concentration gradients from tubule to interstitium.

From these considerations, it is evident that should a local change, which normally represents a physiological stimulus for ADH release, fail to be a true reflection of the overall osmotic or circulatory status of an individual, ADH release may occur with undesirable consequences. Then the release may be excessive for the body's requirements. If release occurs in the absence of any physiological stimulus, it may be regarded as inappropriate. The consequences of these two situations may be similar with the development of a hyponatraemic, hypo-osmolar state as first described in association with ADH-secreting carcinomata of the bronchus.<sup>25</sup> That syndrome of inappropriate ADH secretion (SIADH) is manifested by hyponatraemia in combination with the excretion of a urine which is not maximally dilute in the absence of dehydration, blood volume depletion, or abnormal adrenal function.<sup>26</sup> Continued natriuresis may be observed.

In all 3 infants with hyponatraemia, the low plasma sodium level was related to severe pulmonary dysfunction as shown by hypoxaemia and hypercarbia. Neither on clinical evidence nor on the basis of biochemical and sequential body weight measurements was dehydration present. The clinical history of Case 1, and measured fluid intakes of Cases 2 and 3, indicated that there had been no excessive fluid intake during the 24 hours preceding the detection of hyponatraemia, neither had there been any apparent abnormal fluid losses. In Cases 1 and 3, who had been on free fluids, intake was below normal because of feed refusal. It is noteworthy that Case 2 became hyponatraemic despite fluid restriction (100 ml/kg per day) whereas this degree of restriction was

sufficient to prevent hyponatraemia in Case 4. In Cases 2 and 4, the level of ADH was found to be increased in the acute phase (Table 1).

In Cases 1, 2, and 3 the urine solute concentrations were inappropriately high for the degree of plasma hypotonicity present (Table 1). During the period of maximal hyponatraemia, measured total cation excretion was low in all cases. Whereas in Cases 1 and 2, small volumes of urine with high concentrations of sodium and potassium were excreted, in Case 3 (the most severely hyponatraemic) urine sodium concentration was low. Aldosterone secretion has been shown to continue once a new steady state is reached in SIADH so that sodium excretion reflects intake.<sup>27</sup> Absence of continued natriuresis in the presence of sodium deprivation in SIADH has been noted previously.<sup>28</sup>

Despite the high degree of correlation between serum sodium concentration and the ratio of the sum of the total body exchangeable sodium and potassium to total body water in normal individuals, it has not been possible to account for the degree of hyponatraemia induced by acute water loading on the basis of urinary cation (Na and K) loss and body water gain in cases where ADH is being excessively secreted.<sup>28-30</sup> Attention has also been drawn<sup>28</sup> to the lower than predicted rise in plasma sodium after sodium administration in SIADH. Explanations have invoked the concept of an altered body sodium and potassium distribution, and a low total body exchangeable potassium with increased muscle sodium content has been reported.<sup>30-32</sup>

Of interest in this regard are the observations on sodium and potassium balance in Case 1. During the hyperosmolar sodium chloride administration, balance studies indicated that between 12 and 19 mmol sodium remained unaccounted for (Table 2),

Table 2 (Case 1.) 24-hour sodium and potassium balance

	Input (I) IV (mmol)	Urine (U) (mmol)	Rise in plasma concentration (mmol/l)
Na (mmol/l)	57	11	15
K	0	6	0
Assumed ECF volume as % body weight:		35	40
Calculated increase in ECF sodium (mmol):		21	24
Cation balance $Na_I + K_I -$ $(U_{Na} + U_K + ECF_{Na})$ mmol:		+19	+16
			+12

The measured rise in plasma Na was used to calculate the increase in ECF Na for three estimations of ECF volume<sup>43</sup> (assuming that the volume might be increased due to excessive ADH secretion). The cation balance gives the mmol Na 'unaccounted for' on the basis of the balance calculations.  $Na_I$ : sodium input;  $K_I$ : potassium input;  $U_{Na}$ : urine sodium;  $U_K$ : urine potassium;  $ECF_{Na}$ : extracellular fluid sodium; there was no potassium input over the balance period.

the degree of the discrepancy depending on the infant's extracellular fluid volume for which assumed values were used in the calculations. No stool was passed during this 24-hour balance period, neither was there evidence of gastrointestinal distension nor evidence of overt sweating. It seems probable therefore, that there took place a redistribution of the sodium administered to this infant from extracellular to intracellular compartments in exchange for potassium. In support of this view is the observation that Cases 2 and 3 were both in negative potassium balance and became hypokalaemic with continued urinary potassium loss. Continued natriuresis, which is a common feature of SIADH, was undoubtedly an additional factor responsible for the partial failure to restore plasma sodium to normal in Cases 1 and 2.

**Pathogenesis of excessive ADH secretion.** As a consequence of the pulmonary disease in these infants, several stimuli (Fig. 2) for excessive secretion of ADH may be present. It seems likely that the most important of these is a fall in left atrial tension, a potent cause of ADH secretion.<sup>18,19</sup> This might occur in situations where there is left atrial compression by markedly hyperinflated lungs, as in bronchiolitis and acute asthma, or when the fall in tension is a consequence of diminished left atrial filling secondary to a raised pulmonary vascular resistance<sup>4</sup> as might occur in the above conditions or during the administration of CPAP or positive pressure ventilation.




Fig. 2 Possible mechanisms of origin of 'excessive' antidiuretic hormone release in bronchiolitis and pneumonia.

Such a reduction in left atrial tension, although being a consequence of regional factors, mimics the effect of systemic hypovolaemia by inducing ADH secretion.<sup>33</sup>

Also relevant is the observation that hypoxaemia causes release of ADH by direct hypothalamic stimulation in the fetal lamb<sup>34</sup> and in anaesthetised dogs undergoing a water diuresis<sup>35</sup> although, in addition, ADH release may be mediated via arterial baroreceptors.<sup>35</sup>

Whether ADH is actually secreted from virally infected lung tissue remains to be established. Tuberculous lung parenchyma has been shown to contain ADH, but it is not clear whether the ADH-like material is produced by the infected tissue or has been absorbed on to it from the plasma.<sup>36</sup>

In Cases 1 and 3 the duration of excessive secretion of ADH was brief. The initial, inappropriately high, urine concentration became normal during fluid restriction. In each case the improvement was associated with a rise in arterial oxygen tension although radiological evidence of pulmonary hyperinflation persisted for some days.

The clinical course of Case 2 was different. He maintained a high urine concentration for 5 days during which time small increases in fluid intake resulted in hyponatraemia. Throughout this period he was in positive sodium balance excluding possible losses in sweat (Fig. 1). On the seventh day of fluid restriction the urine specific gravity suddenly fell. It seems probable that there was excessive ADH secretion throughout this period causing water retention. The fact that the body weight remained stable despite increased metabolic needs (pyrexia, tachycardia, tachypnoea) while the baby was receiving a low calorie intake enforced by the fluid restriction, supports this view as does the observation that the body weight on recovery was some 250 g below the admission weight. Catabolism of this amount of body tissue to CO<sub>2</sub> and water would have made a significant contribution to the total water 'load' in this baby. Continuing hypoxaemia was not the explanation for this infant's apparently prolonged excessive secretion of ADH. However, the degree of pulmonary hyperinflation in this infant was striking and this persisted throughout the period of presumed ADH secretion.

High plasma renin levels have been recorded during periods of fluid and sodium deprivation in SIADH<sup>37</sup> and in association with sodium deprivation in normal volunteers.<sup>38</sup> Progressive hypertension developed in Case 2 during the phase of severe fluid restriction, and activation of the renin-angiotension system might therefore have been the cause.<sup>39</sup> At no time was there clinical or biochemical

evidence (in the form of a metabolic acidosis) of reduced organ perfusion during the period of fluid restriction and increasing hypertension.

It is probable that rapidly developing hyponatraemia was the cause of the seizures in Cases 1 and 3. Indeed the rapidity with which hyponatraemia can develop is well documented by the course of Case 2. In rats, hyponatraemia increases brain epileptogenicity.<sup>40</sup>

## Conclusions

We suggest that routine sequential estimations be made of sodium and potassium concentrations and of osmolalities on plasma and urine samples during periods of deteriorating pulmonary function in infants with pneumonia or bronchiolitis. Avoidance of hypoxaemia may reduce the incidence of excessive ADH secretion; the value of instituting moderate fluid restriction with continued administration of sodium and potassium requirements remains to be investigated.

Sodium chloride administration, as a hypertonic solution, has been shown to be effective in achieving a rapid rise in plasma sodium in the short term despite associated high urine sodium concentrations, although in chronic inappropriate ADH secretion, fluid restriction with or without treatment with drugs<sup>41</sup> remains the basis of treatment. Administration of frusemide may also be effective in treating the symptoms of acute hyponatraemia.<sup>42</sup> The more aggressive forms of treatment are justified in the presence of marked hyponatraemia whether or not associated with convulsions. The development of hypertension may be a reflection of excessive fluid and sodium restriction.

Correction of hypoxaemia does not necessarily remove the stimulus for the inappropriate ADH secretion in cases of bronchiolitis with marked pulmonary hyperinflation.

We thank Dr M F Friedman and Dr J Luder for letting us have data on their patients, and Professor C J Dickinson for his help.

## References

- 1 Darrow D C, Hartmann A F. Chemical changes occurring in the body as a result of certain diseases. IV. Primary pneumonia in children. *Am J Dis Child* 1929; 37: 323-34.
- 2 Rutstein D D, Thomson K J, Tolmach D M, Walker W H, Flood R J. Plasma volume and 'extravascular thiocyanate space' in pneumococcus pneumonia. *J Clin Invest* 1945; 24: 11-20.
- 3 Mor J, Ben-Galim E, Abrahamov A. Inappropriate antidiuretic hormone secretion in an infant with severe pneumonia. *Am J Dis Child* 1975; 129: 133-5.

- 4 Cohen L F, di Sant'Agnes P A, Taylor A, Gill J R. The syndrome of inappropriate antidiuretic hormone secretion as a cause of hyponatremia in cystic fibrosis. *J Pediatr* 1977; **90**: 574-8.
- 5 Le Moing G, Gendrell D, Couvreur J, Gaultier Cl, Tournier G, Gerbeaux J. Sécrétion inappropriée d'hormone antidiurétique au cours des infections respiratoires aiguës à adénovirus chez l'enfant. *Nouv Presse Med* 1977; **6**: 3825-33.
- 6 Paxson C L, Jr, Stoerner J W, Denson S E, Adcock E W, III, Morriss F H, Jr. Syndrome of inappropriate antidiuretic hormone secretion in neonates with pneumothorax or atelectasis. *J Pediatr* 1977; **91**: 459-63.
- 7 Higer R W, Holliday M A. Acute bronchial asthma and possible inappropriate secretion of antidiuretic hormone (abstract). *Pediatr Res* 1973; **7**: 428.
- 8 Baker J W, Yerger S, Segar W E. Elevated plasma antidiuretic hormone levels in status asthmaticus. *Mayo Clin Proc* 1976; **51**: 31-4.
- 9 Pomarède R, Moriette G, Czernichow P, Relier J-P. Étude de la vasopressine plasmatique chez les enfants prématures soumis à las ventilation artificielle. *Arch Fr Pediatr* 1978; **35**: Supplement 2, 75-83.
- 10 McQuillin J, Gardner P S. Rapid diagnosis of respiratory syncytial virus infection by immunofluorescent antibody techniques. *Br Med J* 1968; **i**: 602-5.
- 11 Sturdy P M, McQuillin J, Gardner P S. A comparative study of methods for the diagnosis of respiratory virus infections in childhood. *J Hyg (Camb)* 1969; **67**: 659-70.
- 12 Karonen S-L, Morsky P, Siren M, Seuderling U. An enzymatic solid-phase method for trace iodination of proteins and peptides with 125-iodine. *Anal Biochem* 1975; **67**: 1-10.
- 13 Chard T, Forsling M L. Bioassay and radioimmunoassay of oxytocin and vasopressin. In: Antoniades H N, ed. *Hormones in human blood: detection and assay*. Cambridge, Mass: Harvard University Press, 1976; 488-516.
- 14 Hays R M. Antidiuretic hormone. *N Engl J Med* 1976; **295**: 659-65.
- 15 Verney E B. The antidiuretic hormone and the factors which determine its release. *Proc R Soc Lond [Biol]* 1947; **135**: 25-106.
- 16 Haberich F J. Osmoreceptors in the portal circulation and their significance for the regulation of water balance. *Triangle* 1971; **10**: 123-30.
- 17 Davies R, Forsling M L, Slater J D H. The inter-relationship between the release of renin and vasopressin as defined by orthostasis and propranolol. *J Clin Invest* 1977; **60**: 1438-41.
- 18 Gauer O H, Henry J P, Sieker H O. Cardiac receptors and fluid volume control. *Prog Cardiovasc Dis* 1961; **4**: 1-26.
- 19 Goetz K L, Bond G C, Hermreck A S, Trank J W. Plasma ADH levels following a decrease in mean atrial transmural pressure in dogs. *Am J Physiol* 1970; **219**: 1424-8.
- 20 Share L. Vasopressin, its bioassay, and the physiological control of its release. *Am J Med* 1967; **42**: 701-12.
- 21 Schrier R W, Lieberman R, Ufferman R C. Mechanism of antidiuretic effect of beta adrenergic stimulation. *J Clin Invest* 1972; **51**: 97-111.
- 22 Berl T, Cadnapaphornchai P, Harbottle J A, Schrier R W. Mechanism of stimulation of vasopressin release during beta-adrenergic stimulation with isoproterenol. *J Clin Invest* 1974; **53**: 857-67.
- 23 Orloff J, Handler J S. The cellular mode of action of antidiuretic hormone. *Am J Med* 1964; **36**: 686-97.
- 24 Jamison R L, Maffly R H. The urinary concentrating mechanism. *N Engl J Med* 1976; **295**: 1059-67.
- 25 Schwartz W B, Bennett W, Curelop S, Bartter F C. A syndrome of renal sodium loss and hyponatremia probably resulting from inappropriate secretion of antidiuretic hormone. *Am J Med* 1957; **23**: 529-42.
- 26 Bartter F C, Schwartz W B. The syndrome of inappropriate secretion of antidiuretic hormone. *Am J Med* 1967; **42**: 790-806.
- 27 Fichman M P, Bethune J E. The role of adrenocorticoids in the inappropriate antidiuretic hormone syndrome. *Ann Int Med* 1968; **68**: 806-20.
- 28 Nolph K D, Schrier R W. Sodium, potassium, and water metabolism in the syndrome of inappropriate antidiuretic hormone secretion. *Am J Med* 1970; **49**: 534-45.
- 29 Gonzalez C F, Finberg L, Bluestein D D. Electrolyte concentration during acute infections. *Am J Dis Child* 1964; **107**: 476-82.
- 30 Edelman I S, Leibman J, O'Meara M P, Birkenfeld L W. Interrelations between serum sodium concentration, serum osmolality and total exchangeable sodium, total exchangeable potassium, and total body water. *J Clin Invest* 1958; **37**: 1236-56.
- 31 Jones N F, Barraclough M A, Forsling M L, Petch C P. Inappropriate production of vasopressin, potassium deficiency, and cerebrovascular disease. *Am J Med* 1968; **45**: 474-9.
- 32 Kaye M. An investigation into the cause of hyponatremia in the syndrome of inappropriate secretion of antidiuretic hormone. *Am J Med* 1966; **41**: 910-26.
- 33 Friedman A L, Segar W E. Antidiuretic hormone excess. *J Pediatr* 1979; **94**: 521-6.
- 34 Alexander D P, Forsling M L, Martin M J, et al. The effect of maternal hypoxia on fetal pituitary hormone release in the sheep. *Biol Neonate* 1972; **21**: 219-28.
- 35 Anderson R J, Pluss R G, Berns A S, et al. Mechanism of effect of hypoxia on renal water excretion. *J Clin Invest* 1978; **62**: 769-77.
- 36 Vorherr H, Massry S G, Fallet R, Kaplan L, Kleeman C R. Antidiuretic principle in tuberculous lung tissue of a patient with pulmonary tuberculosis and hyponatremia. *Ann Int Med* 1970; **72**: 383-7.
- 37 Skowsky W R, Fisher D A. Intermittent, idiopathic, inappropriate vasopressin secretion in a child. *J Pediatr* 1973; **83**: 62-8.
- 38 Brennan L A, Malvin R L. Concentrations of antidiuretic hormone in plasma during human sodium restriction. *Nephron* 1977; **19**: 284-7.
- 39 Skeggs L T, Dorer F E, Kahn J R, Lentz K E, Levine M. The biochemistry of the renin-angiotensin system and its role in hypertension. *Am J Med* 1976; **60**: 737-48.
- 40 Swinyard E A, Toman J E P, Goodman L S. The effect of cellular hydration on experimental electroshock convulsions. *J Neurophysiol* 1946; **9**: 47-54.
- 41 Cherrill D A, Stote R M, Birge J R, Singer I. Demeclocycline treatment in the syndrome of inappropriate antidiuretic hormone secretion. *Ann Int Med* 1975; **83**: 654-6.
- 42 Hantman D, Rossier B, Zohlman R, Schrier R. Rapid correction of hyponatremia in the syndrome of inappropriate secretion of antidiuretic hormone. *Ann Int Med* 1973; **78**: 870-5.
- 43 Friis-Hansen B. Changes in body water compartments during growth. *Acta Paediatr Scand [Suppl]* 1957; **46**: Supplement 110, 1-68.

Correspondence to Dr R P A Rivers, Department of Paediatrics, St Mary's Hospital Medical School, London W2 1PG.

Received 25 February 1980