

Appendix 1: Scoring criteria for the Sequential Organ-Failure Assessment (SOFA) score*

Variable	Score				
	0	1	2	3	4
Pao ₂ /FIO ₂ , mm Hg	> 400	≤ 400	≤ 300	≤ 200	≤ 100
Platelet count, × 10 ⁶ /L	> 150	≤ 150	≤ 100	≤ 50	≤ 20
Bilirubin level, mg/dL (μmol/L)	< 1.2 (< 20)	1.2–1.9 (20–32)	2.0–5.9 (33–100)	6.0–11.9 (101–203)	> 12 (> 203)
Hypotension†	None	MABP < 70	Dop ≤ 5	Dop > 5 Epi ≤ 0.1 Norepi ≤ 0.1	Dop > 15 Epi > 0.1 Norepi > 0.1
Glasgow Coma score	15	13–14	10–12	6–9	< 6
Creatinine level, mg/dL (μmol/L)	< 1.2 (< 106)	1.2–1.9 (106–168)	2.0–3.4 (169–300)	3.5–4.9 (301–433)	> 5 (> 434)

Note: Pao₂ = partial pressure of arterial oxygen; FIO₂ = fraction of inspired oxygen; MABP = mean arterial blood pressure, in mm Hg;

*Adapted, with permission, from Ferreira FL, Bota DP, Bross A, et al. Serial evaluation of the SOFA score to predict outcome in critically ill patients. *JAMA* 2001;286:1754–8. Copyright © 2001, American Medical Association. All rights reserved.

†Dop (dopamine), epi (epinephrine) and norepi (norepinephrine) doses in μg/kg per min.