

## CASE REPORT

# Polyarticular juvenile idiopathic arthritis treated with methotrexate complicated by the development of non-Hodgkin's lymphoma

A G Cleary, H McDowell, J A Sills

*Arch Dis Child* 2002;**86**:47–49

A 10 year old boy with juvenile idiopathic arthritis is described. He was treated with methotrexate (MTX) for 2 years 8 months, and presented at routine review with hepatosplenomegaly and suspicious bilateral cervical lymphadenopathy, two months after discontinuing therapy. Magnetic resonance scan revealed significant lymphadenopathy around the upper abdominal aorta and coeliac axis. Lymph node biopsy was consistent with non-Hodgkin's lymphoma, similar to that reported in several adults with rheumatic diseases taking low dose MTX therapy. He was successfully treated with a standard cytotoxic chemotherapy regime, but unfortunately his polyarthritis has subsequently flared some months after completion of his treatment.

### CASE REPORT

#### Clinical findings

A 10 year old boy with a four year history of juvenile idiopathic arthritis (JIA) presented at routine review with bilateral non-tender but firm and suspicious cervical lymphadenopathy. He had hepatosplenomegaly. He had been treated with methotrexate (MTX) 7.5 mg per week, for 2 years 8 months; cumulative dose was 1042.5 mg. The MTX had been given by subcutaneous injection for four months prior to its discontinuation because of nausea associated with oral preparations. MTX had been discontinued two months prior to his presentation with lymphadenopathy as his arthritis had been quiescent for one year. He had also intermittently been treated with daily or alternate daily prednisolone throughout the period of his JIA, and with several intra-articular corticosteroid injections. There were no other co-morbid conditions.

He was admitted for urgent investigation. Full blood count, urea and electrolytes, and lactate dehydrogenase were normal. Serology indicated recent Epstein-Barr virus (EBV) infection, with IgG to EBV detected. Cytomegalovirus (CMV) IgG was not detected. Bone marrow aspirate and trephine revealed no evidence of malignant disease, and there were no malignant cells in his cerebrospinal fluid. Abdominal ultrasound confirmed hepatosplenomegaly and the presence of para-aortic nodes. Magnetic resonance scan revealed a diffusely enlarged spleen, 14 cm in length, with normal texture of liver, pancreas, and both kidneys. There was significant lymphadenopathy around the upper abdominal aorta and coeliac axis, with individual nodes measuring 2 cm in diameter. Computed tomography of the chest revealed evidence of bilateral cervical lymphadenopathy, extending to the sternoclavicular joints. There was no evidence of significant mediastinal or axillary lymphadenopathy, and no evidence of parenchymal lung disease.

#### Pathological findings

A tissue diagnosis of non-Hodgkin's lymphoma was obtained. Histopathology of a lymph node biopsy revealed T cell rich B

cell lymphoma. Most of the larger cells were positive for CD79A, CD20Y, and CD45RA (B cell markers), but negative for CD10, CD15, CD30, EBV, and T cell markers (including CD3, CD4, CD5, CD8, CD43, CD43RO), and dendritic reticulum cells (CD21).

#### Treatment

The patient was treated according to a standard treatment protocol (LMB89), which consists of six months of cytotoxic chemotherapy. This comprised vincristine, cyclophosphamide, prednisolone, MTX, doxorubicin, cytarabine, and intrathecal MTX. During this treatment programme his arthritis went into remission. At follow up six months after completion of chemotherapy his lymphoma is in remission, but unfortunately his arthritis has recurred, requiring careful consideration of treatment options.

### DISCUSSION

The literature relating to lymphoma occurring in patients with rheumatoid arthritis treated with weekly low dose MTX has recently been reviewed; at least 50 cases have been reported.<sup>1</sup> To the best of our knowledge, four children with JIA treated with MTX have developed Hodgkin's lymphoma.<sup>2-5</sup> Two of these children had evidence of EBV infection associated with their lymphoproliferative disease.<sup>3,4</sup> Table 1 summarises the characteristics of the cases.

The incidence of non-Hodgkin's lymphoma in the age group 5–9 years for the period 1968–95 was 6.5 per million per year.<sup>6</sup> At present approximately 100 children at our institution with JIA are treated with MTX. Several other paediatric rheumatology units were contacted (Great Ormond Street, Birmingham, Nottingham, Leeds, and Newcastle). Any other cases of lymphoproliferative disease (LPD) occurring during treatment with MTX were sought, as it was acknowledged that since case reports were not the favoured format for most journals, other cases may exist. In fact only one other case was noted; this was in a child who also had Blackfan-Diamond syndrome and therefore was at increased risk of haematological malignancy (Professor Woo, personal communication).

Post-transplant lymphoproliferative disorder (PTLD) in an immunocompromised host following paediatric renal transplantation has been reported. Srivastava *et al* reported PTLD in six of 84 renal transplant recipients in one centre.<sup>7</sup> In all of these patients PTLD was associated with EBV infection (primary in five, reactivation in one). All patients received therapeutic immunosuppression post-transplantation, with prednisolone and various combinations of mycophenolate

**Abbreviations:** Abbreviations: CMV, cytomegalovirus; EBV, Epstein-Barr virus; JIA, juvenile idiopathic arthritis; LPD; lymphoproliferative disease; MTX, methotrexate; PTLD; post-transplant lymphoproliferative disorder

**Table 1** Characteristics of patients with juvenile arthritis developing lymphoma while treated with MTX

| Ref.     | Diagnosis   | Duration of MTX therapy | Other therapy during treatment with MTX | Cell type                      | Spontaneous remission on stopping MTX | Outcome  |
|----------|---|-------------------------|---|--------------------------------|---------------------------------------|--|
| 2        | Systemic JIA Mixed cellularity Hodgkin's lymphoma                 | 16 months               | Prednisolone                            | CD30 CD15 EBV neg              | Yes                                   | Late LPD relapse<br>Autologous stem cell transplantation<br>Remission of LPD and JIA |
| 3        | Systemic JIA Nodular sclerosing Hodgkin's lymphoma                | 33 months*              | Not known                               | CD30 EBV pos                   | Not attempted                         |  |
| 4        | Polyarthritis (RF positive) Nodular sclerosing Hodgkin's lymphoma | 30 months†              | Prednisolone Cyclosporin A              | CD30 CD15 CD20 LMP1 EBV pos    | Not applicable                        | Death (respiratory failure secondary to legionella pneumonia)                        |
| 5        | Polyarthritis (RF negative) Nodular sclerosing Hodgkin's disease  | 21 months‡              | Sulphasalazine                          | EBV negative                   | Partial response                      | Remission of LPD and JIA after 12 months follow up                                   |
| Our case | Polyarthritis Non-Hodgkin's lymphoma                              | 32 months               | Prednisolone Intra-articular steroids   | CD79A CD20 <sup>+</sup> CD45RA | No                                    | Remission of LPD Relapse of polyarthritis  |

\*MTX discontinued for eight weeks during proven EBV infection 21 months prior to presentation with nodular sclerosing Hodgkin's lymphoma.

†MTX discontinued for two months during episode of intercurrent pneumonia.

‡MTX discontinued after 20 months due to disease remission, and recommenced after one year due to relapse.

mofetil, azathioprine, and cyclosporin. Five of the six patients responded completely to a decrease in immunosuppression and antiviral treatment alone. The sixth patient was unusual in that he developed T cell rich Hodgkin's disease and was EBV positive at the time of transplant. He responded to a combination of chemotherapy and CMV hyperimmune globulin.

No cases of LPD have occurred during MTX therapy for JIA in children registered with the United Kingdom Children's Cancer Study Group (D Walker, personal communication). In contrast, 42 cases of transplantation related LPD have been registered. It is important that all cases of malignant disease occurring in patients with JIA treated with immunosuppressants are registered.

The predominant type of lymphoma reported in adult patients with rheumatic disease treated with MTX and steroids is the same as that occurring in the immunosuppressed (for example, post-transplant and AIDS patients), namely large B cell non-Hodgkin's lymphoma with extranodal involvement.<sup>8</sup> Our patient also developed such a disorder, and as such is the first case reported of non-Hodgkin's lymphoma occurring in a patient with juvenile idiopathic arthritis treated with MTX.

The hypothesis that MTX has a role in the aetiology of LPD is supported, at least in part, by the observation of spontaneous remission of lymphoma on cessation of MTX therapy.<sup>9</sup> This has been reported in at least eight adult patients.<sup>1</sup> Resolution of peripheral adenopathy, but not of thoracic or abdominal lymph node and spleen involvement, was noted in one child with JIA on withdrawal of MTX.<sup>5</sup> She subsequently received one of a projected three cycles of combination chemotherapy (MOPP-ABVD) with radiographically documented complete resolution. The role of EBV virus infection in the pathogenesis of LPD in patients treated with MTX is unclear. EBV positive staining was shown in the lymph node biopsy specimens from two of four previously reported patients with JIA who developed lymphoma; similar findings in adult patients have been reported.<sup>10</sup> It may be speculated that MTX interferes with the handling of EBV by T cells, and allows the subsequent proliferation of EBV infected lymphocytes.

MTX is an established and effective treatment for polyarthritis in children.<sup>11</sup> It is not clear how physicians caring for children with polyarthritis should interpret these isolated case reports, as there is no epidemiological or proven causal association between therapeutic immunosuppression and the development of LPD. Another dilemma that has been raised by the subsequent clinical course in our patient has been the relapse of his polyarthritis some months after completing chemotherapy for his non-Hodgkin's lymphoma. His family are understandably very concerned about the potential impact of further immunosuppressant therapy.

Children with polyarthritis are being treated earlier, with larger doses and for longer durations with MTX. Only with data collected prospectively and collaboratively will it be possible to clearly define the risk of LPD in this group. Vigilant clinical observation for the development of lymphoproliferative disorder is recommended as a routine part of the clinical follow up of all children with JIA treated with weekly low dose MTX, especially as routine haematological monitoring is unlikely to alert physicians to the diagnosis.

#### Authors' affiliations

**A G Cleary, H McDowell, J A Sills**, Royal Liverpool Children's Hospital NHS Trust, Eaton Road, Liverpool L12 2AP, UK

Correspondence to: Dr A G Cleary, Royal Liverpool Children's Hospital NHS Trust, Eaton Road, Liverpool L12 2AP, UK; gavin.cleary@talk21.com

Accepted for publication 4 October 2001

## REFERENCES

- 1 **Georgescu L**, Paget SA. Lymphoma in patients with rheumatoid arthritis. *Drug Saf* 1999;**20**:475-87.
- 2 **Padeh S**, Sharon N, Schiby G, *et al*. Hodgkin's lymphoma in systemic onset juvenile rheumatoid arthritis after treatment with low dose methotrexate. *J Rheumatol* 1997;**24**:2035-7.
- 3 **Munro R**, Porter DR, Sturrock RD. Lymphadenopathy in a patient with systemic onset juvenile chronic arthritis. *Ann Rheum Dis* 1998;**57**:513-17.
- 4 **Krugmann J**, Sailer-Hock M, Muller T, *et al*. Epstein-Barr virus-associated Hodgkin's lymphoma and legionella pneumophila infection complicating treatment of juvenile rheumatoid arthritis with methotrexate and cyclosporine A. *Hum Pathol* 2000;**31**:253-5.
- 5 **Londino AV Jr**, Blatt J, Knisely AS. Hodgkin's disease in a patient with juvenile rheumatoid arthritis taking weekly low dose methotrexate. *J Rheumatol* 1998;**25**:1245-6.
- 6 **Cotterill SJ**, Parker L, Malcolm AJ, *et al*. Incidence and survival for cancer in children and young adults in the North of England, 1968-1995: a report from the Northern Region Young Persons' Malignant Disease Registry. *Br J Cancer* 2000;**83**:397-403.
- 7 **Srivastava T**, Zwick D, Rothberg P, Warady B. Posttransplant lymphoproliferative disorder in paediatric renal transplantation. *Pediatr Nephrol* 1999;**13**:748-54.
- 8 **Kamel O**, van de Rijn M, LeBrun D, *et al*. Lymphoid neoplasms in patients with rheumatoid arthritis and dermatomyositis. *Hum Pathol* 1994;**25**:638-43.
- 9 **Salloum E**, Cooper DL, Howe G, *et al*. Spontaneous regression of lymphoproliferative disorders in patients treated with methotrexate for rheumatoid arthritis and other rheumatic diseases. *J Clin Oncol* 1996;**14**:1943-9.
- 10 **Tesch H**, Diehl V. Hodgkin's disease: from basic science to clinical application. *Leukemia* 1996;**10**(suppl 2):S74-7.
- 11 **Giannini EH**, Brewer EJ, Kuzmina N, *et al*. Methotrexate in resistant juvenile rheumatoid arthritis—results of the USA-USSR double-blind, placebo-controlled trial. *N Engl J Med* 1992;**326**:1043-9.

Want full text but don't have  
a subscription?

Pay per view

For just \$8 you can purchase the full text of individual articles using our secure online ordering service. You will have access to the full text of the relevant article for 48 hours during which time you may download and print the pdf file for personal use.

[www.archdischild.com](http://www.archdischild.com)